



Efficacy and Economics of Selected Insecticides against Pod Borer [*Helicoverpa armigera* (Hubner)] on Cowpea [*Vigna unguiculata* (L.)Walp.]

Gundlapalli Ranjith^{a*} and Ashwani Kumar^a

^a Department of Entomology, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92359

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103245>

Original Research Article

Received: 12/05/2023

Accepted: 10/07/2023

Published: 17/07/2023

ABSTRACT

A field trial was conducted at CRF, SHUATS, Naini, Prayagraj during *Kharif* season 2022. Eight treatments were evaluated against *Helicoverpa armigera* i.e. of Chlorantraniliprole 18.5 SC, Emamectin benzoate 5 SG, Flubendiamide 480 SC, Spinosad 45 SC, Thiamethoxam 25 WG, Lambda cyhalothrin5 EC, Indoxacarb 14.5 SC and untreated Control. Results revealed that, among the different treatments lowest larval population of cowpea pod borer was recorded in Chlorantraniliprole 18.5 SC (1.323). Lambda cyhalothrin5 EC (1.478) was found to be the next best treatment followed by Emamectin benzoate 5 SG (1.589), Indoxacarb 14.5 SC (1.734) and Flubendiamide 480 SC (1.922). The least effective treatments were Spinosad 45 SC (2.089) where as Thiamethoxam 25 WG (2.200) was found to be least effective. The plot treated with Chlorantraniliprole 18.5 SC show highest yield (19.46 q/ha) followed Lambda cyhalothrin5 EC (17.82 q/ha) the best and most economical treatment was Chlorantraniliprole 18.5 SC (1: 3.89).

*Corresponding author: E-mail: ranjithcobabs@gmail.com;

Among the treatments studied, the best and most economical treatment was Chlorantraniliprole 18.5 SC (1: 3.89) followed by Lambda cyhalothrin 5 EC (1:3.67), Emamectin benzoate 5 SG (1:3.41), Indoxacarb 14.5 SC (1:2.95), Flubendiamide 480 SC (1:2.56), Spinosad 45 SC (1:2.27) and Thiamethoxam 25 WG (1:2.17) as compared to untreated control plot (1:1.60).

Keywords: Cost-benefit ratio; cowpea; efficacy; *H. armigera*; insecticides; larval population.

1. INTRODUCTION

Cowpea [*Vigna unguiculata* (L.) Walp.] is an important grain legume mainly grown in tropical and subtropical regions for vegetables, grains, and fodder. Cowpea belongs to the family Fabaceae and sub-family Faboidea, and it is a self-pollinating crop with low and narrow genetic diversity, making it susceptible to various environmental factors" [1]. "It requires very few inputs, as the plants root nodules are able to fix atmospheric nitrogen, making it a valuable crop for resource poor farmers and well suited to intercropping with other crops" [2]. "Cowpea is known as vegetable meat due to high amount of protein in the grain with better biological value on dry weight basis. The grain contains 26.61% protein, 3.99% lipid, 56.24% carbohydrates, 8.60% moisture, 3.84 % ash, 1.38% crude fibre, 1.51 % gross energy, and 54.85% nitrogen free extract" [3]. "There are about 21 insect pests of different groups have been recorded damaging the cowpea crop from germination to maturity. The avoidable losses in yield due to insect pests have been recorded in the range of 66 to 100 per cent in cowpea" [4]. "The podborer complex posing serious threat to cowpea cultivation includes Bean pod borer (*Maruca vitrata*) (Fabricius), Blue butterfly (*Lampides boeticus*) (L.), Cow pea pod borer (*Helicoverpa armigera*) (Hubner), Pulse pod borer moth (*Etiella zinckenella*), Field bean pos borer (*Adisura atkinsoni*) and Red gram pod fly (*Exelastis atomosa*) (Walsingham)" [5].

"*Helicoverpa armigera* (Hubner) is a major pest that attacks wide range of different vegetable crops and causes serious economic damage and yield loss. The caterpillars of gram pod borer not only defoliate the leaves but also feed on seeds. While feeding on the developing seeds it's nearly half of the anterior body portion remains inside while the rest of the half portion remains hanging outside. A single larva may destroy 30-40 pods before it reaches maturity. The caterpillars feed on their fellows if suitable vegetation is not available i.e., cannibalism. They pupate in the soil. The pod damage due to *Helicoverpa armigera* on cowpea

crop could increase up to 100% in India. The best way to overcome this damage is to destroy the pest at its initial stage of the life cycle" [6].

Despite all the drawbacks that come with chemical control of pests, there are still many reasons that can persuade farmers to still choose to use them. Regardless of several control strategies that were found effective in managing this pest the chemical control plays a vital role because of quick action, readily available and very easy to use unlike alternative methods, such as biological control and other similar methods which can take a long while to plan and often don't have an immediate effect on pests. Therefore, keeping in view the above facts the present investigation was carried out with the aim to develop a new management strategy for control of pest at farmer's farm economically.

2. MATERIALS AND METHODS

The experiment was conducted at the experimental research plot of the Department of Entomology, Central Research Farm, Sam Higginbottom University of Agriculture Technology And Sciences, during the *Kharif* season of 2022, in a Randomized Block Design with eight treatments replicated three times using variety Kashi kanchan seeds in a plot size of 2m×1m at a spacing of 30 cm × 15cm with a recommended package of practices excluding plant protection. The soil of the experimental site was well drained and medium high. Research field situated at 25°27" North latitude 80°05" East longitudes and at an altitude of 98meter above sea level. The maximum temperature reaches upto 47°C in summer and drops down to 2°C in winter.

Pest population was estimated by observing five plants selected randomly from each treatment for presence of egg masses and larvae at one day prior to insecticide application and at 3rd, 7th and 14th days after each application. The larval population over control against pod borer (*H. armigera*) was calculated by considering the mean of three observations recorded at 3rd, 7th and 14th days after first and second spraying.

The healthy marketable yield obtained from different treatments was collected separately and weighed. The cost of insecticides used in this experiment was recorded during *Kharif* season. The cost of botanicals used was obtained from nearby market. The total cost of plant protection consisted of cost of treatments, sprayer rent and labour charges for the spray. There are two sprays throughout the research period and the overall plant protection expenses was calculated. Total income was realized by multiplying the total yield per hectare by the prevailing market price, while the net benefit is obtained by subtracting the total cost of plant protection from total income. Benefit over the control for each sprayed treatment was obtained by subtracting the income of the control treatment from that of each sprayed treatment” [4]. The C:B ratio was calculated by formula:

$$\text{Gross return} = \text{Marketable yield} \times \text{Market price}$$

$$\text{Cost: Benefit Ratio} = \text{Gross return} / \text{Total cost of cultivation}$$

3. RESULTS AND DISCUSSION

The results of the field trail with insecticides revealed that among the insecticides treated against gram pod borer after first spray Chlorantraniliprole 18.5% SC (1.645) was found significantly superior in reducing the larval population which was followed by Lambda cyhalothrin5 EC (1.755) was found to be the next best treatment followed Emamectin benzoate 5

SG (1.845), Indoxacarb 14.5 SC (2.000), Flubendiamide 480 SC (2.133), Spinosad 45 SC (2.244) and Thiamethoxam25 WG (2.333). After second spray, all the insecticides were found superior over untreated control. The overall mean analysis showed that Chlorantraniliprole 18.5% SC (1.323) and Lambda cyhalothrin5 EC (1.478) were significantly superior than other treatments followed by Emamectin benzoate 5 SG (1.589), Indoxacarb 14.5 SC (1.734), Flubendiamide 480 SC (1.922), Spinosad 45 SC (2.089) whereas Thiamethoxam25 WG (2.200). The treatments were found to be statistically at par with each other.

“Chlorantraniliprole was found to reduce the pod borer larval population to a tune of (1.323). Regarding the yield of cowpea, Chlorantraniliprole registered significantly higher yield (19.46 q/ha) and B:C ratio of 3.89 (Table 1.). The present finding is in line with observation on field application of Chlorantraniliprole 18.5% EC 0.5ml/l against cowpea pod borer and recorded lowest larval population in cowpea” Sreekanth et al., [7], Iqbal et al., [8], Chitralkha et al., [9] and Patil et al., [10].

Lambda cyhalothrin5 EC, Emamectin benzoate 5 SG, Indoxacarb 14.5 SC, Flubendiamide 480 SC, Spinosad 45 SC along with Thiamethoxam25 WG was found effective in reducing larval population Lekha et al., [11], Moosan and Kumar [12], Game et al., [13] and Dinesh et al., [14].

Table 1. Efficacy of certain insecticides against pod borer [*Helicoverpa armigera* (Hubner)] on cowpea [*Vigna unguiculata* (L.)Walp.] overall mean

S.No.	Treatments	Larval population of <i>H. armigera</i> /five plants							Overall Yield mean (q/ha)	C:B ratio	
		First spray				Second spray					
		1DBS	3DAS	7DAS	14DAS	3DAS	7DAS	14DAS			
T1	Chlorantraniliprole 18.5SC	2.467	1.800	1.467	1.667	1.400	0.733	0.867	1.323	19.46	1:3.89
T2	Emamectin benzoate5 SG	2.533	2.000	1.667	1.867	1.667	1.133	1.200	1.589	16.20	1:3.41
T3	Flubendiamide480 SC	2.400	2.267	1.933	2.200	2.000	1.467	1.667	1.922	12.45	1:2.56
T4	Spinosad 45 SC	2.600	2.333	2.000	2.400	2.200	1.733	1.867	2.089	11.40	1:2.27
T5	Thiamethoxam25 WG	2.667	2.400	2.133	2.467	2.333	1.867	2.000	2.200	10.24	1:2.17
T6	Lambda cyhalothrin5EC	2.333	1.933	1.533	1.800	1.533	0.933	1.133	1.478	17.82	1:3.67
T7	Indoxacarb 14.5 SC	2.467	2.200	1.867	1.933	1.800	1.200	1.400	1.734	14.60	1:2.95
T8	Control	2.400	2.533	2.600	2.667	2.733	2.800	2.933	2.711	7.10	1:1.60
	F-test	NS	S	S	S	S	S	S	S		
	S. Ed (±)	-----	0.089	0.057	0.077	0.067	0.057	0.063	0.024		
	C.D. (P = 0.5)	-----	0.189	0.124	0.177	0.143	0.130	0.140	0.063		

Dadas et al. [15] who reported maximum control of gram pod borer with the application of Chlorantraniliprole followed by Lambda cyhalothrin 5 EC. Gudipati and Mondal [16] who reported that Chlorantraniliprole gave the best performance with minimum (1.323) number of larvae followed by Lambda cyhalothrin 5 EC (1.478 larvae), which were at par with each other. The efficacy of newer insecticide for the management of *Helicoverpa armigera*, most effective was Emamectin benzoate 5 SG followed by Indoxacarb 14.5 SC reported by Fakhouri et al., [17] and Meena et al. [18].

Maximum cost benefit ratio (1:3.89) was obtained in Chlorantraniliprole 18.5% SC which was supported by Iqbal et al. [8] who reported that Chlorantraniliprole recorded the highest yield. The next cost benefit ratio obtained in the treatment Lambda cyhalothrin 5 EC (1:3.67) was supported by Jagtap and Kumar [19]. The least cost benefit ratio was observed in Thiamethoxam 25 WG (1:2.17) similar findings made by Nithish et al. [20] but superior as compared to control plot (1:1.60) [21].

4. CONCLUSION

From the analysis of the present findings, it can be concluded that Chlorantraniliprole 18.5 SC is more effective in controlling population of cowpea pod borer followed by Lambda cyhalothrin 5 EC was found to be the next best treatment followed by Emamectin benzoate 5 SG, Indoxacarb 14.5 SC, Flubendiamide 480 SC, Spinosad 45 SC and Thiamethoxam 25 WG in managing *Helicoverpa armigera*. Among the treatments studied, Chlorantraniliprole 18.5 SC gave the highest cost benefit ratio (1:3.89) and marketing yield (19.46 q/ha) followed by Lambda cyhalothrin 5 EC (1:3.67 and 17.82 q/ha), Emamectin benzoate 5 SG (1:3.41 and 16.20 q/ha), Indoxacarb 14.5 SC (1:2.95 and 14.60 q/ha), Flubendiamide 480 SC (1:2.56 and 12.45 q/ha), Spinosad 45 SC (1:2.27 and 11.40 q/ha) and Thiamethoxam 25 WG (1:2.17 and 10.24 q/ha) as such more trials are required in future to validate the findings.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Horn LN, Selma NN, Ueitele I. Cowpea Production Challenges and Contribution

2. to Livelihood in Sub-Saharan Region. Agricultural Sciences. 2022;13:25-32.
2. Lakshmikanth R, Kumar A. Comparative efficacy of selected chemicals and Biopesticides against gram pod borer [*Helicoverpa armigera* (Hubner)] (Lepidoptera: Noctuidae) on cowpea [*Vigna unguiculata* (L.) Walp.]. Journal of Pharmacognosy and Phytochemistry. 2018;7(3):3307-3309.
3. Oyewale RO, Bamaiyi LJ. Management of cowpea insect pests. Scholars Academic Journal of Biosciences. 2013; 1(5):217-226.
4. Ekshinge SN, Kumar A. Efficacy of selected insecticides against fall army worm [*Spodoptera frugiperda* (JE Smith)] on maize (*Zea mays* L.). The Pharma Innovation Journal. 2023;12(5): 4470-4473.
5. Kumar K, Tayde AR. Comparative efficacy of selected botanicals and Novaluran against spotted pod borer [*Maruca vitrata* (Fabricius)] (Lepidoptera:Crambidae) on cowpea [*Vigna unguiculata* (L.) Walp.]. Journal of Pharmacognosy and Phytochemistry. 2017;6(5):540-542.
6. Ahmed K, Awan MS. Integrated management of insect pests of chickpea *Cicer arietinum* (L. Walp) in South Asian Countries: Present status and future strategies– A review. Pakistan Journal of Zoology. 2013;45(4):1125-1145.
7. Sreekanth M, Lakshmi MSM, Rao YK. Bio-efficacy and economics of certain new insecticides against gram pod borer, *Helicoverpa armigera* (Hubner) infesting pigeonpea (*Cajanus cajan* L.). International Journal of Plant, Animal and Environmental Sciences. 2014;4(1): 11-15.
8. Iqbal J, Farooq SU, Jamil M, Khan HAA, Younis M. Relative efficacy of selective insecticides against gram pod borer (*Helicoverpa armigera* H.) of chickpea. Myopathy. 2014;12(2):119-122.
9. Chitralkha, Yadav GS, Verma T. Efficacy of insecticides against *Helicoverpa armigera* on chickpea, Journal of Entomology and Zoology Studies. 2018;6(3):1058-1061.
10. Patil PV, Pawar SA, Kadu RV, Pawar DB. Bio-efficacy of newer insecticides, botanicals and microbial against tomato fruit borer *Helicoverpa armigera* (Hubner) infesting tomato. Journal of Entomology

- and Zoology Studies. 2018;6(5):2006-2011.
11. Lekha, Ameta OP, Swami H. Evaluation of new generation pesticides to control pod borer, *Helicoverpa armigera* (Hubner) infesting chickpea. Legume Research- An International Journal. 2017;40(2):384-387.
 12. Moosan M, Kumar A. Efficacy and economics of different insecticides against gram pod borer, *Helicoverpa armigera* (Hubner).The Pharma Innovation Journal. 2022;11(7):4471-4474.
 13. Game L, Nagar S, Simon S. Comparative efficacy of selected chemical insecticides and neem products against tomato fruit borer [*Helicoverpa armigera* (Hubner)] in Allahabad. Journal of Pharmacognosy and Phytochemistry. 2018;7(4):2215-2218.
 14. Dinesh K, Singh RB, Kavita K, Choudhary RS. Relative efficacy of newer insecticides and biopesticides against *Helicoverpa armigera* (hub.) in chickpea, Journal of Entomology and Zoology Studies. 2017;5(3):455-462.
 15. Dadas SM, Gosalwad SS, Patil SK. Efficacy of different newer insecticides against pigeon pea pod borers, Journal of Entomology and Zoology Studies. 2019;7(5):784- 791.
 16. Gudipati N, Mondal GS. Efficacy of some insecticides against gram pod borer *Helicoverpa armigera* on Chickpea. Indian Journal of Entomology. 2020;82(3):515- 518.
 17. El Fakhouri K, Boulamtat R, Sabraoui A, El Bouhssini M. The chickpea pod borer, *Helicoverpa armigera* (Hübner): Yield loss estimation and biorational insecticide assessment in Morocco. Agronomy. 2022;12(12):3017.
 18. Meena RK, Naqvi AR, Meena DS, Shivbhagvan. Evaluation of bio-pesticides and indoxacarb against gram pod borer on chickpea. Journal of Entomology and Zoology Studies. 2018; 6(2):2208-2212.
 19. Jagtap PM, Kumar A. Field efficacy and economics of different chemicals and biopesticides against pod borer, *Helicoverpa armigera* (Hubner) on chickpea during rabi season. The Pharma Innovation Journal. 2022;11(7): 3970-3975.
 20. Nithish A, Joshi B, Jayaram CS, Kariyanna B. Bio-efficacy of some newer insecticides against gram pod borer (*Helicoverpa armigera*) in pigeon pea under field conditions. Research Journal of Agricultural Sciences. 2015;6(3):563-566.
 21. Gayathri L, Kumar A. Field efficacy of certain insecticides against pod borer, *Helicoverpa armigera* (Hubner) on chick pea in Prayagraj. Journal of Entomology and Zoology Studies. 2019;9(3):280-283.

© 2023 Ranjith and Kumar; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/103245>