

Facial Teratoma in the Newborn: Diagnosis and Prognosis

**Nadia Ben Jamaa¹, Radhouane Achour^{2*}, Feirouz Ayari³, Nadia Lamari²,
Samia Kacem³, Khaled Neji² and Aida Masmoudi¹**

¹*Department of Fetopathology, Center of Maternity and Neonatology of Tunis, University Tunis El Manar, Tunisia.*

²*Department of Emergency, Center of Maternity and Neonatology of Tunis, University Tunis El Manar, Tunisia.*

³*Neonatal Intensive Care Unit, Center of Maternity and Neonatology of Tunis, University Tunis El Manar, Tunisia.*

Authors' contributions

This work was carried out in collaboration between all the authors. Authors NBJ and AM conducted the collection and development of the observation. Authors RA and NBJ involved in writing the article.

Authors SK, KN and AM involved in collecting and reviewing the bibliography and proposing recommendations in writing the article. Authors FA, NL and RA involved in writing the article and translating it into English. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2017/33890

Editor(s):

- (1) Andrzej Kloczkowski, Battelle Center for Mathematical Medicine, The Research Institute at Nationwide Children's Hospital, USA and Department of Pediatrics, The Ohio State University College of Medicine, USA.
(2) George Perry, Dean and Professor of Biology, University of Texas at San Antonio, USA.

Reviewers:

- (1) Wagih Mommtaz Ghannam, Mansoura University, Egypt.
(2) Alicia García Falgueras, The Official College of Psychologists, Madrid, Spain.
(3) Einar Arnbjörnsson, Lund University, Sweden

Complete Peer review History: <http://www.sciencedomain.org/review-history/19346>

Case Report

Received 2nd May 2017
Accepted 31st May 2017
Published 5th June 2017

ABSTRACT

Facial teratoma is an extremely rare germinal tumor in newborn babies. It is often diagnosed antenatally by means of ultrasound and even MRI. The prognosis of this tumor depends mainly on the size and the location of the lesion (eye, respiratory and digestive tracts). Multidisciplinary examination is needed to ensure the newborn's survival. Pre-operative conditioning is needed to avoid complications and carcinogenic risks. The aim of this article is to Specify the anatomical features of facial teratoma and consider the interests of foetopathology review.

*Corresponding author: E-mail: radhouane.a@live.com;

We report a male baby, prematurely born at 28 WA dead at hour 23. Physical examination reveals a frontal mass on the left side of the face plus a facial dysmorphia. Dissection shows hyperplasia of lungs, hypertrophied liver and a splenomegaly. Histological examination reveals neural tissue associated with areas of cartilage and hair; hence the diagnosis of congenital teratoma of the face. Antenatal diagnosis of congenital teratoma of the face is very crucial allowing a multidisciplinary care involving obstetricians, neonatologists, surgeons and anesthesiologists.

Keywords: Facial teratoma; diagnosis; antenatal; prognosis; foeto pathology.

1. INTRODUCTION

Facial teratoma is an extremely rare germinal tumor in newborn babies. It is often diagnosed antenatally by means of ultrasound and even MRI [1,2,3].

The prognosis of this tumor depends mainly on size and location of the lesion (eye, respiratory and digestive tracts). Multidisciplinary examination is needed to ensure the newborn's survival. Pre-intervention conditioning is needed to avoid complications and carcinogenic risks [4,5,6,7].

1.1 Aim

Specify the anatomical features of facial teratoma and consider the interests of foetopathology review.

2. CASE REPORT

Male baby, prematurely born at 28 weeks, delivered by cesarian section for chorioamnionitis, Death at 23 hours postpartum. This is a non-consanguineous marriage, mother aged 36 gravida 1, para 1, no medical history, absence of medication of the mother or the father and no accident during pregnancy. Pregnancy very poorly followed, which did not allow to diagnose the disease at an early term of pregnancy. Physical examination reveals a frontal mass on the left side of the face plus a facial dysmorphia (Fig. 1). Dissection shows hyperplasia of lungs, hypertrophied liver, a splenomegaly and a frontal tumor about 4 inches in diameter, soft consistency (Fig. 2). Histologic examination reveals Monodermal teratoma, neural tissue associated with areas of cartilage and hair; Without histological signs of malignancy. Hence the diagnosis of congenital teratoma of the face.

3. DISCUSSION

Teratomas are germinal tumors showing tissue from all 3 germ layers which are: Endoderm,

mesoderm and ectoderm. The most frequent type of teratoma in the newborn is sacrococcygeal teratoma which accounts for 80% of cases. 2% to 9% of teratomas are located in the head and neck [2,6].



Fig. 1. Retro-orbital tumor, Cranio-facial dysmorphia: Flat nose, thin lips, retrognathia, large low setted ears

Antenatal diagnosis is possible by ultrasound, which allows us to plan specialized management of the child to be born; but in our case it was a poorly followed pregnancy [3].

The prognosis of this tumor depends mainly on size and location of the lesion.

Teratomas can be divided in 3 categories [8,9,10]:

- Mature (benign).
- Immature (malignant).
- Monodermal: This type of tumor contains differentiated cells producing mature tissue in an inappropriate location, often different from surrounding tissues. This is the case of the teratoma diagnosed in our newborn child.

Facial location is a differential diagnosis to encephalocele which only contains brain tissue.

Prenatal Karyotype and searching for associated abnormalities is obligatory in all teratomas [2,11].




Fig. 2. Hyperplasia of lungs (MAKP III histologic examination); Hepatomegaly and splenomegaly

Besides the delivery should include cesarean section with ex utero intrapartum resection of the tumor mass, which may be performed on a placental support operation to increase post natal survival [3,4,5].

A systematic review reported that 3D ultrasound and MRI may enhance the accuracy of the antenatal diagnosis and may ease the selection of newborns requiring treatment [1,9,6].

Multidisciplinary examination is needed to ensure the newborn's survival. Pre-operative conditioning is needed to avoid complications and carcinogenic risks [7].

4. CONCLUSION

Teratomas represent less than 7% of all infantile tumors [12]. A review of the literature shows that women are twice as affected as males [1,12]. The face is an exceptional site. A teratoma is a congenital tumor from the 3 layers of germ cells [8,9,12]. The cases described in the literature are often cytological benign [1,11,12]. In some cases of large tumors, serious complications can occur, even neonatal death.

After a review of the literature [1-12], the diagnosis of these rare cases of teratomas is possible in antenatal by ultrasound. The prognosis depends essentially on the histological nature, the site and size of the tumor. Generally favorable after surgical treatment and multidisciplinary care; but can be severe with neonatal death in other cases.

CONSENT

Parents' informed consent for eventually publish the information for Scientific aims was obtained.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the ethics

committee of the Maternity and Neonatology Center of Tunis (approval number /reference: 00018/17) and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

ACKNOWLEDGEMENTS

This work was supported by the Maternity and Neonatology Center, Faculty of Medicine of Tunis, Tunisia.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tonni G, De Felice C, Centini G, Ginanneschi C. Cervical and oral teratoma in the fetus, a systematic review of etiology, pathology, diagnosis, treatment and prognosis. *Arch Gynecol Obstet.* 2010;285:355-61.
2. Tian-Ni K, Yi-Hung S, Nga-San H, Hsing CT, Cin-Chuan L, Lin MYS, Chen SH. Multidisciplinary examination for prenatal diagnosis of posterior cervical teratoma in early second trimester. *Taiwanese Journal of Obstetrics and Gynecology.* 2013;270-272.
3. Courtier J, Poder I, Wang ZJ, Westfalen AC, Yeh BM, Coakley FV. Fetal tracheolaryngeal airway obstruction: Prenatal evaluation by sonography and MRI. *Pediatr Radiol.* 2010;40:1800-5.
4. Lakhoo K. Neonatal teratomas. *Early Hum Dev.* 2010;86:643-7.
5. Le Saux A, Poli Merol ML, Zachar D, Josset P, Pezzettigotta S, Doco Fenzy M, Guyot E, Francois-Fiquet C. Cleft palate associated with cervico-facial teratoma: Report of two cases and review of the

- literature. *Int J Pediatr Otorhinolaryngol.* 2012;76(8):1225-7.
6. Azizkhan RG, Haase GM, Applebaum H, Dillon PW, Coran AG, King PA. Diagnosis, management, and outcome of cervico facial teratomas in neonates: A children's cancer group study. *J Pediatr Surg.* 1995;30:312–6.
 7. Aubry MC, Aubry JP. Diagnostic anténatal des malformations et tumeurs cervico-faciales. *Rev Prat.* 1991;41:16-20.
 8. Fernando Pellerano, Elvis Guillermo, Gloreley Garrido, Pedro Berges. Congenital orbital teratoma. *Ocul Oncol Pathol.* 2017;3(1):11–16.
 9. Pablo Padilla Iserte, Amparo Sanroma Pérez, Blanca Ferri Folch, Juan Rubio Moll, Vicente Diago Almela, Alfredo Perales-Marín. Ultrasound evaluation of congenital cervical teratoma and therapeutic management (Ex Utero Intrapartum Treatment). *Case Rep Obstet Gynecol.* 2012;597489. DOI: 10.1155/2012/597489
 10. Rafał Zieliński, Maria Respondek-Liberska. The role of prenatal ultrasound assessment in management of fetal cervicofacial tumors. *Arch Med Sci.* 2016;12(4):850–855.
 11. Sahar N Saleem. Fetal MRI: An approach to practice: A review. *J Adv Res.* 2014;5(5):507–523.
 12. Huge orbital teratoma with intracranial extension: A case report. *Journal of Pediatric Surgery.* 2010;45:E27–E31.

© 2017 Jamaa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/19346>*