



Common Fixed Point Theorems for (\mathfrak{F}, ϕ) - Contractive Mappings on C^* -algebra Valued \mathfrak{B} -metric Spaces

K. Maheshwaran ^{a*}, R. Jahir Hussain ^a
and Nesrin Manav Tatar ^b

^aPG & Research Department of Mathematics, Jamal Mohamed College (Autonomous) (Affiliated to Bharathidasan University), Tiruchirappalli-620020, Tamilnadu, India.

^bDepartment of Mathematics, Science and Art Faculty, Erzincan Binali Yildirim University, Erzincan, Turkey.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJOM/2023/v19i9713

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103141>

Received: 12/05/2023

Accepted: 14/07/2023

Published: 21/07/2023

Original Research Article

Abstract

This paper, we extends the new concept of common fixed point theorems in C^* -algebra valued \mathfrak{b} -metric space ($AVBMS$) via (\mathfrak{F}, ϕ) - contractive mappings. Investigated are the common fixed points criteria for existence and uniqueness. Additionally, provide an illustrate an example.

Keywords: C^* -algebra valued; \mathfrak{b} -metric spaces; common fixed point.

2010 Mathematics Subject Classification: 47H10, 54H25, 54H20.

*Corresponding author: E-mail: mahesksamy@gmail.com;

Asian Res. J. Math., vol. 19, no. 9, pp. 218-225, 2023

1 Introduction

In 2014, Ma et al. [1] proposed the idea of C^* -algebra valued metric space (AVMS) and investigated certain fixed point theorems for self-mapping under various contractive circumstances. Furthermore, the concept of C^* -(AVMS) is generalized to that of C^* -(AVBMS), where \mathfrak{B} is an element of C^* -algebra greater than 1 and the triangle inequality is altered into $D_b(\varsigma, \eta) \leq \tau(D_b(\varsigma, \mathfrak{z}) + D_b(\mathfrak{z}, \eta))$. Theorems for self-map with contractive condition are then established using various fixed point theorems [2]. Besides, Alsulami et al. [3] the classic Banach fixed point theorems have been examined for their fixed point outcomes can be used to produce C^* -(AVMS), C^* -(AVBMS) in fixed point theory. [4, 5]. We recall the concept of C^* -(AVBMS). Since C^* -algebra is an important topic in functional analysis, operator theory and algebras which is important in non-commutative geometry and theoretical physics, such as quantum mechanics and string theory [6]. During this article, we indicate \mathfrak{A} as an unitary C^* -algebra and $\mathfrak{A}_h = \{a \in \mathfrak{A} : a = a^*\}$. Especially, an aspect $a \in \mathfrak{A}$ is a positive factor, if $a = a^*$ and $\sigma(a) \subseteq \mathcal{R}^+$, where $\sigma(a)$ is the spectrum of a . There is a natural partial order in \mathfrak{A}_h placed by $a \leq b$ iff $\theta \leq (b - a)$, where θ implies the zero factor in \mathfrak{A} . Then, let \mathfrak{A}^+ and \mathfrak{A}' represent the set $\{a \in \mathfrak{A} : \theta \leq a\}$ and the set $\{a \in \mathfrak{A} : ab = ba, \forall b \in \mathfrak{A}\}$, respectively and $|a| = (a^*a)^{\frac{1}{2}}$.

Definition 1.1. [1] Let χ be a non-empty set and $\tau \in \mathfrak{A}$ such that $\tau \geq I$. Suppose that the mapping $D_b : \chi \times \chi \rightarrow \mathfrak{A}$ is held, the following constraints.

- (i) $\theta \leq D_b(\varsigma, \eta)$ and $D_b(\varsigma, \eta) = \theta$ iff $(\varsigma = \eta)$;
- (ii) $D_b(\varsigma, \eta) = D_b(\eta, \varsigma)$;
- (iii) $D_b(\varsigma, \eta) \leq \tau(D_b(\varsigma, \mathfrak{z}) + D_b(\mathfrak{z}, \eta)) \forall \varsigma, \eta, \mathfrak{z} \in \chi$.

Then, D_b is called C^* -(AVBM) on χ and $(\chi, \mathfrak{A}, D_b)$ is called C^* -(AVBMS).

Definition 1.2. [1]. Let $(\chi, \mathfrak{A}, D_b)$ be C^* -(AVBMS). Imagine that $\{\varsigma_n\}$ is an order in χ and $\varsigma \in \chi$. If at all $\epsilon > \theta, \exists \mathcal{N} \ni \forall n > \mathcal{N}, \|D_b(\varsigma_n, \tau)\| \leq \epsilon$ then $\{\varsigma_n\}$ is allegedly convergent with respect to \mathfrak{A} , and $\{\varsigma_n\}$ converges to ς , i.e., we use $\lim_{n \rightarrow \infty} \varsigma_n = \varsigma$. If at all $\epsilon > \theta, \exists \mathcal{N} \ni \forall n, m > \mathcal{N}, \|D_b(\varsigma_n, \varsigma_m)\| \leq \epsilon$, then $\{\varsigma_n\}$ is referred to as a Cauchy sequence in χ . $(\chi, \mathfrak{A}, D_b)$ is referred to as a complete C^* -(AVBMS) if each Cauchy sequence is convergent in χ .

Remark 1.1. [7] Let \mathfrak{A} be a C^* -algebra and assume that ϕ is a linear functional on \mathfrak{A} . Define $\phi^*(a) = \phi(a^*) \forall a \in \mathfrak{A}$. Then, ϕ^* likewise has a linear function \mathfrak{A} . And the function ϕ is known as a self-adjoint if $\phi^* = \phi$.

A linear function ϕ on \mathfrak{A} is called positive if $\phi(a^*a) \geq 0$ for all $a \in \mathfrak{A}$. We indicate the positivity of ϕ by $\phi \geq \theta$. For two self-adjoint linear function ϕ_1, ϕ_2 , There are $(\phi_2 - \phi_1) \geq \theta$ when $\phi_2 \geq \phi_1$.

Definition 1.3. [7] If $\phi : \mathfrak{A} \rightarrow \mathfrak{B}$ is a linear mapping in C^* -algebra, It is said to be positive if $\phi(\mathfrak{A}^+) \subseteq \phi(\mathfrak{B}^+)$. When this occurs, $\phi(\mathfrak{A}_h) \subseteq \phi(\mathfrak{B}_h)$, and the map of restriction $\phi : \mathfrak{A}_h \rightarrow \mathfrak{B}_h$ is increasing. if $\mathfrak{B} = \mathfrak{A}$ the positive linear map is thus referred to as positive functional, and it meets the following propositions 1.1 and 1.2.

Proposition 1.1. [7] Let \mathfrak{A} be a C^* -algebra with 1, thereafter, a positive functional is bounded and $\phi(1) = \|\phi\|$.

Proposition 1.2. [7] Let \mathfrak{A} be a C^* -algebra with 1 and let ϕ be a bounded linear functional on $\mathfrak{A} \ni \phi(a) = \|\phi\| \|a\|$. There exists positive component $a \in \mathfrak{A} \ni \phi$ is a positive functional.

Definition 1.4. [8] Let the non-decreasing function $\mathfrak{F} : \mathfrak{A}^+ \rightarrow \mathfrak{A}^+$ be a positive linear map that complies with the following restrictions:

- (i) \mathfrak{F} is continuous;
- (ii) $\mathfrak{F}(a) = \theta$ iff $a = \theta$;
- (iii) $\lim_{n \rightarrow \infty} \mathfrak{F}^n(a) = \theta$.

Definition 1.5. [8] Suppose that \mathfrak{A} and \mathfrak{B} are C^* -algebra. \mathfrak{A} mapping $\mathfrak{F} : \mathfrak{A} \rightarrow \mathfrak{B}$ is purported to be C^* -homomorphism if :

- (i) $\mathfrak{F}(a\varsigma + b\eta) = a\mathfrak{F}(\varsigma) + b\mathfrak{F}(\eta) \forall a, b \in C$ and $\varsigma, \eta \in \mathfrak{A}$;
- (ii) $\mathfrak{F}(\varsigma, \eta) = \mathfrak{F}(\varsigma)\mathfrak{F}(\eta) \forall \varsigma, \eta \in \mathfrak{A}$;
- (iii) $\mathfrak{F}(\varsigma^*) = \mathfrak{F}(\varsigma)^* \forall \varsigma \in \mathfrak{A}$;
- (iv) \mathfrak{F} maps the unit in \mathfrak{A} to the unit in \mathfrak{B} .

Definition 1.6. [9] Let \mathfrak{A} and \mathfrak{B} be a C^* -algebra space and let $\mathfrak{F} : \mathfrak{A} \rightarrow \mathfrak{B}$ be a homomorphism, then \mathfrak{F} is called an $*$ -isomorphism if it is one-to-one $*$ -isomorphism. We declare that C^* -algebra \mathfrak{A} is $*$ -isomorphism to a C^* -algebra \mathfrak{B} if \exists $*$ -isomorphism of \mathfrak{A} onto \mathfrak{B} .

property 1.1. [9] Let \mathfrak{A} and \mathfrak{B} be C^* -algebra space and $\mathfrak{F} : \mathfrak{A} \rightarrow \mathfrak{B}$ is a C^* -homomorphism $\forall \varsigma \in \mathfrak{A}$, there are $\sigma(\mathfrak{F}(\varsigma)) \subset \sigma(\varsigma)$ and $\|\mathfrak{F}(\varsigma)\| \leq \|\varsigma\|$.

Corollary 1.1. [10] Every C^* -homomorphism is bounded.

Corollary 1.2. [10] Suppose that \mathfrak{F} is C^* -isomorphism from \mathfrak{A} to \mathfrak{B} , then $\sigma(\mathfrak{F}(\varsigma)) \subset \sigma(\varsigma)$ and $\|\mathfrak{F}(\varsigma)\| \leq \|\varsigma\| \forall \varsigma \in \mathfrak{A}$.

Lemma 1.3. [10] Every $*$ -homomorphism is positive.

2 Main Results

Theorem 2.1. Let $(\chi, \mathfrak{A}, D_b)$ is a complete C^* -(AVBMS). Let $L, M : \chi \rightarrow \chi$ be a contractive mapping and

$$\mathfrak{F}(D_b(L\varsigma, M\eta)) \leq \mathfrak{F}(N(\varsigma, \eta)) - \phi(D_b(\varsigma, \eta)) \tag{2.1}$$

$$N(\varsigma, \eta) \leq \left[\begin{array}{c} \alpha D_b(\varsigma, \eta) + \frac{\beta[1+D_b(\varsigma, L\varsigma)]D_b(\eta, M\eta)}{1+D_b(\varsigma, \eta)} \\ +\gamma [D_b(\varsigma, L\varsigma) + D_b(\eta, M\eta)] + \delta [D_b(\varsigma, M\eta) + D_b(\eta, L\varsigma)] \end{array} \right]$$

For all $\varsigma, \eta \in \chi$, where $\tau \in \mathfrak{A}'_+$, $\alpha + \beta + \gamma + \delta \geq 0$ with $\tau\alpha + \beta + \gamma(\tau + 1) + \delta(\tau(\tau + 1)) < 1$. \mathfrak{F} and ϕ are $*$ -homomorphisms and the constraint $\mathfrak{F}(\mathbf{a}) \leq \phi(\mathbf{a})$. Then L and M have a unique common fixed point in χ .

Proof. Let $\varsigma_0 \in \chi$ and define $\varsigma_n = L\varsigma_{n-1}$, $\varsigma_{n+1} = M\varsigma_n$ we have

$$\begin{aligned} \mathfrak{F}(D_b(\varsigma_n, \varsigma_{n+1})) &= \mathfrak{F}(D_b(L\varsigma_{n-1}, M\varsigma_n)) \\ &\leq \mathfrak{F}(N(\varsigma_{n-1}, \varsigma_n)) - \phi(D_b(\varsigma_{n-1}, \varsigma_n)) \\ &= \mathfrak{F} \left(\begin{array}{c} \alpha D_b(\varsigma_{n-1}, \varsigma_n) + \frac{\beta[1+D_b(\varsigma_{n-1}, L\varsigma_{n-1})]D_b(\varsigma_n, M\varsigma_n)}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \\ +\gamma [D_b(\varsigma_{n-1}, L\varsigma_{n-1}) + D_b(\varsigma_n, M\varsigma_n)] \\ +\delta [D_b(\varsigma_{n-1}, M\varsigma_n) + D_b(\varsigma_n, L\varsigma_{n-1})] \end{array} \right) - \phi(D_b(\varsigma_{n-1}, \varsigma_n)) \\ &= \mathfrak{F}(\alpha)\mathfrak{F}(D_b(\varsigma_{n-1}, \varsigma_n)) + \frac{\mathfrak{F}(\beta)\mathfrak{F}[1+D_b(\varsigma_{n-1}, L\varsigma_{n-1})]D_b(\varsigma_n, M\varsigma_n)}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \\ &\quad + \mathfrak{F}(\gamma)\mathfrak{F}[D_b(\varsigma_{n-1}, L\varsigma_{n-1}) + D_b(\varsigma_n, M\varsigma_n)] \\ &\quad + \mathfrak{F}(\delta)\mathfrak{F}[D_b(\varsigma_{n-1}, M\varsigma_n) + D_b(\varsigma_n, L\varsigma_{n-1})] - \phi(D_b(\varsigma_{n-1}, \varsigma_n)). \end{aligned}$$

Therefore

$$\begin{aligned} \|\mathfrak{F}(D_b(\varsigma_n, \varsigma_{n+1}))\| &= \|\mathfrak{F}(D_b(L\varsigma_{n-1}, M\varsigma_n))\| \\ &\leq \|\mathfrak{F}(\alpha)\| \|\mathfrak{F}(D_b(\varsigma_{n-1}, \varsigma_n))\| + \|\mathfrak{F}(\beta)\| \left\| \frac{\mathfrak{F}[1+D_b(\varsigma_{n-1}, L\varsigma_{n-1})]D_b(\varsigma_n, M\varsigma_n)}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \right\| \\ &\quad + \|\mathfrak{F}(\gamma)\| \|\mathfrak{F}[D_b(\varsigma_{n-1}, L\varsigma_{n-1}) + D_b(\varsigma_n, M\varsigma_n)]\| \\ &\quad + \|\mathfrak{F}(\delta)\| \|\mathfrak{F}[D_b(\varsigma_{n-1}, M\varsigma_n) + D_b(\varsigma_n, L\varsigma_{n-1})]\| \\ &\quad - \|\phi(D_b(\varsigma_{n-1}, \varsigma_n))\| \rightarrow 0 \text{ as } n \rightarrow +\infty. \end{aligned}$$

Consider the fact that ϕ and \mathfrak{F} are strongly monotone functions. There are

$$\begin{aligned}
 D_b(\varsigma_n, \varsigma_{n+1}) &= D_b(L\varsigma_{n-1}, M\varsigma_n) \\
 &\leq \alpha D_b(\varsigma_{n-1}, \varsigma_n) + \frac{\beta[1+D_b(\varsigma_{n-1}, L\varsigma_{n-1})]D_b(\varsigma_n, M\varsigma_n)}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \\
 &+ \gamma [D_b(\varsigma_{n-1}, L\varsigma_{n-1}) + D_b(\varsigma_n, M\varsigma_n)] + \delta [D_b(\varsigma_{n-1}, M\varsigma_n) + D_b(\varsigma_n, L\varsigma_{n-1})] \\
 &= \alpha D_b(\varsigma_{n-1}, \varsigma_n) + \frac{\beta[1+D_b(\varsigma_{n-1}, \varsigma_n)]D_b(\varsigma_n, \varsigma_{n+1})}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \\
 &+ \gamma [D_b(\varsigma_{n-1}, \varsigma_n) + D_b(\varsigma_n, \varsigma_{n+1})] + \delta [D_b(\varsigma_{n-1}, \varsigma_{n+1}) + D_b(\varsigma_n, \varsigma_n)] \\
 &= (\alpha + \gamma)D_b(\varsigma_{n-1}, \varsigma_n) + (\beta + \gamma)D_b(\varsigma_n, \varsigma_{n+1}) + \delta (D_b(\varsigma_{n-1}, \varsigma_{n+1})) \\
 &\leq (\alpha + \gamma)D_b(\varsigma_{n-1}, \varsigma_n) + (\beta + \gamma)D_b(\varsigma_n, \varsigma_{n+1}) \\
 &+ \tau\delta (D_b(\varsigma_{n-1}, \varsigma_n) + D_b(\varsigma_n, \varsigma_{n+1})) \\
 &= (\alpha + \gamma + \tau\delta)D_b(\varsigma_{n-1}, \varsigma_n) + (\beta + \gamma + \tau\delta)D_b(\varsigma_n, \varsigma_{n+1}).
 \end{aligned}$$

This indicates that

$$\begin{aligned}
 D_b(\varsigma_n, \varsigma_{n+1}) &\leq \frac{\alpha + \gamma + \tau\delta}{\beta + \gamma + \tau\delta} D_b(\varsigma_n, \varsigma_{n-1}) \\
 D_b(\varsigma_n, \varsigma_{n+1}) &\leq h D_b(\varsigma_n, \varsigma_{n-1})
 \end{aligned} \tag{2.2}$$

where,

$$h = \frac{\alpha + \gamma + \tau\delta}{\beta + \gamma + \tau\delta} < 1.$$

As a result,

$$\| D_b(\varsigma_{n-1}, \varsigma_n) \| \| D_b(\varsigma_n, \varsigma_{n+1}) \| \leq \| h \| \| D_b(\varsigma_n, \varsigma_{n-1}) \| \rightarrow 0, \text{ as } n, m \rightarrow +\infty.$$

Let $n > m$

$$D_b(\varsigma_n, \varsigma_m) \leq \tau D_b(\varsigma_n, \varsigma_{n-1}) + \tau^2 D_b(\varsigma_{n-1}, \varsigma_{n-2}) + \dots + \tau^{n-m} D_b(\varsigma_{m-1}, \varsigma_m)$$

When using the theorem's constraint,

$$\begin{aligned}
 \mathfrak{F}(D_b(\varsigma_n, \varsigma_m)) &\leq \mathfrak{F}(\tau D_b(\varsigma_n, \varsigma_{n-1})) + \mathfrak{F}(\tau^2 D_b(\varsigma_{n-1}, \varsigma_{n-2})) + \dots + \mathfrak{F}(\tau^{n-m} D_b(\varsigma_{m-1}, \varsigma_m)) \\
 &\leq \mathfrak{F}(\tau) \mathfrak{F}(D_b(\varsigma_n, \varsigma_{n-1})) + \mathfrak{F}(\tau^2) \mathfrak{F}(D_b(\varsigma_{n-1}, \varsigma_{n-2})) + \dots \\
 &+ \mathfrak{F}(\tau^{n-m}) \mathfrak{F}(D_b(\varsigma_{m-1}, \varsigma_m)) \\
 &\leq \mathfrak{F}(\tau N(\varsigma_n, \varsigma_{n-1})) - \phi(\tau D_b(\varsigma_n, \varsigma_{n-1})) + \mathfrak{F}(\tau^2 N(\varsigma_{n-1}, \varsigma_{n-2})) \\
 &- \phi(\tau^2 D_b(\varsigma_{n-1}, \varsigma_{n-2})) + \dots + \mathfrak{F}(\tau^{n-m} N(\varsigma_{m-1}, \varsigma_m)) \\
 &- \phi(\tau^{n-m} D_b(\varsigma_{m-1}, \varsigma_m)) \\
 &= \mathfrak{F} \left(\begin{aligned} &\alpha D_b(\varsigma_n, \varsigma_{n-1}) + \frac{\beta[1+D_b(\varsigma_{n-1}, \varsigma_n)]D_b(\varsigma_n, \varsigma_{n+1})}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \\ &+ \gamma \tau [D_b(\varsigma_{n-1}, \varsigma_n) + D_b(\varsigma_n, \varsigma_{n+1})] \\ &+ \delta \tau [D_b(\varsigma_{n-1}, \varsigma_{n+1}) + D_b(\varsigma_n, \varsigma_n)] \end{aligned} \right) - \phi(\tau D_b(\varsigma_n, \varsigma_{n-1})) + \dots \\
 &+ \mathfrak{F} \left(\begin{aligned} &\alpha \tau^{n-m} D_b(\varsigma_{m-1}, \varsigma_m) + \frac{\beta \tau^{n-m} [1+D_b(\varsigma_m, \varsigma_{m-1})] D_b(\varsigma_{m-1}, \varsigma_{m-2})}{1+D_b(\varsigma_m, \varsigma_{m-1})} \\ &+ \gamma \tau^{n-m} [D_b(\varsigma_m, \varsigma_{m-1}) + D_b(\varsigma_{m-1}, \varsigma_{m-2})] \\ &+ \delta \tau^{n-m} [D_b(\varsigma_m, \varsigma_{m-2}) + D_b(\varsigma_{m-1}, \varsigma_{m-1})] \end{aligned} \right) \\
 &- \phi(\tau^{n-m} D_b(\varsigma_{m-1}, \varsigma_m)).
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \mathfrak{F}(D_b(\varsigma_n, \varsigma_m)) &= \left(\begin{aligned} &\mathfrak{F}(\alpha) \mathfrak{F}(\tau) \mathfrak{F}(D_b(\varsigma_n, \varsigma_{n-1})) \\ &+ \mathfrak{F}(\beta) \mathfrak{F}(\tau) \mathfrak{F} \left(\frac{[1+D_b(\varsigma_{n-1}, \varsigma_n)] D_b(\varsigma_n, \varsigma_{n+1})}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \right) \\ &+ \mathfrak{F}(\gamma) \mathfrak{F}(\tau) \mathfrak{F} [D_b(\varsigma_{n-1}, \varsigma_n) + D_b(\varsigma_n, \varsigma_{n+1})] \\ &+ \mathfrak{F}(\delta) \mathfrak{F}(\tau) \mathfrak{F} [D_b(\varsigma_{n-1}, \varsigma_{n+1}) + D_b(\varsigma_n, \varsigma_n)] \end{aligned} \right) - \phi(\tau D_b(\varsigma_n, \varsigma_{n-1})) + \dots \\
 &+ \left(\begin{aligned} &\mathfrak{F}(\alpha) \mathfrak{F}(\tau^{n-m}) \mathfrak{F}(D_b(\varsigma_{m-1}, \varsigma_m)) \\ &+ \mathfrak{F}(\beta) \mathfrak{F}(\tau^{n-m}) \mathfrak{F} \left(\frac{[1+D_b(\varsigma_m, \varsigma_{m-1})] D_b(\varsigma_{m-1}, \varsigma_{m-2})}{1+D_b(\varsigma_m, \varsigma_{m-1})} \right) \\ &+ \mathfrak{F}(\gamma) \mathfrak{F}(\tau^{n-m}) \mathfrak{F} [D_b(\varsigma_m, \varsigma_{m-1}) + D_b(\varsigma_{m-1}, \varsigma_{m-2})] \\ &+ \mathfrak{F}(\delta) \mathfrak{F}(\tau^{n-m}) \mathfrak{F} [D_b(\varsigma_m, \varsigma_{m-2}) + D_b(\varsigma_{m-1}, \varsigma_{m-1})] \end{aligned} \right) - \phi(\tau^{n-m} D_b(\varsigma_{m-1}, \varsigma_m)).
 \end{aligned}$$

Consider the fact that ϕ and \mathfrak{F} are strongly monotone functions. There are

$$\begin{aligned}
 D_b(\varsigma_n, \varsigma_m) &\leq \alpha\tau D_b(\varsigma_n, \varsigma_{n-1}) + \frac{\beta[1+D_b(\varsigma_{n-1}, \varsigma_n)]D_b(\varsigma_n, \varsigma_{n+1})}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \\
 &+ \gamma\tau [D_b(\varsigma_{n-1}, \varsigma_n) + D_b(\varsigma_n, \varsigma_{n+1})] + \delta\tau [D_b(\varsigma_{n-1}, \varsigma_{n+1}) + D_b(\varsigma_n, \varsigma_n)] + \dots \\
 &+ \alpha\tau^{n-m} D_b(\varsigma_{m-1}, \varsigma_m) + \frac{\beta\tau^{n-m}[1+D_b(\varsigma_m, \varsigma_{m-1})]D_b(\varsigma_{m-1}, \varsigma_{m-2})}{1+D_b(\varsigma_m, \varsigma_{m-1})} \\
 &+ \gamma\tau^{n-m} [D_b(\varsigma_m, \varsigma_{m-1}) + D_b(\varsigma_{m-1}, \varsigma_{m-2})] \\
 &+ \delta\tau^{n-m} [D_b(\varsigma_m, \varsigma_{m-2}) + D_b(\varsigma_{m-1}, \varsigma_{m-1})].
 \end{aligned}$$

So,

$$D_b(\varsigma_n, \varsigma_m) \leq \left(\begin{aligned} &\| \alpha \| \| \tau \| \| D_b(\varsigma_n, \varsigma_{n-1}) \| \\ &+ \| \beta \| \| \tau \| \| \frac{[1+D_b(\varsigma_{n-1}, \varsigma_n)]D_b(\varsigma_n, \varsigma_{n+1})}{1+D_b(\varsigma_{n-1}, \varsigma_n)} \| \\ &+ \| \gamma \| \| \tau \| \| D_b(\varsigma_{n-1}, \varsigma_n) + D_b(\varsigma_n, \varsigma_{n+1}) \| \\ &+ \| \delta \| \| \tau \| \| D_b(\varsigma_{n-1}, \varsigma_{n+1}) + D_b(\varsigma_n, \varsigma_n) \| + \dots \\ &+ \| \alpha \| \| \tau^{n-m} \| \| D_b(\varsigma_{m-1}, \varsigma_m) \| \\ &+ \| \beta \| \| \tau^{n-m} \| \| \frac{[1+D_b(\varsigma_m, \varsigma_{m-1})]D_b(\varsigma_{m-1}, \varsigma_{m-2})}{1+D_b(\varsigma_m, \varsigma_{m-1})} \| \\ &+ \| \gamma \| \| \tau^{n-m} \| \| D_b(\varsigma_m, \varsigma_{m-1}) + D_b(\varsigma_{m-1}, \varsigma_{m-2}) \| \\ &+ \| \delta \| \| \tau^{n-m} \| \| D_b(\varsigma_m, \varsigma_{m-2}) + D_b(\varsigma_{m-1}, \varsigma_{m-1}) \| \end{aligned} \right) \rightarrow 0, \text{ as } n, m \rightarrow +\infty.$$

Then $\{\varsigma_n\}$ is Cauchy sequence. Since $(\chi, \mathfrak{A}, D_b)$ is a complete C^* -(AVBMS) $\exists u \in \chi \ni \varsigma_n \rightarrow u$ as $n \rightarrow \infty$. Now that

$$\begin{aligned}
 D_b(u, Mu) &\leq \tau [D_b(u, \varsigma_{n+1}) + D_b(\varsigma_{n+1}, Mu)] \\
 &= \tau [D_b(\varsigma_{n+1}, Mu) + D_b(u, \varsigma_{n+1})] \\
 &= \tau [D_b(L\varsigma_n, Mu) + D_b(u, \varsigma_{n+1})] \\
 \mathfrak{F}(D_b(u, Mu)) &= \tau [\mathfrak{F}(D_b(L\varsigma_n, Mu)) + \mathfrak{F}(D_b(u, \varsigma_{n+1}))] \\
 &\leq \tau [\mathfrak{F}(N(\varsigma_n, u)) - \phi D_b(\varsigma_n, u)] + \tau [\mathfrak{F}D_b(u, \varsigma_{n+1})] \\
 \| \mathfrak{F}(D_b(L\varsigma_n, Mu)) \| &\leq \| \tau \| \| \mathfrak{F}(D_b(u, \varsigma_{n+1})) \| + \| \tau \| \| \mathfrak{F}\alpha \| \| D_b(\varsigma_n, u) \| \\
 &+ \| \tau \| \| \mathfrak{F}\beta \| \| \frac{[1+D_b(\varsigma_n, L\varsigma_n)]D_b(u, Mu)}{1+D_b(\varsigma_n, u)} \| \\
 &+ \| \tau \| \| \mathfrak{F}\gamma \| \| D_b(\varsigma_n, \varsigma_{n+1}) + D_b(u, Mu) \| \\
 &+ \| \tau \| \| \mathfrak{F}\delta \| \| D_b(\varsigma_n, Mu) + D_b(u, L\varsigma_n) \| - \| \tau \| \| \phi D_b(\varsigma_n, u) \|.
 \end{aligned}$$

Using the property of ϕ , we have

$$\begin{aligned}
 \| \mathfrak{F}(D_b(L\varsigma_n, Mu)) \| &\leq \| \tau \| \| \mathfrak{F}(D_b(u, \varsigma_{n+1})) \| + \| \tau \| \| \mathfrak{F}\alpha \| \| D_b(\varsigma_n, u) \| \\
 &+ \| \tau \| \| \mathfrak{F}\beta \| \| \frac{[1+D_b(\varsigma_n, L\varsigma_n)]D_b(u, Mu)}{1+D_b(\varsigma_n, u)} \| \\
 &+ \| \tau \| \| \mathfrak{F}\gamma \| \| D_b(\varsigma_n, \varsigma_{n+1}) + D_b(u, Mu) \| \\
 &+ \| \tau \| \| \mathfrak{F}\delta \| \| D_b(\varsigma_n, Mu) + D_b(u, L\varsigma_n) \|
 \end{aligned}$$

where \mathfrak{F} is strongly monotone, then

$$\begin{aligned}
 \| (D_b(L\varsigma_n, Mu)) \| &\leq \| \tau \| \| (D_b(u, \varsigma_{n+1})) \| + \| \tau \| \| \alpha \| \| D_b(\varsigma_n, u) \| \\
 &+ \| \tau \| \| \beta \| \| \frac{[1+D_b(\varsigma_n, L\varsigma_n)]D_b(u, Mu)}{1+D_b(\varsigma_n, u)} \| \\
 &+ \| \tau \| \| \gamma \| \| D_b(\varsigma_n, \varsigma_{n+1}) + D_b(u, Mu) \| \\
 &+ \| \tau \| \| \delta \| \| D_b(\varsigma_n, Mu) + D_b(u, L\varsigma_n) \| \\
 &= \| \tau \| \| (D_b(u, \varsigma_{n+1})) \| \\
 &+ \| \tau \| \left[\begin{aligned} &\| \alpha \| \| D_b(\varsigma_n, u) \| \\ &+ \| \beta \| \| \frac{[1+D_b(\varsigma_n, \varsigma_{n+1})]D_b(u, Mu)}{1+D_b(\varsigma_n, u)} \| \\ &+ \| \gamma \| \| D_b(\varsigma_n, \varsigma_{n+1}) + D_b(u, Mu) \| \\ &+ \| \delta \| \| D_b(\varsigma_n, Mu) + D_b(u, \varsigma_{n+1}) \| \end{aligned} \right]
 \end{aligned}$$

as $\varsigma_n \rightarrow u$ and $\varsigma_{n+1} \rightarrow u$ as $n \rightarrow \infty$, we obtain

$$\| 1 - \tau\beta - \tau\gamma - \tau\delta \| \| D_b(u, Mu) \| \leq \left(\| \tau \| \| \alpha \| \| D_b(\varsigma_n, u) \| + \| \tau \| \| 1 + \delta \| \| D_b(u, \varsigma_{n+1}) \| \right) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Hence $\| D_b (Mu, u) \| = 0$ since $\| 1 - \tau\beta - \tau\gamma - \tau\delta \| > 0$. As a result, $Mu = u$ which is uis a fixed point of M . Similarly we can demonstrate this $Lu = u$. Hence $Lu = Mu = u$. This show that u is common fixed point of L and M .

Let v be a different fixed point common to L and M . (i.e) $Lv = Mv = v \ni u \neq v$ we have $D_b (u, v) = D_b (Lu, Mv)$ then,

$$\begin{aligned} \mathfrak{F} (D_b (u, v)) &= \mathfrak{F} (D_b (Lu, Mv)) \leq \mathfrak{F} (N (u, v)) - \phi (D_b (u, v)) \\ \| \mathfrak{F} (D_b (Lu, Mv)) \| &\leq \| \mathfrak{F}\alpha \| \| D_b (u, v) \| + \| \mathfrak{F}\beta \| \| \frac{[1+D_b(u, Lu)]D_b(v, Mv)}{1+D_b(u, v)} \| \\ &+ \| \mathfrak{F}\gamma \| \| D_b (u, Lu) + D_b (v, Mv) \| \\ &+ \| \mathfrak{F}\delta \| \| D_b (u, Mv) + D_b (v, Lu) \| - \| \phi D_b (u, v) \| \end{aligned}$$

Using the property of ϕ , we get

$$\begin{aligned} \| \mathfrak{F} (D_b (Lu, Mv)) \| &\leq \| \mathfrak{F}\alpha \| \| D_b (u, v) \| + \| \mathfrak{F}\beta \| \| \frac{[1+D_b(u, Lu)]D_b(v, Mv)}{1+D_b(u, v)} \| \\ &+ \| \mathfrak{F}\gamma \| \| D_b (u, Lu) + D_b (v, Mv) \| \\ &+ \| \mathfrak{F}\delta \| \| D_b (u, Mv) + D_b (v, Lu) \| \end{aligned}$$

Where \mathfrak{F} is strongly monotone, then

$$\begin{aligned} \| (D_b (Lu, Mv)) \| &\leq \| \alpha \| \| D_b (u, v) \| + \| \beta \| \| \frac{[1+D_b(u, Lu)]D_b(v, Mv)}{1+D_b(u, v)} \| \\ &+ \| \gamma \| \| D_b (u, Lu) + D_b (v, Mv) \| \\ &+ \| \delta \| \| D_b (u, Mv) + D_b (v, Lu) \| \\ &= \| \alpha + 2\delta \| \| D_b (u, v) \| \\ &\leq \| \tau\alpha + \beta + (\tau + 1)\gamma + \tau(\tau + 1)\delta \| \| D_b (u, v) \| \\ &< \| D_b (u, v) \| . \end{aligned}$$

Which is a contradiction. Hence $\| D_b (u, v) \| = 0$ and $u = v$. Thus u is a unique common fixed point of L and M . \square

Corollary 2.2. Let $(\chi, \mathfrak{A}, D_b)$ is a complete C^* -(AVbMS). Let $M : \chi \rightarrow \chi$ be a contractive mapping and

$$\mathfrak{F} (D_b (M^n \varsigma, M^n \eta)) \leq \mathfrak{F} (L (\varsigma, \eta)) - \phi (D_b (\varsigma, \eta)) \tag{2.3}$$

$$L (\varsigma, \eta) \leq \alpha D_b (\varsigma, \eta) + \gamma [D_b (\varsigma, M^n \eta) + D_b (\eta, M^n \eta)] + \delta [D_b (\varsigma, M^n \eta) + D_b (\eta, M^n \eta)]$$

for all $\varsigma, \eta \in \chi$, where $\tau \in \mathfrak{A}'_+$, $\alpha + \gamma + \delta \geq 0$ with $\tau\alpha + \gamma(\tau + 1) + \delta(\tau(\tau + 1)) < 1$. \mathfrak{F} and ϕ are *-homomorphisms and the constraint $\mathfrak{F} (a) \leq \phi(a)$. Then M have a unique fixed point in X .

Corollary 2.3. Let $(\chi, \mathfrak{A}, D_b)$ is a complete C^* -(AVbMS). Let $L : \chi \rightarrow \chi$ be a contractive mapping and

$$\mathfrak{F} (D_b (L^n \varsigma, L^n \eta)) \leq \mathfrak{F} (N (\varsigma, \eta)) - \phi (D_b (\varsigma, \eta)) \tag{2.4}$$

$$\begin{aligned} N (\varsigma, \eta) &\leq \alpha D_b (\varsigma, \eta) + \frac{\beta[1+D_b(\varsigma, L^n \varsigma)]D_b(\eta, L^n \eta)}{1+D_b(\varsigma, \eta)} \\ &+ \gamma [D_b (\varsigma, L^n \eta) + D_b (\eta, L^n \eta)] + \delta [D_b (\varsigma, L^n \eta) + D_b (\eta, L^n \eta)] \end{aligned}$$

for all $\varsigma, \eta \in \chi$, where $\tau \in \mathfrak{A}'_+$, $\alpha + \beta + \gamma + \delta \geq 0$ with $\tau\alpha + \beta + \gamma(\tau + 1) + \delta(\tau(\tau + 1)) < 1$. \mathfrak{F} and ϕ are *-homomorphisms and the constraint $\mathfrak{F} (a) \leq \phi(a)$. Then L have a unique fixed point in χ .

Example 2.4. Let $\chi = [0, 1]$ and $\mathfrak{A} = \mathbb{C}$ with a norm $\| \varsigma \| = |\varsigma|$ be a real C^* -algebra. We define $\mathbb{C}^+ = \{(\varsigma, \eta) \in \mathbb{C} : \varsigma = Re(\xi) \geq 0, \eta = Im(\xi) \geq 0\}$. The partial order \leq with respect to the C^* -algebra \mathbb{C} . $Re(\varsigma_1) \leq Re(\varsigma_2)$ and $Im(\eta_1) \leq Im(\eta_2) \forall (\varsigma_1, \eta_1), (\varsigma_2, \eta_2) \in \mathbb{C}$. Let $D_b : \chi \times \chi \rightarrow \mathbb{C}$ suppose that $D_b (\varsigma, \eta) = 2(|\varsigma - \eta|, |\varsigma - \eta|)$ for $\varsigma, \eta \in \chi$. Then, $(\chi, \mathfrak{A}, D_b)$ is a C^* -algebra valued \mathfrak{b} -metric space where $\tau = 1$ in theorem 2.1. Let $\mathfrak{F}, \phi : \mathbb{C}^+ \rightarrow \mathbb{C}^+$ be the mappings defined as follows: For $\mathcal{T} = (\varsigma, \eta) \in \mathbb{C}^+$

$$\psi (\mathcal{T}) = \begin{cases} (\varsigma, \eta), & \text{if } \varsigma \leq 1 \text{ and } \eta \leq 1, \\ (\varsigma^2, \eta), & \text{if } \varsigma > 1 \text{ and } \eta \leq 1, \\ (\varsigma, \eta^2), & \text{if } \varsigma \leq 1 \text{ and } \eta > 1, \\ (\varsigma^2, \eta^2), & \text{if } \varsigma > 1 \text{ and } \eta > 1. \end{cases}$$

and for $\mathcal{G} = (\mathcal{G}_1, \mathcal{G}_2) \in \mathbb{C}^+$ with $\mathcal{P} = \min \{\mathcal{G}_1, \mathcal{G}_2\}$,

$$\phi(\mathcal{G}) = \begin{cases} \left(\frac{\mathcal{P}^2}{2}, \frac{\mathcal{P}^2}{2}\right), & \text{if } \mathcal{P} \leq 1 \\ \left(\frac{1}{2}, \frac{1}{2}\right), & \text{if } \mathcal{P} > 2 \end{cases}$$

Then, \mathfrak{F} and ϕ are satisfying in definition 1.4 and 1.5. Let $L, M : \chi \rightarrow \chi$ be defined as follows:

$$L(\varsigma) = \begin{cases} 0, & \text{if } 0 \leq \varsigma \leq \frac{1}{2} \\ \frac{1}{4}, & \text{if } \frac{1}{2} < \varsigma \leq 1 \end{cases} ; M(\varsigma) = \frac{1}{32}, \text{ for } \varsigma \in \chi$$

Then, L and M are satisfying in theorem 2.1. Let $\alpha = \frac{1}{8}$, $\beta = 0$, $\gamma = \frac{1}{16}$ and $\delta = \frac{1}{16}$. It demonstrates that:

$$\mathfrak{F}(D_b(L\varsigma, M\eta)) \leq \mathfrak{F}(N(\varsigma, \eta)) - \phi(D_b(\varsigma, \eta)) \quad \forall \varsigma, \eta \in \chi \text{ with } \eta \leq \varsigma.$$

Hence, Theorem 2.1 is satisfied. Then demonstrate that 0 is a unique common fixed point of L and M .

3 Conclusions

In Theorem 2.1 we have formulated a contractive conditions to modify and extend the concept of common fixed point theorem for C^* -algebra valued b -metric space via (ϕ, \mathfrak{F}) -contractive mapping. The existence and uniqueness of the result is presented in this article. We have also given some example which satisfies the contractive condition of our main result. Our result may be the vision for other authors to extend and improve several results in such spaces and applications to other related areas.

Acknowledgements

The authors thanks the management, Asian Research Journal of Mathematics for their constant support towards the successful completion of this work. We wish to thank the anonymous reviewers for a careful reading of manuscript and for very useful comments and suggestions.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Ma Z, Jiang L, Sun H. C^* -algebra-valued metric spaces and related fixed point theorems. Fixed Point Theory Application. 2014;206.
- [2] Ma Z, Jiang L. C^* -algebra-valued b-metric spaces and related fixed point theorems. Fixed Point Theory Appl. 2015;1:1-12.
- [3] Alsulami HH, Agarwal RP, Karapinar E, Khojasteh F. A short note on C^* -valued contraction mappings. J. Inequalities, Appl. 2016;50.
- [4] Kamran T, Postolache M, Ghiura A, Batul S, Ali R. The banach contraction principle in C^* -algebra-valued b-metric spaces with application. Fixed Point Theory Application. 2016;10.
- [5] Ozer O, Omran S. On the generalized C^* -valued metric spaces related with Banach fixed point theory. Int. J. Adv. Appl.sci. 2017;4:35-37.
- [6] Schroer B. Lectures on algebraic quantum field theory and operator algebra. Acta Math. Sin. 2001;23:2205-2212.
- [7] Murphy GJ. C^* -algebras and operator theory. Academic Press: London, UK. 1990.

- [8] Rahmah Mustafa, Saleh Omran, Quang Ngoc Nguyen. Fixed point theory using \mathfrak{F} contractive mapping in C^* -algebra valued B-metric space. Mathematics. 2021;9:92.
- [9] Sakai S. C^* -algebras and W^* -algebras. Springer: Berlin, Heidelberg; 1998.
Print ISBN 978-3-540-63633-5, Online ISBN 978-3-642-61993-9.
- [10] Zhu K. An introduction to operator algebras. CRC Press: Boca Raton, FL, USA; 1961.

© 2023 Maheshwaran et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/103141>