



Towards Sustainable Abundance: The Advancements in Tamarind Production Technology

Vasanthkumar S. S.^{a++*}, Rubika R.^{a++}, Sabarivasan M.^{a++},
Pooja U. K.^{b++} and Udhayakumar K.^{c++}

^a Department of Plantation, Spices, Medicinal and Aromatic Crops, HC and RI- TNAU, Periyakulam, India.

^b Department of Plantation, Spices, Medicinal and Aromatic Crops, HC and RI- TNAU, Coimbatore, India.

^c Department of Fruit Science, HC and RI- TNAU, Periyakulam, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i203932

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107126>

Review Article

Received: 28/07/2023

Accepted: 04/10/2023

Published: 11/10/2023

ABSTRACT

Tamarind, a tropical fruit-bearing tree, is essential for food, pharmaceutical, and industrial applications. Advancements in tamarind production technology are needed to ensure sustainable cultivation and efficient processing. This review focuses on innovative cultivation practices, breeding strategies, pest management, post-harvest practices, and value addition. It explores intercropping, agroforestry, and water-efficient irrigation systems to maximize land utilization and water resources. Integrated pest management strategies, such as biological control and organic pesticides, are proposed to reduce chemical dependency, minimize environmental impact, and maintain ecological balance. Post-harvest technology, including innovative harvesting, cleaning, storage, and

⁺⁺ PG Scholar;

*Corresponding author: E-mail: kalaivasanth17@gmail.com;

processing techniques, is evaluated to maintain product quality and shelf life. The review also examines the economic viability and value addition opportunities within the tamarind industry, identifying potential product diversification and reducing the carbon footprint.

Keywords: *Imli; tamarind cultivation; Tamarindus indica; sustainable practices; tamarind processing.*

1. INTRODUCTION

Tamarind (*Tamarindus indica* L.) is a monotypic genus belonging to the subfamily Caesalpinioideae of the family Leguminosae (Fabaceae) with a somatic chromosome number of $2n = 24$ [1]. The fruit of the tamarind tree, or "Assam tree," is known as "imli" or "Indian date." This multipurpose tree can grow in any type of soil and is frost- and drought-tolerant [2]. Mostly utilized for its fruits, which are either consumed raw or processed, used as a spice or condiment, or the fruits and seeds are processed for various uses [3]. India produces roughly 2.5 lakh metric tonnes of tamarind pulp each year, and there is potential for exporting juice concentrates and pulp powder to European nations. When properly dried with salt, tamarind pulp has a great ability to preserve food [4]. Since the tamarind has so many beneficial qualities, both rural and urban residents use almost every component of the tree [5].

2. ORIGIN, HISTORY AND DISTRIBUTION

Tamarindus indica L. is a tropical plant native to tropical Eastern Africa [6]. Ethiopian traders introduced it to India centuries ago [7]. Tamarind, originating from Persian and Arab sources, reached India and was called 'Tamar-i-hind', the Persian word means 'Date of India' Its Sanskrit name 'Amlika' indicates its ancient presence, with mention in Indian Brahmasamhita scriptures from 1200 to 200 BC [8]. It is mentioned in Ayurvedic literature as "amla" (Caraka Samhita, and Sushruta Samhita) [9]. The fruit has medicinal properties and culinary applications, with starch-rich seeds used in textile industries and consumption. Tamarind is said to be pan tropical [7]. Widespread in tropics and subtropics, cultivated in 54 countries, particularly in African, Asian, and South American regions [5,10]. Cultivating in various states of India, including Madhya Pradesh, Bihar, Andhra Pradesh, Karnataka, Tamil Nadu, West Bengal, Orissa, Kerala [11].

Taxonomy: Tamarind belongs to the 3rd largest family of angiosperms, Fabaceae with 700

genera and nearly 17000 species [5]. It is a member of monotypic genus *Tamarindus* and comprises one species, *indica*.

Kingdom	: Plantae
Subkingdom	: Tracheobionta
Superdivision	: Spermatophyta
Division	: Magnoliophyta
Class	: Magnoliopsida
Subclass	: Rosidae
Superorder	: Rosanae
Order	: Fabales
Family	: Fabaceae
Subfamily	: Caesalpinioideae
Tribe	: Detarieae
Genus	: <i>Tamarindus</i>
Species	: <i>Tamarindus indica</i>

[9]

Botanical Description: Tamarind is an evergreen tree with a 30m tall, 8m diameter trunk, light grey to brown color and short fissured, scaly trunk [5]. Its crown is dense and spans up to a circumference of 12 m in diameter [12]. The plant has wide, crooked, and droopy branches with red exudate when damaged. Its roots are deep, with tap root and lateral root systems. Leaves are paripinnate, distributed alternately, and have a short petiole. The leaves fold in absence of light due to lutein degradation. The inflorescence is a terminal raceme with up to 18 flowers, with flower buds developing within 20 days. The flower development takes 18 to 26 days, with pedicellate and bracteate flowers. The flowers are bisexual, protogynous, nectariferous, and entomophilous, leading to cross pollination. Anthesis occurs at 6 AM, with a cream, pale yellow, pink, or white turbinate calyx and a cream, pale yellow, pink, or white corolla. The androecium consists of three fertile stamens, with anthers that dehisce longitudinally after opening the flowers. The gynoecium is superior, with an adnate ovary and up to 18 ovules.

The style is green, long, and hooked with a subcapitate stigma. Stigma is most receptive on the day of flower opening. Flowers are irregular in length and diameter.

Table 1. Nutraceuticals of tamarind

Source	[26]	[27]	[28]	[29]
Nutrient	Food value			
Water	17.8–35.8g	-	-	35.29 g per 100 g
Thiamin	0.33 mg	-	-	-
Protein	2-3g	9.15%	2.00-9.10%	-
Riboflavin	0.1 mg	-	-	-
Fat	0.6g	6.24%	0.50-3.10%	-
Niacin	1.0 mg	-	-	-
Carbohydrates	41.1–61.4g	60.02%	56.70-82.60	50.07 g per 100 g)
Phosphorus	34-78 mg	-	-	-
Iron	0.2-0.9 mg	-	-	-
Pectin (calcium pectate)	2.4 %	-	2.00-4.00	-
Vitamin C	44 mg	-	-	-
Fiber	2.9g	7.16%	2.20-18.30	4.13 g per 100 g
Ash	2.6-3.9g	6.24%	2.10-3.30	-
Tartaric acid (free)	9.8 %	-	8.40-12.40	-
TSS	54-69.9 %	-	-	-
Calcium	34-94 mg	-	-	-
Tartaric acid(combined)	6.7 %	-	8.00-18.00	-
Invert sugar	38.2 %	-	25.00-45.00	-
Moisture	-	11.19%	15.00-30.00%	-
Starch	-	-	5.70	-
Tannin, (mg)	-	-	600.00	-
Ascorbic acid, (mg)	-	-	3.00-9.00	-
E-carotene equivalent (Pg)	-	-	10.00-60.00	-
Thiamine (mg)	-	-	0.18-0.22	-
Riboflavin (mg)	-	-	0.07-0.09	-
Niacin (mg)	-	-	0.60	-

Table 2. Some chemical composition of tamarind pulp

Parts	Chemical constituents [18]
Leaves.	Pulps contain invert sugar, citric acid, pipercolic acid, nicotinic acid, 1-malic acid, volatile oils, vitamins B3, C, vitexin, isovitexin, benzyl benzoate, cinnamates, serine, pectin, beta alanine, proline, phenylalanine, leucine, potassium, 1-malic acid, tannin, and glycosides.
Fruits.	Fruits contain Furan derivatives and carboxylic acid. Phlorotannins, apple acid, grape acid, succinic acid, citric acid, tartaric acid, pectin, invert sugar.
Seeds.	Seed extract contains campesterol, β -amyirin, β -sitosterol, palmitic acid, oleic acid, linoleic acid, and eicosanoic acid. Mucilage, arabinose, xylose, galactose pectin, glucose, and uranic acid were also found. A new bufadienolide and cardioid were identified. Cellulose, albuminoid, amyloids, phytohemagglutinins, and chitinase were also found.
Stem bark	Tannins, saponins, glycosides, peroxidase and lipids
Root bark	N-hexacosane, eicosanoic acid, β -sinosterol, (+)-pinitol, octacosanyl ferulate, 21-oxobehenic acid.

Table 3. Physico-chemical properties of tamarind

Physico-chemical properties	Pulp	Seed	References
Saponification value (mg KOH g-1)	301.3	266.6	[29]
Iodine value	120.6	78.1	
Unsaponified matter (g kg-1)	139.0	31.3	
Acid value (g kg-1)	896.0	292.6	
Free fatty acid (g kg-1)	448.0	46.3	
Peroxidase value (m Eq kg-1)	123.3	98.9	

Table 4. Mineral composition of tamarind

Mineral Composition of fruit Pulp Mineral mg/100g	Pulp	Pulp [27]	Seed	Seed [30]	Kernel	Testa
Calcium	81.0-466.0	21.57	9.3-786.0	10.0	120.0	100.0
Phosphorus	86.0-190.0	17.10-18.40	68.4-165.0	25.5		
Magnesium	25.0-72.0	10.54	17.5-118.3	15.0	180.0	120.0
Potassium	62.0-570.0	187.73	272.8-610.0	21.0	1020.0	240.0
Sodium	3.0-76.7	112.76	19.2-28.8	2.1	210.0	240.0
Copper	0.8-1.2		1.6-19.0			
Iron	1.3-10.9	1.05	6.5	75.9	80.0	80.0
Zinc	0.8-1.1		2.8		100.0	120.0
Nickel	0.5					
manganese		0.13	0.9			

Source: [31]

Table 5. Uses of tamarind

Food	Fruit pulp is used as a preservative and in various culinary dishes like chutney, tamarind juice concentrate, pulp powder, jams, syrups, candy, and souring porridge. It is also consumed raw as a snack.
Fodder	Animal feed made up of fried seeds and foliage
Agriculture	Flowers are used to make honey, and their roots fix nitrogen in the soil.
Fuel	Tamarind wood is potential for charcoal and firewood, but not widely used in Southeast Asia.
Timber	The sapwood is light yellow and the heartwood is dark purplish brown, marketed in the USA as "madeira mahogany," is used for various tools, furniture, utensils, poles, posts, boats, and construction materials.
Tannins	Tannins found in both leaves and bark can be used to make ink and dye.
Lipids	Seed oil is used in paints, varnishes, and oil lamps.
Medicine	Bark decoction, leaf decoction, pulp, and seeds are all plant parts with medicinal value for treating sores, ulcers, throat infections, rheumatism, cough, fever, intestinal worms, conjunctivitis, scurvy, dysentery, and diarrhea.
Other uses	Wood is pulverized for mulch, cleaned for utensils, and used for fabric size. Trees serve as windbreaks, living fences, and attractive plants in parks, roadways, and riverbanks.

Source: [32-34]

Table 6. Tamarind plantations in four states

Tamil Nadu	Maharashtra	Karnataka	Andhra Pradesh
85 clones evaluated at Periyakulam Horticultural College [35].	120 accessions evaluated at Pune College of Agriculture [37].	Forest department of Karnataka and	Department of Forest records 52 tree genotypes from tamarind tree surveys in Chittoor and Anantapur districts.
Institute of Forest Genetics and Tree Breeding identified 12 phenotypes in 1994 at Kanyakumari [36].	87 genotypes were identified and categorized as sour, sweet and red type [38].	University of Agricultural Sciences, Bangalore.	Tamarind genotypes were classified into five groups based on fruit characteristics: long, bold, medium, small, curved, irregular, and sweet [40].
	Fruit Research Station, Aurangabad has 350 trees under germplasm evaluation maintained under Marathwada Agricultural University, Parbhani	collaborate to evaluate 230 plus trees and 40 plus tree lines. [39].	
	Out of 195 plus elite trees, 5 elite types (T-9, T-10, T-11, T-12 and T-13) were identified at Ahamadnagar and Satara districts [25].		

The pods, oblong, curved, or straight fruits, are 5-10 cm long and 2 cm wide. When raw, they are brittle and light grey to brown. When ripe, the pulp softens and the seeds are hard, red to purple-brown, orbicular to rhomboid, flat, glossy, and have a central depression. Pods take 10 months to ripen and are indehiscent [13].

Properties of Tamarind: A typical fruit or pod has 55% pulp, 34% seeds, and 11% shell and fibers [14]. The testa, or seed coat, makes up 20–30% of the seed, and the endosperm, or kernel, makes up 70–75% [15]. The pods, which have a variety of uses and may be used either processed or raw. The other fruit parts, which are also eaten, contain minerals and nutrients [16]. Pulp contains high sugar content (30-40%), making it suitable for commercial applications like concentrates, pickles, confections, and powders [17].

Physical description: Tamarind had tall pods, brown shell, red flesh, and six seeds in its fruit. The fruits had a transverse diameter between 2.50 and 3.00 cm and a longitudinal diameter between 1.30 and 1.70 cm. The length ranged from 3.70 to 11.10 cm [18].

Chemical Composition: Tamarind pulp is rich in minerals, protein, and carbs but lacks water. It contains 20.6 percent water, 3.1% protein, 0.4 percent fat, 70.8 percent carbohydrates, 3.0% fiber, and 2.1% ash. The fruit's exact composition varies depending on its cultivation [19].

Physicochemical characterization: Tartaric acid, a rare biological source, is the primary acid in tamarind pulp. Its [SS/TA] ratio measures tamarind maturity and flavor, indicating the equilibrium between sugars and organic acids. In comparison to other fruits, Tamarind contains higher levels of vitamin C (4.79 mg/100 g) and folates (59.35 g/100 g) compared to other fruits, but only 108.78 g of vitamin E per 100 grams [9].

Antioxidant Properties of *Tamarindus indica*: Polyphenols are the primary class of natural antioxidants found in plants, found in various parts like bark, fruits, leaves, nuts, roots, and seeds [20]. Flavonoids, a subclass of polyphenols, are the second-largest family. *Tamarindus indica*, a plant with remarkable antioxidant properties, is a notable example [21].

Genetic Resources: All Tamarind plantations in India have originated from seedlings and are heterogenous having diverse characteristics. Seedling plants exhibit heterozygosity in

morphology, physiology and phenology [22]. This extends the range of desirable traits for crop improvement. Brown pulped and Red pulped types were reported [23]. Tamarind plantations in Maharashtra and Madhya Pradesh showed wide variations in sweetness, acidity, size, shape and bearing habits [24,25].

3. SWEET TAMARIND

Pithecellobium dulce (Roxb.) Benth. Various nutrients and antioxidants are abundant in delicious tamarind fruits. Root and bark decoctions are taken orally for diarrhea, hemoptysis, chest congestion, and internal ulcers. Seed juice is inhaled for chest congestion, and pulverized seeds are ingested for internal ulcers. Leaves can be applied as a plaster, and when taken with salt, can cure indigestion but also produce abortion. Root bark may also be used for dysentery and as a febrifuge [41].

Crop Improvement: Significant crop improvement can be done by selecting plus trees and maintaining by multiplication through clonal propagation.

Varieties: The regional varieties fall into two categories: "Brown sour types" and "Reddish sweet types."

Local selections are: Cumbum selection from Madurai district; Rahuri selection from Maharashtra; Karur urigam tamarind from Karur; Kangadevanapalli tamarind and Dever Ullimangalam tamarind [40] Bangalore Tamarind, Tumkur Tamarind, Hosur Tamarind, Krishnagiri Tamarind, Natham Tamarind, Nagarkoil Tamarind, Villupuram Tamarind, Ranchi Tamarind, are some of the most well-known local varieties [42].

Below is a list of some tamarind improvements.

1. **Periyakulam-1 (PKM-1):** It was released in 1992 by Horticultural College and Research Institute, Periyakulam. It is a clonal selection from germplasm. It is an early bearer. Its high pulp recovery is 39%. The interior side of the pulp exhibits the distinctive white colour will be, signifying its exceptional quality. This choice offers significant levels of ascorbic acid (3.95 mg/100 g) and tartaric acid (17.1%). It has a high yield of 263 kg/ha, Typically, seeds and grafts are used for propagation [43,44] findings showed that variety PKM-1 was

superior in terms of plant height (445.33–569 cm), TSS (480 Brix), total sugar (40.33–40.33 g), and lower acidity (6.7 mg) in West Bengal.

2. **Urigan:** In Tamil Nadu's Dharmapuri district, it is a superior local genotype that may be found near Urigan, a small village near Thenkanikottai. The tree is little and has branches that resemble those on umbrella. The pods are long, fleshy, and tasteful. The pods have a typical inward curve and are flat. Each fruit weighs between 150 and 200 grams. The range of seeds is 10 to 12. Flowers start to bloom slightly later than usual, in March, and fruits are ready to be picked in July. In a trial by [44] the variety Urigan performed better than other varieties in West [45].
3. **Prathisthan and HO 263:** These cultivars were released for commercial growth in the area by Fruit Research Station, Himal Bagh, Aurangabad, Maharashtra [40].
4. **NTI-19 (DST-1):** This variety produces well in the Karnataka district of Dharwad. The pods are large in size. In 4-5 years after planting, the grafted plants begin to produce enormous pods. The fifth year, each plant produces an average of roughly 3 kg.
5. **Red tamarind:** The Tamil Nadu Agricultural University advises cultivating this type. The pulp, which is red in colour and in high demand in Arabian nations.
6. **GKVK-6 and GKVK-33:** These two are local varieties made released by the Gandhi Krishi Vignana Kendra, Bangalore's UAS Division of Horticulture [42].
7. **GKVK 17:** When compared to GKVK-6, this cultivar exhibits a regular fruit-bearing habit and a tree canopy resembling an umbrella with average yield of 100.81 kg/tree. It has been recommended for the Southern regions of Karnataka [4].
8. **Yogeswari,** an attractive red pulp variety that has been regarded as a special variety, is gaining in popularity [40].

Soil: Tamarind tree thrives in various soil types, including poor, rocky terrain, ravines, and degraded terrain in India. It tolerates sodic and salty soils and requires minimal cultivation [46]. Tamarind growth thrives in loamy, deep, well-drained alluvial soil, promoting long tap roots, while red sandy loam soils in Andhra Pradesh, India, are suitable [47] silt-clay and clay soils were not suitable [48]. The tree cannot stand

waterlogging it thrive next to termite or ant mounds. The ideal pH for tamarind being 5.5–6.8, which is slightly acidic [29].

Climate: Tamarind has a wide geographic distribution in the sub- and semi-arid tropics. Tamarind thrives on evenly distributed 500-1500mm annual rainfall [49]. Effect of climatic conditions on flowering and fruiting of *Tamarindus indica* (Fabaceae). Tamarind can withstand up to 4000 mm of annual rainfall. It is a tropical climate tree, can tolerate temperatures as low as 9.5 °C and as high as 33 °C annually. Young tamarind trees are vulnerable to fire, frost, and strong winds. With thick branches and a deep root system, they are nicknamed "hurricane-resistant" due to their ability to withstand powerful typhoons and cyclones [50]. Grows at 0-2000 m, but absence at higher elevations is more due to temperature drop than elevation [51].

Propagation: Seed dissemination is the most common method, but it doesn't breed true to type due to heterozygosity and prolonged juvenile periods in plants [52]. In recent years, methods for vegetative propagation have been developed.

Seed propagation: Harvest fresh, mature fruits by examining a brittle epicarp, and cut or shake pods if nearby [53].

Ripe pods are harvested in March and April, dried in the sun, and seeds removed. They are separated from the edible pulp by hand kneading and washing. Clean seeds are dried in the shade and stored in gunny bags. Seeds have a shelf life of six months, and optimal harvesting occurs at peak maturity. An established tree can yield three quintals of fruit annually, containing around 80 kg of seeds. Direct sowing requires 20 kg of seeds per hectare of land [40]. Monkeys are the primary seed disperser in South India. Tamarind seeds germinate 13 days, potentially up to a month [54]. To achieve optimal germination, wait 45 days after planting. Hastened germination can be achieved by removing hard seededness with acid or mechanical scarification and soaking in hot water [29]

Soaking seeds in hot water breaks dormancy, promoting germination. Exposed batches to 50% sulphuric acid for 60 minutes had the highest germination rate [55,56] Tamarind seeds germination increased with methanol pretreatment, while upright planting with micropylar end facing upwards increased vigor and germination [57].

Nursery beds sown seeds in March and April, and within a year, seedlings reach a height of at least 80 cm. In India, seedlings are typically planted between 0.4 and 2 meters tall. In Tamil Nadu, transplants are better after two years. Overgrown seedlings can be successfully transplanted as stumped seedlings with stem and tap root lengths cut to 5 cm and 20-25 cm, respectively [28]

Vegetative propagation: Vegetative propagation procedures for large-scale multiplication and genotype characteristics include root and stem cuttings, air and stem-layering, budding, and grafting for tamarind [29,58]. Stem cuttings are an affordable and simple method for propagating tamarind, with a 39.9% success rate in October, followed by November and June [40]. Grafts gain popularity due to compact growth and early yields, bearing fruit in the third or fourth year after planting [59]. Vegetative reproduction techniques like shield and patch budding, cleft grafting, whip grafting, and approach grafting are effective but costly and time-consuming. Veneer and soft wood grafting have shown potential for propagating tamarind, with graft success rates of 49% and 68%, respectively. Wood grafting was successfully used to increase tamarind yield in March and April on 8-month-old seedlings with 22-32 cm height and 0.3-0.4 cm diameter [60]. Research shows 74% success rate in approach grafting compared to 25% standard grafting [61]. Study suggests patch-budding for faster tamarind reproduction on large scale. Tamarind has been tested using air layering, with shoots rooted at 37% when 4000 ppm IBA was applied. Higher hormone concentrations led to higher rooting percentages [40].

Tissue culture: On MS medium containing BAP, found high frequency regeneration from excised cotyledon. On MS, rooting was accomplished using IAA Micro shoots from tamarind seedling explants produced several shoots and rooted [62]. MS medium containing 0.5 mg/IBA, 0.5 mg/1 KN, and 0.1 mg/1 IAA was sufficient for generating 2-3 multiple shoots. On MS medium containing 0.2 mg/l of IAA, IBA, and NAA, Rhizogenesis was promoted [63].

The study found that ideal in vitro culture conditions for high-frequency plant regeneration from Urigam seedling stem pieces were found to have the highest response to shoot bud differentiation, earliest reaction, and the greatest number of multiple shoots per culture. Axillary buds of Urigam induced numerous shoots [64]

Orchard establishment: Plant from July to November, with a distance of 5-13 meters between plants. For a pure stand plantation, the final spacing should be 13 x 13 meters square or on a triangle. For optimal spacing, use larger density plants and thinning twice to achieve 8-15 x 8-15 meters [29]. Seedlings require 10mx10m or 13mx13m spacing, while grafts and buddings require 6mx6m spacing [42]. Tamarind is grown at 8 x 8 m, 8 x 12 m, or 12 x 12 m in some regions of India. For commercial planting, a spacing of 10 x 5 m has been advised, and 250 trees will be sufficient to cover one hectare [29] and 100 plants per acre are needed in 1 m³ trenches, filled with FYM, topsoil, and 10% dust. Plant seedlings or grafts in the center, keeping the graft-union above ground. Stack young seedlings to protect against wind [59].

Tissue culture-raised plants bloom at an average height of 3.7 meters, compared to seed-raised plants at 8 meters. To ensure optimal growth, it is recommended to plant more tissue culture-developed plants per unit area, with 156 plants per hectare at 9 m x 9 m spacing. Trees transplanted should be secured against animals and monkeys to prevent damage to seedlings and fruit consumption [65].

Irrigation: Irrigation aids early growth establishment, particularly during dry seasons [66]. Rainwater harvesting promotes growth and fruiting in dry areas; irrigation benefits tamarind [67].

Manuring: Neem cake, groundnut cake, and superphosphate combined with liquid manuring stimulated seedling growth in six-month-old trees [68].

Organic manures like farm yard manure and neem cake increased crop yield during rainy days [32]. A complete fertilizer mixture comprising 200 g of nitrogen, 150 g of phosphorus and 250 g of potassium along with 25 kg of farm yard manure applied annually to each tree is beneficial for a steady state of productivity [69].

Use of biofertilizers: Tamarind seedlings respond positively to soil inoculation of up to thirteen VAM fungi, resulting in increased plant height, leaf number, stem girth, and biomass production. This response increases phosphorous and zinc levels, root colonization, and external hyphal networks. Tamarind responds positively to *Gigaspora margarita* and *Glomus fasciculatum*, and VAM inoculation

significantly improves eight-month-old container-grown seedling growth. Rock phosphate is the best addition for optimal growth [70].

Pruning and training: Tamarind is a compact tree with symmetrical branches. Young trees require pruning to develop 3-5 well-spaced branches into the main scaffold structure. Regular pruning rejuvenates fruiting wood and regulates tree size in closely planted orchards [71].

Weed management: In new and old orchard, hoeing, hand weeding and ploughing the land 2-3 times in a year is done to suppress weed growth. Intercropping and mulching may be followed to control weeds [5].

Flowering, pollination and fruit set: Tamarind trees in South India bloom from April to July, with peak flowering occurring in May-June. New leaves appear in May, followed by flowers. Flower opening starts at 5:30 am, with peak anthesis at 6:30 am. Mature anthers are reddish brown and dehisce by longitudinal splitting of lobes. Pollen grains are round with an average diameter of 15 μ m, with a 96.5% fertile percentage. Stigma is receptive from one day before anthesis to two days after anthesis. Tamarind is predominantly cross-pollinated, with red ants playing a crucial role in pollination [40].

Tamarind fruit set is poor, causing large abscission of flowers and fruits. Factors contributing to poor fruit set include fertilization failure, sterility, defective pollen grains, slow growth, or early degeneration [72].

Tamarind flowers profusely, but fruit set is low under natural conditions (3-5%). Artificial cross-pollination can increase it to 50-55%, due to pollen limitation and floral abnormalities [24].

Tamarind's fruit set increased in controlled cross-pollinated flowers, indicating partial self-incompatibility. High pollinators and visiting trees led to increased fruit set, indicating tamarind's self-incompatibility [73]. 4-CPA, NAA, and GA were found to be more effective in increasing tamarind fruit set [74].

Floral biology: Inflorescence racemes are small, drooping racemes measuring 5-10 cm in length, with terminal and lateral drooping. The flowers are bisexual and 2 to 2.5 cm in diameter, with pedicels about 5 mm long. The bracts are fragrant and zygomorphic, nearly as long as the

flower bud. There are two boat-shaped bractioles, 8 mm long, and four sepals, up to 1.5 cm long. The plant has five petals, with the posterior and lateral ones being large and showy, slightly exceeding the calyx. There are three fertile stamens, with filaments connate and alternate with bristle-like staminodes. Anthers are transverse, reddish brown, and dehisce longitudinally. The ovary is superior and unicarpellary, with up to 18 ovules and a green, hooked style [75].

Fruit Growth and Development: The fruit growth process involves three phases: prematuration (63%), lignifications and stone cell formation (67%), and ripening (17%). Prematuration involves maximum cellular activity, while lignifications and stone cell formation increase fruit weight and size. Ripening occurs 29 weeks after anthesis and lasts 17% of the growth period. During ripening, the peel separates from the pulp, moisture content decreases, total sugars increase, and acid content doubles. The growth curve is sigmoid after maturation, and the fruit takes 245 days from set to harvest [23].

Crop diversification: Intercropping and multistory cropping are practices that grow annuals or short-duration crops in interspaces during their formative years. Intercropping maximizes land and space use efficiency, generates additional income during the unproductive phase, and protects interspace from losses from weeds, erosion, radiation, temperature, wind, and water. These practices enrich interspace with nitrogen-fixing legume crops like peas, gram, lentil, black gram, cowpea, cluster beet, and okra [5].

Pest: Tamarind faces various pests, including shothole borers, toy beetles, caterpillars, bagworms, mealy bugs, and scale insect *Ramaswamiella subnudula* and *Halothrips ceylonicus* Schmutz are yellowish thrips that attack tamarind, Controlling them is possible using dimethioate spray at 20-30 ml/10 litres of water or Fenthion spray at 10-15 ml/10 litres of water [76].

Eulemma unguifera Moore attacks flowers, weakening them and causing them to dry and fall from trees. Tamarind mealybug, *Drosichiella tamarindus* (Green), Tamarind weevil, *Sitophilus linearis* Tamarind fruit borer, *Phycita orthoclina* Meyrick [77].

Cydia palamedes Meyrick larvae bore into tender buds, causing caterpillars to cover flowers and buds with webs, causing crop losses. *Latolia lepila*, a slug caterpillar, can be a serious pest [78].

Disease: Nursery seedlings are affected by powdery mildew caused by *Oidium sp.*, causing defoliation and retarding early growth. Cercospora leaf disease attacks tamarind, causing severe defoliation and causing early growth retardation. In order to have effective control 3-4 sprays of wettable sulphur at 15-day intervals is the most economical method [79]. Seedling blight disease caused by *Macrophomina phaseolina* and *Rhizoctonia solani* is crucial during nursery stage. Raising seedlings in amended soils can reduce blight incidence [80].

Alternate Bearing: Regular bearing in cropping is not systematic, while alternate bearing is a genetic trait identified in tamarind trees. It can be managed through cultural practices and hormonal treatments [5].

Maturity index: Flesh color varies from green to brown, dark brown, red, or black, depending on variety [71]. At maturity, exocarp hardens and separates from pulp. Finger pressing tamarind fruits produces a hollow, loose sound, indicating pulp shrinkage and fruit readiness for harvesting, while shell brittleness develops [10]

Harvesting: Tamarind trees can be propagated through seeds or vegetatively, with tamarind trees bearing fruits 8-12 years after planting. Economical yields can be achieved after 7-10 years, with a lifespan of 50-60 years. Flowering typically occurs from June to July, and fruits are mature and ready for harvest in India during summer [81]. Mature Fruits-Fully ripen fruits are harvested. They are used for the preparation of various products like Sauce etc., [72]. Fruits are harvested by shaking the branches and the fallen fruits are collected. During the second harvest the left-over fruits are picked using fruit picker. Mechanical innovations in fruit bunch picking include branch and trunk shakers, suited for hard trunk trees like tamarind and soft trunk trees like apricot.

Yield: PKM 1 variety yields 250 kg per tree after 9 years, starting from 4-5 years, with grafts starting from 4-5 years [82]. A fully mature tree can produce 20 to 250 kg of fruits annually [72].

Processing: Tamarind can be processed using wet and dry methods, including drying, dehulling, defining, deseeding, pressing into cakes, and storage. It can also be pulped to remove seeds, fiber, and cellulose, creating edible pulp with minimal water. This pulp is then compacted into blocks similar to cheese and drum-dried [10]

Drying: [10] Tamarind dehulling and deseeding are crucial for drying. A local type was dried in the sun and subjected to plate drying at 50, 60, and 70°C. Mechanical drying at 70°C increased dampness expulsion rate [72]. Tamarind requires deseeding and sun drying for pulp preservation.

Dehulling: Tamarind can be dehulled by drying it completely in the sun and then removing the hull and pulp with sticks. The dehuller for tamarind was invented by TNAU, Coimbatore, India and has a deshelling rate of 100 kg/h, dehulling capacity of 94% [83].

The feed hopper was saturated with dried tamarind fruit, and rotating beaters applied impact force. Dehulled, un-hulling, and hull-less fruits were sent to the outlet. A dehuller was created at PHT center, UAS, Bangalore. Large curved fruits had a higher hulling percentage (79.48%), with a capacity of 500 kg per hour [71].

Deseeding: In Tamil Nadu, seeds are eliminated by hand beating tamarinds, A hammer type tamarind deseeder invented at Tamil Nadu Agricultural university [10].

Tamarind seed processing: Each tree produces 60–200 kg of seed. Which has numerous medicinal properties. There are three significant activities in Tamarind seed handling, roasting, decorticating and color sorting [84].

Roasting: Tamarind seed is roasted in a horizontal cyclone furnace. The furnace is having temperature of 220°C to brittle out the external layer of the tamarind seed [85].

Decorticating: Roasted seed is shipped off the decorticator for eliminating the external layer of the seed. After decortications husk will be removed [86].

Color sorting: Variety sorters sort seeds based on their external layer (Testa) and internal tone (creamy white). Un-decorticated seeds are removed for decorticating activity [87].

Value added products:

Tamarind seed Kernel Powder (TKP):

Tamarind seed Kernel Powder (TKP) is a pulverized tamarind used as an adhesive in the paper industry [88]. Tamarind seed powder is widely used in vegetable and food industries for thickening, stabilizing, and gelling food through tamarind gum or tamarin xyloglucan.

Tamarind gum: Tamarind gum is a user-friendly dye made by boiling tamarind seed powder in water, used for dyeing cotton and silk fabrics. This process is more efficient than solution dye preparation in powder form [89].

Tamarind seed oil: Tamarind seeds contain golden yellow oil about 7-9% oil content, used as a light and varnishing agent. Amber oil is also used [29].

Pulp based products:

Tamarind toffees: Toffee is a sweetmeat made from fruit pulp and sugar, wrapped in paper or after fiber loss. Tamarind balls are a type of sweetmeat made from sugar and pulp in India [90].

Tamarind beverage: Tamarind pulp is a popular beverage in southern and central America and Asia, used to create ready-to-serve drinks, syrups, and concentrates with a shelf life of six months. The pulp is mixed with water and hand squeezed for delicious drinks [91].

Tamarind pulp powder (TPP): TPP is an easy food product made by concentrating, drying, and milling pulp, with physicochemical characteristics varying depending on manufacturing method [92].

Tamarind pickle: Tamarind pickle is made from commercially available pulp. [10] The flavor of pickles is hot, spicy, and salty, making them long-lasting. Salt, increased acidity, and spices are responsible for preserving [93].

Tamarind candy: Tamarind candy is popular due to its natural sour-sweet flavor. It is made by boiling tamarind pulp with sugar and simmering it with minimal water. Sweetened tamarind fruit is prepared by peeling ripe fruits and pouring boiling sugar syrup at 62°C [94]

Sous: Traditional Jordanian drink made with tamarind, *Glycyrrhiza glabra* roots, and mulahatti and yastimadhu [95].

Tamarind Ade: The perfect blend of ripe pulp with sugar and water is used to make this delicious tamarind drink, which is also made in the Philippines and many tropical American countries [10].

Tamarind juice concentrate: Tamarind pulp is extracted from pods, separated juice, and hand-pulled into pulp, ensuring purity and free from contaminants [96].

Tamarind jam: Tamarind jam is made by shelling ripe fruits, boiling pulp at 100°C for 10 minutes, cooling, packing in sterile jars, and sealing [97].

Tamarind fruit leather: The soft, rubbery and sweet taste of fruit leather is a result of drying sheets of tamarind fruit pulp [10] Study examines drying techniques on tamarind leather, revealing that cabin drier and solar drier affect color.

Storage: The moisture content during storage dropped from 30% to approximately 20% [98]. Compared to storage in white polythene wrap, and plastic containers, the initial brown color might continue up to 4 months in metal and black polythene [71].

4. CONCLUSION

Tamarind production technology is essential for sustainable growth and development. Innovative practices like intercropping and agroforestry optimize land utilization, conserve resources, and promote biodiversity. Water-efficient irrigation systems overcome challenges, and variety selection improves productivity. Integrated pest management protects trees, and efficient post-harvest practices preserve nutritional content. Product diversification enhances revenue streams and attracts investors. Eco-friendly techniques promote responsible tamarind production, aligning with global sustainability goals.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Rajamanickam C, Ravindran C, Sundaraiya K, Vani V. Genetic diversity in tamarind (*Tamarindus indica* L.). *Pharma Innovation*. 2023;12(5):1921-1926.
- Rajamanickam C. Genetic variability, correlation and path analysis in tamarind (*Tamarindus indica* L.). *Journal of Spices and Aromatic Crops*. 2020;29(2).
- Shah N. *Tamarindus indica*: Introduction in India and culinary, medicinal, and industrial uses. *Asian Agrihist*. 2014;18(4):343-55.
- Venkataravana P, Sivappa MM, Priyadarshini S. GKVK-17: A new high yielding variety of Tamarind (*Tamarindus indica* L.) for Southern region of Karnataka. *IJCS*. 2020;8(1):1883-6.
- Singh S, Mishra D, Singh A. Tamarind (*Tamarindus indica* L.). *Tropical Fruit Crops Theory to Practical*. 2021;610-31.
- Sookying S, Duangjai A, Saokaew S, Phisalprapa P. Botanical aspects, phytochemicals, and toxicity of *Tamarindus indica* leaf and a systematic review of antioxidant capacities of *T. indica* leaf extracts. *Frontiers in Nutrition*. 2022; 9:977015.
- Chimsah F, Nyarko G, Abubakari A. A review of explored uses and study of nutritional potential of tamarind (*Tamarindus indica* L.) in Northern Ghana; 2020.
- Ravindran P, Divakaran M, Pillai G. *Handbook of herbs and spice*. Volume. 2012;1:86-115.
- Joshi AV, Urmila A, Swapnil G, Shahrukh S. Review on *Tamarindus indica*. *Journal of Pharmacognosy and Phytochemistry*. 2023;12(1):303-8.
- Sudha P, Rajkumar P, Joice AA, Sudagar I, Arulmari R. Postharvest technology of tamarind. *postharvest technology-recent advances, new perspectives and applications: IntechOpen*; 2022.
- Idris MM, Karamba H, Ayoubi R. *Tamarindus indica*-A Review Paper on Efficacy of Tamarind plant in Water Treatment and Purification. *Int J Sci Eng Res*. 2018;9.
- Rao AS, Kumar AA. Tamarind seed processing and by-products. *Agricultural Engineering International: CIGR Journal*. 2015;17:200-4.
- Choudhary S, Singh A, Rani M. Rhizosphere: Hotspot of soil-microbes-plant interaction. *A Monthly Peer Reviewed Magazine for Agriculture and Allied Sciences*. 2022;80.
- Muzaffar K, Kumar P. Tamarind-a mini review. *MOJ Food Process Technol*. 2017;5(3):296-7.
- Shankaracharya N. Tamarind-chemistry, technology and uses-a critical appraisal. *Journal of food Science and Technology*. 1998;35(3):193-208.
- Okello J, Okullo JB, Eilu G, Nyeko P, Obua J. Mineral composition of *Tamarindus indica* LINN (Tamarind) pulp and seeds from different agro-ecological zones of Uganda. *Food science & nutrition*. 2017; 5(5):959-66.
- Suliman AME, Alawad SM, Osman MA, Abdelmageed EA. Physicochemical characteristics of local varieties of tamarind (*Tamarindus indica* L), Sudan. *International Journal of Plant Research*. 2015;5(1):13-8.
- Hamacek FR, Santos PRG, de Moraes Cardoso L, Pinheiro-Sant'Ana HM. Nutritional composition of tamarind (*Tamarindus indica* L.) from the Cerrado of Minas Gerais, Brazil. *Fruits*. 2013;68(5): 381-95.
- De Caluwé E, Halamouá K, Van Damme P. *Tamarindus indica* L.–A review of traditional uses, phytochemistry and pharmacology. *Afrika focus*. 2010;23(1):53-83.
- Devi B, Boruah T. Tamarind (*Tamarindus indica*). *Antioxidants in Fruits: Properties and Health Benefits*. 2020:317-32.
- Lu M, Yuan B, Zeng M, Chen J. Antioxidant capacity and major phenolic compounds of spices commonly consumed in China. *Food Research International*. 2011;44(2):530-6.
- Pareek O, Awasthi O. Genetic resources of tamarind. *Indian Journal of Plant Genetic Resources*. 2002;15(3):197-202.
- Hernandez-Unzon H, Lakshminarayana S. Developmental physiology of tamarind fruit (*Tamarindus indica* L.) 1. *HortScience*. 1982;17(6):938-40.
- Karale A, Wagh A, Pawar B, More T. Association of fruit characters in tamarind. *JOURNAL-Maharashtra Agricultural Universities*. 1999;24(3):319-20.
- Keskar B, Karale A, Dhawale B, Choudhari K. Improvement of Tamarind (*Tamarindus indica* L.) by selection. *Maharashtra Journal of Horticulture*. 1989;4(2):121-4.
- Singh S, Naruka I, Singh A. Tamarind (*Tamarindus indica* L.). 2021;439-56.
- Ishaku G, Ardo Pariya B, Hayatuddeen A, Andrew F. Nutritional composition of

- Tamarindus indica* fruit pulp. Journal of Chemistry and Chemical Sciences. 2016;6:695-9.
28. Martin I. Fruits for the Future. Tamarind. (*Tamarindus indica* L.) By K. El-Siddeg, HPM Gunesana, BA Prasad, DKNP Pushpukumara, KVR Ramana, P. Vijayanand and JT Williams. Southampton, UK: Southampton Centre for Underutilized Crops (2006), pp. 188, available free on request to national scientists of developing countries. ISBN 854328599. Experimental Agriculture. 2007;43(3):407.
 29. El-Siddig K. Tamarind: *Tamarindus indica* L: Crops for the Future; 2006.
 30. Yusuf A, Mofio B, Ahmed A. Proximate and mineral composition of *Tamarindus indica* Linn 1753 seeds. Science world journal. 2007;2(1).
 31. Parvez SS, Parvez MM, Nishihara E, Gemma H, Fujii Y. *Tamarindus indica* L. leaf is a source of allelopathic substance. Plant Growth Regulation. 2003;40:107-15.
 32. Kumar S, Thombare P, Kale P. Tamarind: Value Added Products and its Health Benefits. 2020.
 33. Narina SS, Catanzaro CJ. Tamarind (*Tamarindus indica* L.), an underutilized fruit crop with potential nutritional value for cultivation in the United States of America: A review. Asian Food Science Journal. 2018;5(1):1-15.
 34. Raja MM, Poonguzhali S, Rao BV, Choudhury A, Devarajan A. A scientific evidence-based review of tamarind usage in Indian Folklore Medicine. Journal of Natural Remedies. 2022:347-62.
 35. Azhakiyamanavalan R, Vadivel E, editors. Dry land tamarind PKM-1, pre and post-release status. proceedings of national symposium on *Tamarindus indica*, Tirupathi, Andhra Pradesh; 1997.
 36. Bennet S, Durai A, Nicodemus A, Gireesan K, Nagarajan B, Varghees M, editors. Genetic improvement of *Tamarindus indica* L. for increased productivity. Proceedings of National Symposium on *Tamarindus indica* L Tirupati, Andhra Pradesh, India; 1997.
 37. Sharma S, Bhardwaj R. Tamarind—a suitable fruit crop for dry arid regions. Proc Nat Sym on *Tamarindus indica*. 1997;4-6.
 38. Menon JS, Asna A, Menon MV, Pooja A, Gopinath PP, Singh A. In situ characterization of tamarind (*Tamarindus indica* L.) fruit and spotting sweet tamarind types in Palakkad gap of Kerala. Plant Genetic Resources. 2023;21(2):166-73.
 39. Murthy K. Tamarind tree improvement works in Karnataka. MYFOREST. 1997; 33:457-64.
 40. Ravindran P. Advances in spices research: Agrobios (India); 2006.
 41. Meena VS, Gora JS, Singh A, Ram C, Meena NK, Roupael Y, et al. Underutilized fruit crops of Indian arid and semi-arid regions: Importance, conservation and utilization strategies. Horticulturae. 2022; 8(2):171.
 42. Farooqi AA, Sreeramu B, Srinivasappa K. Cultivation of spice crops: Universities press; 2005.
 43. Nair KP, Nair KP. Tamarind. Minor spices and condiments: Global economic potential. 2021;177-84.
 44. Tania C, Das M, Basanta T, Chatterjee R, Hnamte V, Chattopadhyay P. Assessment of tamarind (*Tamarindus indica* L.) varieties for growth, flowering, fruiting, yield and quality. International Journal of Current Microbiology and Applied Sciences. 2018;7(11):1708-13.
 45. Geetha V. Urigam tamarind-A unique variety. Spice India. 1995;8(8):8-9.
 46. Ndiaye NI, Qun CL, Nkoh JN. Importance of soil amendments with biochar and/or Arbuscular Mycorrhizal fungi to mitigate aluminum toxicity in tamarind (*Tamarindus indica* L.) on an acidic soil: A greenhouse study. Heliyon. 2022;8(2).
 47. Rao L, Joseph B, Sreemannarayana B. Growth and biomass production of some important multipurpose tree species on rainfed sandy loam soils. Indian forester. 2000;126(7):772-81.
 48. Kelly B, Cuny P. Afforestation using indigenous forest tree species on hydromorphic soil. Results of an experiment in southern Mali. Revue Forestière Française. 2000;52(5):453-66.
 49. Fandohan AB, Salako VK, Assogbadjo AE, Diallo BO, Van Damme P, Sinsin B. Effect of climatic conditions on flowering and fruiting of *Tamarindus indica* (Fabaceae). J Hortic For. 2015;7(8):186-92.
 50. Rachie KO. Tropical legumes: resources for the future. Tropical legumes: Resources for the future. 1979.
 51. Azad MS. Tamarindo— *Tamarindus indica*. 2018;403-12.
 52. Karale A, Kavlgud S, More T, editors. Studies of tamarind in Maharashtra. Proc Nat Symp on *Tamarindus indica* L,

- Tirupathi (AP), Organized by Forest Dept of AP, India; 1997.
53. Morton JF. Fruits of warm climates: JF Morton; 1987.
 54. Jøker D. *Tamarindus indica* L. Seed Leaflet-Danida Forest Seed Centre. 2000;(45).
 55. Muhammad S, Amusa N. Effects of sulphuric acid and hot water treatments on seed germination of tamarind (*Tamarindus indica* L). African journal of biotechnology. 2003;2(9):376-279.
 56. MacDonald I, Omonhinmin A, Ogboghodo I. Germination ecology of two savanna tree species, *Tamarindus indica* and *Prosopis africana*. Seed Technology. 2002;103-7.
 57. Parameswari K, Srimathi P, Maiarkodi K. Standardisation of dormancy breaking treatment in tamarind (*Tamarindus indica* L.) seeds. Legume Research-An International Journal. 2001;24(1):60-2.
 58. Yahia E, Salih N-E. Tamarind (*Tamarindus indica* L.). Postharvest biology and technology of tropical and subtropical fruits: Elsevier. 2011;442-58e.
 59. Pruthi JS. Minor spices and condiments: crop management and post-harvest technology. Minor spices and condiments: crop management and post-harvest technology; 2001.
 60. Purushotham K, Narasimharao S. Propagation of tamarind by veneer and soft-wood grafting. South Indian Horticulture. 1990;38(4).
 61. Anil M, Deshmukh U, Singh J, Nishad D, Kumar PPD. Studies on softwood grafting of tamarind (*Tamarindus indica* L.) under different growing conditions; 2022.
 62. Mowobi GG, Osuji C, Salisu A, Yahaya FM. In vitro regeneration of Tamarind (*Tamarindus indica* L.) explants. Journal of Environment and Life Sciences. 2016;1(1):26-31.
 63. Olomola D, Osunlaja O, Adekunle E, Kolawole I, Aguda S, Oyediran R. In-vitro propagation of *Tamarindus indica* L. through Seed Culture Technique towards Plantation Establishment. In-vitro. 2019;5(11).
 64. Pattepur SV, Mokashi A, Ajjappalavara P. Effect of cytokinins and auxin on shoot proliferation of cotyledonary nodes derived from axenic seedling of tamarind (*Tamarindus indica* L.). Asn J Hort. 2010;5(1):185-8.
 65. Simon O. Scaling Up the Production and Commercialization of Tamarind Fruit in Kenya: The Missing Value Chain Links; 2019.
 66. Yaacob O, Subhadrabandhu S. The production of economic fruits in South-East Asia: Oxford University Press; 1995.
 67. de Lima Neto AJ, Cavalcante LF, Nunes JC, de Luna Souto AG, Bezerra FTC, Cavalcante AG. Biometric variables and photosynthetic pigments in tamarind seedlings irrigated with saline water and biofertilizers. Semina: Ciências Agrárias. 2018;39(5):1909-20.
 68. Kundu S, Gajbhiye P, Srinivasarao C, Bheemaiah G. Effect of integrated nutrient management on yield attributes, yield, nutrient uptake and economics of growing maize in tamarind-based cropping system. Indian Journal of Dryland Agricultural Research and Development. 2009;24(1): 81-6.
 69. Mayavel A, Prasath CH, Muthuraj P, Nagarajan B. Effect of Manuring, Growth Regulators and Chemicals on Vegetative and Reproductive Characters of Seed Origin Plantation in Tamarind (*Tamarindus indica*). Int J Curr Microbiol App Sci. 2018;7(10):1821-6.
 70. Reena J, Bagyaraj D. Growth stimulation of *Tamarindus indica* by selected VA mycorrhizal fungi. World Journal of Microbiology and Biotechnology. 1990;6:59-63.
 71. Parthasarathy VA. PU, Kumar A. Spice Crops: Vol: II: Tree Spices. Today and Tomorrow, New Delhi. 2008.
 72. Das P. Spice crops production technology: Scientific Publishers; 2014.
 73. Nagarajan B, Nicodemus A, Mandal AK, Verma RK, Gireesan K, Mahadevan NP. Phenology and controlled pollination studies in Tamarind. Silvae Genetica. 1998;47:237-40.
 74. Hiwale S, Hiwale S. Tamarind (*Tamarindus indica* L.). Sustainable Horticulture in Semiarid Dry Lands. 2015;197-212.
 75. Purselglove J. Tropical Crops: Dicotyledons Longman Group Ltd. Burnt Hill; 1987.
 76. Wadnerkar D. Pest management in tamarind. Production and management of rainfed fruit crops. 2006;131-4.
 77. Joshi RC, David BV. Nematode, mite and insect pests of tamarind: A review. Agriculture for Development. 2018;35:52-63.
 78. Babu A, Sheeba G, Sen A, Ignacimuthu S. Record of slug caterpillar on *Ixora*

- parviflora Vahl. Insect Environment. 2000;6(1).
79. Gatan MG. Identification and assessment of major phytopathogenic fungi of tamarind leaves in Central Luzon, Philippines. Philippines (October 7, 2021); 2021.
80. Muhammad S, Suberu H, Amusa N, Agaji M. The effect of soil amendment with sawdust and rice husks on the growth and incidence of seedling blight of *Tamarindus indica* Linn caused by *Macrophomina phaseolina* and *Rhizoctonia solani*. Moor J Agric Res. 2001;2:40-6.
81. Nybe E. Spices: New India Publishing; 2007.
82. Prasath CH, Balasubramanian A, Radhakrishnan S. Effect of canopy management (pruning) on fruit yield in tamarind plantation at Harur Taluk, Dharmapuri, Tamil Nadu, India. Indian Journal of Plant and Soil. 2019;6(1).
83. Balasubramanian A, Sudha P, Prasath CH, Sangareswari M, Radhakrishnan S, Suresh K. Tamarind science and technology: Scientific Publishers; 2018.
84. Rao AS, Kumar AA, Ramana M. Tamarind seed processing and by-products. Agric Eng Int: CIGR J. 2015;17:200-4.
85. Feng L, Liu H, Li L, Wang X, Kitazawa H, Guo Y. Improving the property of a reproducible bioplastic film of glutenin and its application in retarding senescence of postharvest *Agaricus bisporus*. Food Bioscience. 2022;48:101796.
86. ARUDRA AK. Tamarind seed processing and by-products. Agricultural Engineering International: CIGR Journal. 2015; 17(2).
87. Rajkumar P, Deepa J, Rani CI. Mathematical Modeling on Drying of Tamarind Fruits; 2023.
88. Prabhu K, Teli M. Eco-dyeing using *Tamarindus indica* L. seed coat tannin as a natural mordant for textiles with antibacterial activity. Journal of Saudi Chemical Society. 2014;18(6):864-72.
89. Malviya R, Sundram S, Fuloria S, Subramaniyan V, Sathasivam KV, Azad AK, et al. Evaluation and characterization of tamarind gum polysaccharide: the biopolymer. Polymers. 2021;13(18):3023.
90. Donno D, Mellano MG, Hassani S, De Biaggi M, Riondato I, Gamba G, et al. Assessing nutritional traits and phytochemical composition of artisan jams produced in comoros islands: Using indigenous fruits with high health-impact as an example of biodiversity integration and food security in rural development. Molecules. 2018;23(10):2707.
91. Lima RS, Bolini HMA. Impact of non-nutritive sweeteners on the sensory profile and acceptance of a functional tamarind beverage. Journal of Food and Nutrition Research. 2020;8(1):26-32.
92. Bhusari SN, Kumar P, editors. Antioxidant activities of spray dried tamarind pulp powder as affected by carrier type and their addition rate. International Conference on Food, Biological and Medical Sciences (FBMS-2014), Bangkok (Thailand); 2014.
93. Israel KS, Murthy C, Patil B, Hosamani R. Value addition of tamarind products in Karnataka. Journal of Pharmacognosy and Phytochemistry. 2019;8(6):726-30.
94. Kiranmai E, Rajeswari K, Sharma S, Maheswari KU. Standardization and development of tamarind candy by blending with mango pulp. Journal of Pharmacognosy and Phytochemistry. 2018; 7(2):2042-7.
95. Latoch A, Libera J. Quality and safety of pork steak marinated in fermented dairy products and sous-vide cooked. Sustainability. 2019;11(20):5644.
96. Maia JL, Wurlitzer NJ, Lima JR, Borges MdF, Maia MdO, Damião BS, Oliveira LdS. Quality and storage stability of tamarind juice. Brazilian Journal of Food Technology. 2021;24.
97. Okudu H, Umoh E, Ojinnaka M, Chianakwalam O. Nutritional, functional and sensory attributes of jam from velvet tamarind pulp. African Journal of Food Science. 2017;11(2):44-9.
98. Kumar B, Patel S, Mishra N. A Review on-post harvest handling practices of tamarind (*Tamarindus indica* L.) fruits.

© 2023 Vasanthkumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/107126>