



Foliar Application of Liquid Micronutrient Formulation for Improving Quality of Panner Grapes

Yogaswathy D. ^{a++}, D. Jegadeeswari ^{a#*}, S. Suganya ^{a†}
and C. Kavitha ^{b‡}

^a Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore - 641 003, India.

^b Department of Fruit Science, HC&RI, Tamil Nadu Agricultural University, Coimbatore - 641 003, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102766

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105557>

Original Research Article

Received: 17/06/2023

Accepted: 23/08/2023

Published: 25/08/2023

ABSTRACT

Micronutrient deficiencies play a significant role in impacting the yield and quality of grapes in vineyards. To address the deficiencies, a field study was conducted during the summer, of 2022 by using foliar micronutrient mixtures in three different concentrations (0.5%, 1.0%, and 1.5%), with one spray during pre-fruit set stage and twice at pre-fruit set/berry growth stages. The study aimed to investigate the effects of foliar micronutrient application on the biochemical quality characteristics of the Muscat Hamburg grapevine cultivar. The results of the study showed that the application of micronutrients spray showed a beneficial impact on all the yield parameters, such as number of

⁺⁺ PG Scholar;

[#] Professor (SS&AC);

[†] Associate Professor (SS&AC);

[‡] Associate Professor;

*Corresponding author: E-mail: jagadeeswari.d@tnau.ac.in;

bunches per vine (ranging from 28 to 44), the yield per vine (varied from 5.56 to 6.55 kg per vine), with overall yield range between 20.7 to 24.2 t ha⁻¹. The biochemical properties of the berries, including sugars (reducing, non-reducing, and total sugars), juice pH and content, TSS (Total Soluble Solids), Titrable acidity, and berry firmness, were found to be high in grapevines sprayed at 1.0% twice. Results showed that total sugars ranged from 15.2% to 10.7%, with a juice pH of 3.47 to 3.94, juice content of 83.5 to 95.5%, and TSS from 15.6 to 17.40° Brix. Regarding the Titrable acidity and firmness of the berries, the outcomes showed a range of 0.92% to 1.14% for acidity and 9.4 to 12.9 N for firmness. Findings from the study clearly showed that applying 1.0% micronutrients twice at fruit-set and berry stages increased yield and improved quality indices of Paneer grapes.

Keywords: Micronutrients; foliar spray; grape; fruit quality.

1. INTRODUCTION

Grape (*Vitis vinifera* L.) member of Vitaceae family, is one of the top-ranked fruit in the world due to its multifaceted uses and benefit. It has gained substantial importance due to its high nutritional content, flavor, varied uses, and higher profit to the farmers [1]. It is one of the most important commercial fruit crop, from temperate to tropical regions, covering more land than any other fruit and accounting for more than half of global fruit production [2]. Micronutrients are necessary ingredients for plant growth and production [3,4] and each of these component is crucial for plant growth and development. Poor soil conditions bind these nutrients, reducing their availability and absorption. In inadequate conditions, the extrinsic administration of these chemicals via an integrated method can efficiently manage plant vitality. These supplements account for the decreased delivery of nutrients from the soil at various phases of development [5].

Nonetheless, employing a well-suited and harmonized approach to applying both essential macro and micronutrients has demonstrated its feasibility and lasting effectiveness in bolstering crop productivity. This approach has also proven effective in raising micronutrient levels and enhancing yield and biochemical quality parameters [6]. Efficient micronutrient management, such as micronutrient soil and foliar delivery in a variety of soil-crop scenarios, could help alleviate micronutrient shortages in soils and crops, boosting agricultural output and crop quality [7]. To meet the requirement at critical stages, foliar application is gaining popularity because of several advantages, including its swift effects and the little amount of fertilizer required in the solution. Foliar fertilizers are more effective when there is a strong demand for nutrients, particularly whilst

soil supply and root uptake are limited [8]. With this insight, the current study is being conducted in order to investigate the impact of micronutrient foliar treatment on the yield and quality characteristics of grapes.

2. MATERIALS AND METHODS

2.1 Experimental Vineyard

The field experiment was carried out during the summer of 2022 under irrigated conditions in a farmer's vineyard located at Mathipalayam village of Thondamuthur block, Coimbatore district, Tamil Nadu. The precise coordinates of the location are 10° 56'12.8" N latitude and 76° 50'44.0" E longitude. Field initial soil sample was collected at a depth of 0-15 cm and analysed for various physical and chemical attributes. Texture of the soil was silty clay [9], with neutral in soil reaction (7.74) [10] low electrical conductivity (0.28dS m⁻¹) [10] and high organic carbon content (1.23 g kg⁻¹) [11]. Available nitrogen (N) status was low (236 kg ha⁻¹) [12], while available potassium (K) (198 kg ha⁻¹) [13] and phosphorus (P) (21 kg ha⁻¹) [14] were at medium levels. In terms of available micronutrients, all micronutrients were present in sufficient quantities except for iron and boron, which exhibited a deficiency at 3.41 and 0.42 mg kg⁻¹ respectively [15].

This experiment was conducted using Randomized Block Design (RBD) with seven treatments, as varied concentrations i.e., micronutrients (Fe, Zn, Mn, Cu, and B), with concentrations of 0, 0.5%, 1.0%, and 1.5% applied once and twice. The foliar spraying of the micronutrient combinations was carried out during the fruit set and berry growth stages. The standard package of practices for nutrient management (200:160:600 NPK g/vine) and viticulture was followed in the vineyard,

regardless of the treatment applied. Routine cultural practices were followed, berry was harvested, yield and quality parameters were determined.

2.2 Yield Parameters

The plants were grown until they reached full maturity, and the mature clusters were collected treatment wise. Various physical measurements were computed manually, encompassing metrics such as cluster count per vine, bunch weight (g), bunch length (cm), number of berries per cluster, berry diameter (mm), and berry weight (g) [16].

2.3 Biochemical Quality Analysis

A sample of berries was taken from each replicate's bunch and juice was extracted for estimating total soluble solids (TSS), pH, titratable acidity, berry firmness, reducing sugars, total sugars, and non-reducing sugars. The TSS of grape juice was determined with a hand-held refractometer and expressed in degrees Brix, while the pH was determined with a digital pH meter. Titratable acidity is determined by titrating the extracted sample of grape juice with an alkaline solution (such as 0.1N NaOH) with phenolphthalein as an indicator and expressed as a percentage of tartaric acid [17]. TSS/acid ratio was calculated by dividing TSS ($^{\circ}$ brix) by acidity (%). Berry firmness is determined using a standard penetrometer and expressed in Newton (N) [18]. The grape juice percentage was calculated, reducing, non-reducing, and total sugar content in the fruit was estimated using Fehling's solution, as reported by Ranganna [19]. The sugar-acid ratio was determined by dividing total sugar content by acidity.

2.4 Statistical Analysis

The data collected from the experiment were statistically analysed utilizing AGRESS software version 7.01. A significance level of $P < 0.05$ was employed, and Critical Difference (CD) values were computed for $P < 0.05$ whenever the "F" test yielded significant results [20].

3. RESULTS AND DISCUSSION

3.1 Yield Parameters

The impact of applying various levels of micronutrients to crops via their leaves was investigated, and the results were documented in Table 1. The foliar treatment of micronutrients

had a notable influence on the fruit yield such as the number of fruit bunches per plant, with a range between 28 to 44. Significantly higher bunch yield was recorded and the results were, 5.56 to 6.55 kg per plant, while the overall yield per hectare spanned from 20.7 to 24.2 t ha⁻¹. The higher yield per plant and per hectare was observed in 1.0% micronutrient spray twice, with a value of 6.55 kg per plant and 24.2 t ha⁻¹ respectively. This was closely followed by the application of 0.5% micronutrient spray twice, which yielded 6.30 kg per plant and fruit yield of 23.9 t ha⁻¹. In contrast, the lowest yield of at 5.56 kg and 20.7 t ha⁻¹ was recorded in the control with water spray alone. Our findings align with earlier research, which demonstrated that applying Fe, Zn, and B through foliar methods led to substantial enhancements in vine productivity, bunch weight, and grapevine quality [21]. In a recent study conducted by Saha *et al.* in 2023, it was found that simultaneous application of both Zn and B led to a noteworthy augmentation in crucial quality attributes of tomatoes. Furthermore, the influence of Fe was observed in direct aspects of fruit development, such as fruit set, retention percentage, bunch quantity, bunch length, berry count per bunch, berry diameter, bunch weight, compactness, berry weight, and overall vine yield [22] (Singh *et al.*, 2015).

3.2 Biochemical Quality Parameters

The quality attributes of grapes, such as juice content, pH, TSS, Titrable acidity, berry firmness, total sugars, reducing sugars, and non-reducing sugars, were notably affected by the application of micronutrient combination at varying doses. The juice content (%) ranged from 95.5% to 83.5%, with the highest juice content observed in the vines treated with 1.0% micronutrient spray twice (95.5%), followed by 1.5% micronutrient spray (93.0%). Conversely, the lowest juice content was recorded in the vines sprayed with water alone (83.5%). This difference in juice content can be attributed to the greater photosynthetic activity and biomass production in the vines, leading to an increased presence of metabolites. As the berries continue to grow and develop, significant amounts of water and other metabolites are transported into the berries. These results validate the conclusions drawn by previous studies [23,24] in the context of sweet oranges. The pH of the juice varied from 3.94 to 3.47, with the highest pH recorded in the vines treated with 1.0% micronutrient spray twice (3.94), followed by the ones treated with 0.5% micronutrient spray twice (3.84). Whereas, the lowest pH was observed in the control (3.47).

Table 1. Effect of foliar spraying of micronutrients on the yield of grapes

Treatments	Yield (kg vine ⁻¹)	Yield (t ha ⁻¹)
T ₁ - Control	5.56 ^b	20.7 ^d
T ₂ - 0.5% MN spray once	5.92 ^a	21.5 ^d
T ₃ - 1.0% MN spray once	6.18 ^a	23.9 ^{ab}
T ₄ - 1.5% MN spray once	5.29 ^{ab}	23.1 ^{bc}
T ₅ - 0.5% MN spray twice	6.30 ^a	23.9 ^{ab}
T ₆ - 1.0% MN spray twice	6.55 ^a	24.2 ^a
T ₇ - 1.5% MN spray twice	5.30 ^{ab}	22.5 ^c
SEd	0.66	0.41
CD (P=0.05)	1.39	0.86

*MN – Micronutrients. Mean separation was conducted using the LSD test. Treatments with different lowercase letters indicate significant differences at the $P < 0.05$ level

Fig. 1 shows the sugar contents (%) in the berries and demonstrates a direct relationship between the concentration of micronutrients and sugar content (%). The lowest amount of non-reducing sugars (3.22%) was evidenced in control with water spray alone without micronutrients, whereas the highest amount of non-reducing sugars (5.23%) was observed in grapevines treated with micronutrient foliar spray at 1.0% twice. Similar results were obtained for reducing sugars with the results varying from 9.74% to 7.31%.

The lowest amount of reducing sugars (7.31%) was noticed in control, while the application of micronutrients at various concentrations increased the concentration of reducing sugars in all the treatments, with statistically significant differences compared to the control. Furthermore, the total sugars in grapevines, ranged from 15.2% to 10.7%, showed a significant increase in concentration due to the application of micronutrients, compared to the control (10.7%). The grapevines treated with the micronutrients at 1.0% twice exhibited the highest level of total sugars (15.2%). The rise in sugar levels in grape berries was clearly associated with elevated chlorophyll contents and a higher rate of photosynthesis due to increased levels of Fe, B, and Zn [25]. Similar observations have been documented in apple and grapevine studies [21,26,27]. Moreover, previous reports also documented that foliar sprays of micronutrients have led to increased sugar contents in strawberry, pomegranate, and grape crops [28-30].

The concentration of quality parameters such as total soluble solids (TSS), Titrable acidity, and berry firmness is a crucial factor for assessing fruit quality (Table 3). In this investigation,

grapevines treated with a combination of micronutrients at 1.0% twice displayed the highest TSS content, measuring 17.4° Brix. As the foliar dose of these micronutrients increased from the control to various concentrations, the TSS content also increased, ranging from 17.4 to 15.6° Brix. In contrast, the lowest TSS was observed in control (15.6° Brix). The gradual rise in TSS could be due to prime role of B and indicates a direct correlation between B and TSS, possibly attributed due to significant role in photosynthesis, flower and fruit set characteristics [31,32]. These findings align with previous studies on grapevines, where foliar treatment of B significantly elevated TSS levels in grape berries. Additionally, Zn, as an essential micronutrient, aids in activating enzymes (fructose 1 and 6-bis phosphatase) which is crucial for biochemical reactions that contribute towards sugar synthesis and accumulation in fruits [33,34].

With regard to Titrable acidity of the juice, the results ranged from 0.92 to 1.14%. Similar to other biochemical parameters, in TA also micronutrient spray showed positive effect. Highest acidity (1.14%) was observed in grapevines that did not receive any supplemental micronutrients, while the application of micronutrients led to a significant reduction in acidity percentage. The lowest Titrable acidity (0.92%) was found in grapevines treated with a 1.0% micronutrient spray twice. These results are consistent with the earlier findings of Parthiban et al. (2021). Similarly, the studies conducted by Ali et al. [16] on grape "Flame Seedless" and Mostafa et al. [35] also reported that foliar application of boric acid progressively increased the total soluble solids (TSS) and reduced Titrable acidity in grapes, supporting the present observations.

Table 2. Influence of foliar micronutrients on the pH and juice content (%) of grape juice

Treatments	Juice pH	Juice content (%)
T ₁ - Control	3.47 ^c	83.5 ^d
T ₂ - 0.5% MN spray once	3.53 ^{bc}	85.0 ^{cd}
T ₃ - 1.0% MN spray once	3.72 ^{abc}	89.2 ^{bc}
T ₄ - 1.5% MN spray once	3.81 ^{ab}	93.0 ^{ab}
T ₅ - 0.5% MN spray twice	3.84 ^{ab}	90.5 ^{ab}
T ₆ - 1.0% MN spray twice	3.94 ^a	95.5 ^a
T ₇ - 1.5% MN spray twice	3.83 ^{ab}	93.0 ^{ab}
SEd	0.15	2.60
CD (P=0.05)	0.31	5.46

*MN – Micronutrients. Mean separation was conducted using the LSD test. Treatments with different lowercase letters indicate significant differences at the P < 0.05 level

Table 3. Impact of foliar nutrition of micronutrients on the TSS, Titrable acidity, and berry firmness

Treatments	TSS (° brix)	Titration acidity (%)	Berry firmness (N)
T ₁ - Control	15.6 ^d	1.14 ^d	9.4 ^e
T ₂ - 0.5% MN spray once	16.0 ^{cd}	1.06 ^{cd}	9.9 ^d
T ₃ - 1.0% MN spray once	16.7 ^b	0.94 ^{ab}	10.8 ^c
T ₄ - 1.5% MN spray once	16.1 ^c	0.99 ^{bc}	10.1 ^d
T ₅ - 0.5% MN spray twice	16.5 ^{bc}	0.95 ^{ab}	11.5 ^b
T ₆ - 1.0% MN spray twice	17.4 ^a	0.92 ^a	12.9 ^a
T ₇ - 1.5% MN spray twice	16.1 ^c	1.10 ^{de}	10.3 ^d
SEd	0.24	0.04	0.24
CD (P=0.05)	0.50	0.07	0.51

*MN – Micronutrients. Mean separation was conducted using the LSD test. Treatments with different lowercase letters indicate significant differences at the P < 0.05 level

The berry firmness exhibited significant variations among the treatments, ranging from 12.9 N to 9.40 N. Among all the treatments, 1.0% micronutrient spray twice resulted in the highest berry firmness (12.9 N), followed by the 0.5% micronutrient spray twice (10.5 N). Conversely, the lowest berry firmness was recorded in the water spray treatment (9.40 N). Fruits that had higher uptake of micronutrients demonstrated better firmness compared to the control, as these elements potentially contribute to strengthening pectin bonding and stabilizing the cell wall structure, involvement in metabolism and translocation as well. These findings are consistent with the observations made by Swathi et al. [32]. In addition, Mostafa et al. [36] reported that a combination of boric acid and potassium sulphate increased fruit firmness in Anna apple. Similarly, research by Yadav et al. [37] showed that foliar spraying of peach trees with 0.1% H₃BO₃ + 0.5% ZnSO₄.7H₂O + 0.5% FeSO₄.7H₂O led to an increase in fruit firmness in peach.

The sugar/acid ratio and TSS/acid ratio (Fig. 2) indicate the organoleptic taste of the fruit which

appeals to the consumers' acceptance and thus market preference. The sugar-acid ratio of the berries ranged from 16.6 to 9.42 wherein the highest was observed in the treatment sprayed with 1.0% micronutrient spray twice (16.6) followed by 0.5% spray twice (14.5). Similarly, higher TSS/acid ratio (19.0) was calculated from the vines sprayed with 1.0% foliar-sprayed micronutrient blend followed by 1.0% micronutrient spray once (17.8). Balanced fertilization particularly micronutrients helps the plant to grow and produce potential growth steadily without any interruption provided the environment favours. In the present study also significant results were noticed for all the parameters, the results of sugar/acid ratio and TSS/acid ratio are in accordance with the findings of Pal et al., [38] who recorded that foliar application of micronutrients improved the fruit quality in respect of a higher TSS/ acid ratio. Prabhu and Singaram [39] also observed with higher TSS/acid ratio in the Muscat grape with foliar application of 0.5% ZnSO₄ and 0.2% borax [40,41].




Fig. 1. The influence of foliar micronutrients on the sugar content (%) of grape juice




Fig. 2. The influence of foliar micronutrients on sugar-acid and TSS-acid ratio on grapes

4. CONCLUSION

The present study aimed to enhance the yield and quality of "Muscat Hamburg" table grapes through the foliar application of micronutrients (Fe, Zn, Mn, Cu, and B). Our discovery highlights that the combined utilization of micronutrients

serves as an efficient approach for enhancing grape fruit yield, yield-related characteristics, and their constituents. Increasing doses of micronutrients led to an increased number of bunches per vine, yield per vine, and yield per hectare. Among all treatments, the most effective was the foliar spraying of 1.0% micronutrient

spray twice, at the fruit set stage and at berry growth stages. Significantly higher fruit yield (24.2 t ha^{-1}) as well as improved quality attributes were recorded in foliar spraying of 1.0% micronutrient spray twice, at the fruit set stage and at berry growth stages. Similarly, the application of micronutrients had a positive impact on grape quality characteristics such as TSS, sugar content, and berry firmness (12.9 N) when compared to control without micronutrient spray (water alone). As the micronutrient spray increased, Titrable acidity in grapes decreased. Based on the findings, it was concluded that applying 1.0% micronutrient spray twice during fruit set and berry growth stages resulted in increased fruit yield (Paneer grape variety), along with enhanced quality parameters.

ACKNOWLEDGEMENTS

Authors gratefully acknowledge the Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore for providing lab and research facilities to conduct the research study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Imran M, Rauf A, Imran A, Nadeem M, Ahmad Z, Atif M, Waqar AB. Health benefits of grapes polyphenols. *J. Environ. Agric. Sci.* 2017;10:40–51.
2. Nache Gowda V, Keshava SA, Shyamamma S. Growth, yield and quality of bangalore blue grapes as influenced by foliar applied polyfeed and multi-K. In *International Symposium on Grape Production and Processing*. 2006;785:207-212.
3. Mahmood-ul-Hassan M, Akhtar M, Nabi G. Boron and zinc transport through intact columns of calcareous soils. *Pedosphere*. 2008;18:524–532.
4. Palmer CM, Guerinot ML. Facing the challenges of Cu, Fe and Zn homeostasis in plants. *Nature Chemical Biology*. 2009;5(5):333-340.
5. Keller M. Deficit irrigation and vine mineral nutrition. *Am. J. Enol. Vitic.* 2005;56:267–283.
6. Saha B, Saha S, Kumar J, Bairwa R. Nutritional enrichment of tomato (*Lycopersicon esculentum* L.) through zinc (Zn) and boron (B) fertilization. *Journal of Plant Nutrition*. 2023;46(9):2155-2166.
7. Shukla AK, Behera SK. Biofortification for overcoming zinc and iron malnutrition in Indian population: Current research status and way forward. *Indian Journal of Fertilisers*. 2020;16(12):1262-1276.
8. Fernández V, Sotiropoulos T, Brown PH. Foliar fertilization: scientific principles and field practices. *International Fertilizer Industry Association*; 2013.
9. Piper AM. A graphic procedure in the geochemical interpretation of water-analyses. *Eos, Transactions American Geophysical Union*. 1944;25(6): 914-928.
10. Jackson ML. *Soil chemical analysis: Advanced course*; 2005.
11. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science*. 1934;37(1):29-38.
12. Subbiah BV, Asija GL. A rapid method for the estimation of nitrogen in soil. *Current Science*. 1956;26(5):259-60.
13. Stanford G, English L. Use of the flame photometer in rapid soil tests for K and Ca; 1949.
14. Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate; 1954.
15. Lindsay R, Norvell WA. Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil science society of American Journal*. 1978;42(3):421-428.
16. Ali I, Wang X, Abbas WM, Hassan MU, Shafique M, Tareen MJ, et al. Quality responses of table grapes 'flame seedless' as effected by foliarly applied micronutrients. *Horticulturae*. 2021;7(11): 462.
17. Genanew T. Effect of postharvest treatments on storage behavior and quality of tomato fruits. *World Journal of Agricultural Sciences*. 2013;9(1):29-37.
18. Leng F, Wang C, Sun L, Li P, Cao J, Wang Y, et al. Effects of different treatments on physicochemical characteristics of 'kyoho' grapes during storage at low temperature. *Horticulturae*. 2022;8(2):94.
19. Ranganna S. *Hand Book of analysis and quality control for fruits and vegetables products*. 3, & edition, Tata McGraw-Hill publishing company Ltd. New Delhi. 1997:12-16.

20. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley & Sons; 1984.
21. Ali I, Wang X, Tareen MJ, Wattoo FM, Qayyum A, Hassan MU, Shafique M, Liaquat M, Asghar S, Hussain T, et al. Foliar application of salicylic acid at different phenological stages of peach fruit cv. 'flordaking' improves harvest quality and reduces chilling injury during low temperature storage. *Plants*. 2021;10:1981.
22. Nawaz H, Zubair M, Derawadan H. Interactive effects of nitrogen, phosphorus and zinc on growth and yield of tomato (*Solanum lycopersicum*). *African Journal of Agricultural Research*. 2012;7(26):3792-3769.
23. Ghosh SN, Besra KC. Effect of zinc, boron and iron spray on yield and fruit quality of sweet orange cv. mosambi grown under rainfed laterite soil. *Indian Agriculturist*. 2000;44(3/4):147-151.
24. Kulkarni NH. Effect of growth regulators and micronutrients on fruit drop, yield and quality in sweet orange (*Citrus sinensis* Osbeck). Unpublished Ph. D. Thesis, Marathwada Agricultural University, Parbhani, India; 2004.
25. Perveen R, Wang X, Jamil Y, Ali Q, Ali S, Zakaria MQ, Afzaal M, Kasana RA, Saleem MH, Fiaz S. Quantitative determination of the effects of He-Ne laser irradiation on seed thermodynamics, germination attributes and metabolites of safflower (*Carthamus tinctorius* L.) in relation with the activities of germination enzymes. *Agronomy*. 2021;11:1411.
26. Spinelli F, Fiori G, Noferini M, Sprocatti M, Costa G. Perspectives on the use of a seaweed extract to moderate the negative effects of alternate bearing in apple trees. *The Journal of Horticultural Science and Biotechnology*. 2009;84(6):131-137.
27. Khan SU, Wang X, Mehmood T, Latif S, Khan SU, Fiaz S, Qayyum A. Comparison of organic and inorganic mulching for weed suppression in wheat under rain-fed conditions of Haripur, Pakistan. *Agronomy*. 2021;11:1131.
28. Manaf A, Wang X, Tariq F, Jhanzab HM, Bibi Y, Sher A, Razzaq A, Fiaz S, Tanveer SK, Qayyum A. Antioxidant enzyme activities correlated with growth parameters of wheat sprayed with silver and gold nanoparticle suspensions. *Agronomy*. 2021;11:1494.
29. Wang X, Saleem MH, Parveen A, Mumtaz S, Hassan A, Adnan M, Fiaz S, Ali S, Iqbal Khan Z, Ali S, et al. Proximate composition and nutritive value of some leafy vegetables from Faisalabad, Pakistan. *Sustainability*. 2021;13:8444.
30. Naqve M, Wang X, Shahbaz M, Fiaz S, Naqvi W, Naseer M, Mahmood A, Ali H. Foliar spray of alpha-tocopherol modulates antioxidant potential of okra fruit under salt stress. *Plants*. 2021;10:1382.
31. Güneş A, Köse C, Turan M. Yield and mineral composition of grapevine (*Vitis vinifera* L. cv. Karaerik) as affected by boron management. *Turkish Journal of Agriculture and Forestry*. 2015;39(5):742-752.
32. Swathi AS, Jegadeeswari D, Chitdeshwari T, Kavitha C. Effect of foliar nutrition of calcium and boron on the yield and quality attributes of grape. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(3):3625-3629.
33. Bybordi A, Shabanov JA. Effects of the foliar application of magnesium and zinc on the yield and quality of three grape cultivars grown in the calcareous soils of Iran. *Notulae Scientia Biologicae*. 2010;2(2):81-86.
34. Nikkhah R, Nafar H, Rastgoo S, Dorostkar M. Effect of foliar application of boron and zinc on qualitative and quantitative fruit characteristics of grapevine (*Vitis vinifera* L.). *International Journal of Agriculture and Crop Sciences (IJACS)*. 2013;6(9):485-492.
35. Mostafa EAM, El-Shamma MS, Hagagg LF. Correction of B deficiency in grapevines of Bez-El-Anaza cultivar. *Eurasian J Agric Environ Sci*. 2006;1:301-305.
36. Mostafa MF, El-Boray MS, Iraqi MA, EL-Bary AA. Effect of potassium and boron application on yield, fruit quality and leaf mineral content of Anna apple trees. *J. Agric. Sci. Mansoura Univ*. 1999;24(9):4965-4977.
37. Yadav V, Singh PN, Yadav P. Effect of foliar fertilization of boron, zinc and iron on fruit growth and yield of low-chill peach cv. Sharbati. *International Journal of Scientific and Research Publications*. 2013;8(3):1-6.
38. Pal R, Ghosh SN. Influence of micronutrients on growth dynamics, fruit yield and quality of arka neelamani grape. *Indian Journal of Horticulture*. 2018;75(1): 149-152.

39. Prabu PC, Singaram P. Effect of micronutrients (zinc and boron) on growth and yield of grapes (*Vitis vinifera*) cv. Muscat. *Madras Agricultural Journal*. 2001;88:45-49.
40. Gopalswamy N, Rao VN. Effect of graded dose of boron on yield and quality of grapes Cv. Anab-e-shahi. *South Indian Hort*. 1972;20:41-49.
41. Manjit S, Mahital J, et al. Response of iron and zinc on vegetative and reproductive growth of strawberry (*Fragaria x Ananassa Duch.*) cv. Chandler. *Bangladesh Journal of Botany*. 2015;44(2):337-340.

© 2023 Yogaswathy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105557>