



# Evaluation of Traditional Rice Genotypes (*Oryza sativa* L.) for Seed Quality Traits under Aerobic Condition

Sura Venkata Rachana <sup>a</sup>, B. Basavaraja <sup>a\*</sup>,  
K. S. Shashidhara <sup>b</sup>, K. N. Prakash <sup>c</sup>, M. H. Manjunatha <sup>d</sup>,  
B. N. Radha <sup>a</sup> and D. C. Hanumanthappa <sup>e</sup>

<sup>a</sup> Department of Seed Science and Technology, University of Agricultural Sciences, GKVK, Bengaluru, 560 065, India.

<sup>b</sup> Department of Genetics and Plant Breeding, College of Agriculture, Hassan, 573225, India.

<sup>c</sup> Department of Agricultural Economics, University of Agricultural Sciences, GKVK, Bengaluru, 560 065, India.

<sup>d</sup> Department of Agrometeorology, University of Agricultural Sciences, GKVK, Bengaluru, 560 065, India.

<sup>e</sup> AICRP on Agroforestry, University of Agricultural Sciences, GKVK, Bengaluru, 560 065, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JEAI/2024/v46i42337

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113594>

Original Research Article

Received: 19/12/2023

Accepted: 26/02/2024

Published: 01/03/2024

## ABSTRACT

Freshly harvested seeds of 58 traditional rice genotypes and two checks (KRH-4 and Doddabyranellu grown under aerobic condition were subjected to seed quality evaluation. Among traditional rice genotypes lowest electrical conductivity (3.48 dScm<sup>-1</sup> g<sup>-1</sup>), highest germination percentage (99.67 %),  $\alpha$ -amylase activity (2.03 cm) and total dehydrogenase activity (1.123) was observed in Neralisale, whereas highest thousand seed weight (30.00 gm), mean seedling length

\*Corresponding author: E-mail: basavaraja.sst@gmail.com;

(37.83 cm), mean seedling dry weight (22.53 mg), seedling vigor index-I (3693) and II (2216) was noticed in Chinnaponni compared to other genotypes. Among checks, Doddabyranellu showed better seed quality traits. From this study it could be concluded that genotypes Chinnaponni and Neralisale are well-suited for aerobic condition, benefiting farmers facing irrigation constraints by ensuring improved seed quality.

*Keywords: Aerobic rice; seed quality; checks; traditional rice genotypes.*

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is one of the staple food crop and primary source of energy for over half of the world's population and is considered as "global grain". Rice is the primary food crop consumed by more than 3.5 billion people and is one of the world's leading cereals and grown under variety of climatic and soil conditions. Rice varieties are composed of carbohydrates (77.53 %), protein (12.35 %), fat (3.23%), ash (1.64 %) and supply 60 to 80 per cent calories of energy. In addition to this, it is a source of magnesium, thiamine, niacin, phosphorous, vitamin B<sub>6</sub>, zinc and copper. In rice production, India is the world's second largest producer after China.

Rice is the largest irrigated water consumer and is responsible for massive greenhouse gas (GHG) emissions which contribute to global warming. Traditional lowland rice production with constant floods requires a greater amount of water. However, rising water scarcity affects the viability of the irrigated rice production system, as well as food security and the livelihoods of rice producers and consumers. Given the rising scarcity of water, there is a need to develop alternative rice cultivation technologies that use less water. The name "Aerobic rice" was recently coined by IRRI to describe a new method of growing rice. Aerobic rice, also known as "upland rice" or "aerobic paddy" is a cultivation method which involves growing of rice plants in non-puddled, well drained fields or upland areas typically using less water rather than in traditional flooded paddy fields. It refers to the process of establishing the crop from dry seeds sown in the field rather than transplanting seedlings from nursery.

Traditional rice genotypes refer to the older, locally adapted rice varieties that have been cultivated by farmers for generations. These varieties are often well suited to specific agroclimatic conditions, local soil types and cultural preferences. They have played a crucial role in ensuring food security and livelihoods for many communities around the world. Traditional

cultivars, which are rich in nutritional and locally superior traits, hold the key to rice cultivation sustainability. These varieties are important in terms of sustainable production and land usage [6, 13, 14].

Numerous hybrids have been evaluated for their performance and seed quality in aerobic condition. The effect of water regime such as upland cultivation of traditional rice genotypes on the seed quality of rice has not been studied exhaustively. Therefore, the present experiment was conducted to evaluate the performance of traditional rice genotypes under aerobic condition for seed quality traits.

## 2. MATERIALS AND METHODS

The study was conducted in Department of Seed Science and Technology laboratory, University of Agricultural Sciences, GKVK, Bangalore during the year 2022-23. Freshly harvested seeds of 58 traditional rice genotypes and two checks (KRH-4 and Doddabyranellu grown under aerobic condition were subjected to seed quality evaluation. Seed quality parameters viz., thousand seed weight (gm), moisture content (%), seed germination (%), shoot length (cm), root length (cm), mean seedling length (cm), mean seedling dry weight (mg), seedling vigour index-I, seedling vigour index-II, electrical conductivity (dScm<sup>-1</sup>g<sup>-1</sup>), total dehydrogenase activity (A<sub>480nm</sub>) and α-amylase activity (cm) were recorded.

### 2.1 Thousand Seed Weight (gm)

1000 seeds are counted from the harvested plants of each replication and are weighed and expressed in grams.

### 2.2 Seed Moisture Content (%)

Three replicates of five grams of seed material are taken for determining the seed moisture content using hot air oven. The powdered seed material is placed in a weighed moisture cup. After removing the lid, moisture cups are placed in hot air oven maintained at 130°C for 2 ± 1 for

2 hours and the contents are allowed to dry. Then it is cooled in a desiccator for 30 min and weighed in an electronic balance along with metal cup and lid. The moisture content is worked out using the following formula and expressed as percentage [2].

M2- M3

Moisture content (%) =  $\frac{\quad}{\quad} \times 100$

M2 - M1

Where,

M1: Weight of the empty moisture cup

M2: Weight of the moisture cup + sample

before drying M3: Weight of the moisture cup + sample after drying

### 2.3 Seed Germination (%)

The germination test is conducted in the laboratory by using between paper method as per ISTA rules, for this the one hundred seeds from each genotype are drawn at random in three replicates and is carried out. Further rolled towels are incubated in a germination chamber maintained at  $25 \pm 1^\circ\text{C}$  and 90 per cent relative humidity. Germinated seedlings are evaluated on 14th day as final count and percentage germination is expressed based on number of normal seedlings present. [2].

### 2.4 Shoot Length (cm)

From the germination test, ten normal seedlings are selected randomly from replication of each genotype on the day of final count. The shoot length is measured and expressed in centimeters.

### 2.5 Root Length (cm)

From the germination test, ten normal seedlings are selected randomly from replication of each genotype on the day of final count. The root length is measured and expressed in centimeters.

### 2.6 Mean Seedling Length (cm)

From the germination test, ten normal seedlings are selected randomly from replication of each genotype on the day of final count. The seedling length is measured from tip of shoot to root tip and the mean length is calculated and expressed as seedling length in centimeters.

### 2.7 Mean Seedling Dry Weight (mg)

The normal seedlings are taken in butter paper after removing the cotyledons and dried in a hot-air oven maintained at  $85 \pm 1^\circ\text{C}$  for 24 hours. Then, seedlings were removed and allowed to cool in a dessicator for 30 minutes before weighing in an electronic balance. The average weight is calculated and expressed as seedling dry weight in mg per 10 seedlings.

### 2.8 Seedling Vigour Index – I

The seedling vigor index - I is computed by multiplying percentage germination and seedling length (cm) and expressed as a whole number [1].

Seedling vigour index - I = Germination (%) x Mean seedling length (cm)

### 2.9 Seedling Vigour Index – II

The seedling vigor index - II is computed by multiplying percentage germination and seedling dry weight (mg) and expressed as a whole number [1].

Seedling vigour index - II = Germination (%) x Seedling dry weight (mg)

### 2.10 Electrical Conductivity ( $\text{dScm}^{-1}\text{g}^{-1}$ )

Electrical conductivity will be measured [2]. Three replications of 25 seeds were weighed on an analytical balance and soaked in 25 ml of distilled water for 24 hours at  $25 \pm 10^\circ\text{C}$ . The electrical conductivity was measured using digital conductivity meter.

### 2.11 Total Dehydrogenase Activity ( $A_{480\text{nm}}$ )

The total dehydrogenase activity was measured by the method suggested by [5]. The seeds imbibed in Tz viability testing were washed thoroughly with distilled water and the red-coloured formazan from the stained regions was extracted by soaking with 5 ml of 2-methoxy ethanol for 8 h in an airtight container. The extract was decanted and the colour intensity was measured in Spectrophotometer (Model Mini Spec 17) at 480 nm with a suitable blank (Methoxy ethanol). The total dehydrogenase activity (TDH) was expressed as absorbance.

## 2.12 A-Amylase Activity (cm)

The  $\alpha$ -amylase activity was estimated in semi quantitatively by adopting „Beld-Rock Model“. The overnight soaked seeds of three replications were dehusked and cut into two equal halves. The embryo was placed equidistantly on a sterilized petri plates containing 2 per cent agar and one per cent potato starch medium in such a way that embryo portion of seed should come on to top and incubated for 24 hours at  $25 \pm 1^\circ\text{C}$  [4]. After incubation, the seeds were removed and iodine solution was poured on to the agar medium. After 30 minutes, the observation on

clear „halo zone“ formed around the embryos indicating the  $\alpha$ -amylase activity. The diameter of  $\alpha$ -amylase activity was measured and expressed in centimeters.

## 3. RESULTS AND DISCUSSION

### 3.1 1000 Seed Weight (g)

The data pertaining to thousand seed weight of traditional rice genotypes cultivated under aerobic condition are depicted in Table 1.

**Table 1. Seed moisture content, seed germination of traditional rice (*Oryza sativa* L.) genotypes cultivated under aerobic condition**

Genotype number	Genotypes	Thousand seed weight (g)	Seed moisture content (%)	Seed germination (%)
1	KRH-4	18.49	11.39	88.00
2	Doddabyranellu	27.00	11.25	97.67
1	Tvrs mukkanasanna	13.81	11.09	97.33
2	Salemsanna -2	14.96	11.50	96.00
3	Belegamibasmati	18.22	12.21	97.67
4	Kadupeple	20.50	11.13	87.67
5	Jeenugodu	21.53	11.62	97.33
6	Neralisale	28.01	11.65	99.67
7	Shashtikashale	20.79	11.14	94.00
8	Tulasiya	14.57	11.52	96.00
9	Rajamudikempu	12.05	11.74	91.00
10	Mysoorumalligae	19.89	11.43	97.33
11	Doddayasandha	24.03	11.56	95.00
12	HMT-1	25.59	11.66	95.67
13	Nirgulabattha	15.29	11.79	96.33
14	RNR	11.01	11.89	95.00
15	Saraswati -1	17.46	12.32	93.67
16	Salemsanna -1	12.58	11.85	94.67
17	Nirgasamba	18.25	12.17	97.33
18	Pattubattha	14.54	12.25	92.33
19	Mutthinassanna	22.24	12.33	95.67
20	Jupodli	22.66	12.46	97.00
21	Gudabattha- 2	21.83	11.38	97.67
22	Ratnasagara	20.87	11.20	93.67
23	Bangaragandu	22.62	12.17	96.00
24	Navali	20.70	11.75	85.00
25	Bidigekannappa	19.52	11.03	92.00
26	Kalainamik -1	11.82	12.18	92.33
27	Doddabattha	21.25	11.41	91.00
28	Ambemohar	19.80	11.76	82.33
29	MSN 99	12.85	11.93	94.67
30	Imbangidda	16.63	11.63	92.00
31	Mobikar	14.91	11.75	93.33
32	Kagesele	18.02	12.19	95.67
33	Belejadu	13.21	11.42	93.33
34	Kuhadisamba	18.34	11.28	97.67
35	Kundipullan	18.69	11.55	96.00

Genotype number	Genotypes	Thousand seed weight (g)	Seed moisture content (%)	Seed germination (%)
36	Sannakki	10.90	12.26	93.67
37	Vannekattu	20.69	11.35	97.00
38	AP 21	16.39	11.16	91.33
39	Thogarsi	27.13	11.97	97.33
40	Ranangali	23.35	11.88	88.33
41	Santhethal	10.49	12.29	96.67
42	SLR 3449	19.88	11.31	82.67
43	Basmati	19.27	11.24	86.33
44	Kyasalai	24.00	11.09	95.00
45	Valakkasanna	19.26	11.29	95.67
46	KMP 225	20.67	11.03	96.00
47	Beppigedoddabyra	22.05	11.87	86.00
48	Karibasmati	18.61	11.07	94.00
49	Sannallibattha	17.05	11.42	89.00
50	Chinnaponni	30.00	12.15	98.33
51	Bebbanna	16.24	11.74	90.67
52	GK-1	20.43	11.82	95.67
53	Sannakempakki	13.43	12.05	97.33
54	AP-6	19.31	11.43	97.67
55	Jeerigesanna	10.16	12.04	96.67
56	Jawahar	19.33	11.42	98.67
57	Ratbad	20.94	11.57	96.33
58	Bellineralu	20.89	11.83	91.67
	<b>Mean</b>	18.75	11.66	93.95
	<b>SEm±</b>	0.49	0.37	1.84
	<b>CD (p=0.05)</b>	1.39	NS	5.16
	<b>CV (%)</b>	4.60	5.55	3.40

\*NS: Non Significant

Different rice genotypes and checks resulted in significant differences in thousand seed weight. Genotype Chinnaponni recorded highest thousand seed weight (30.00 g), while lowest was recorded in genotype Jeeerigesanna (10.16 g). Among checks Doddabyranellu recorded higher test weight (27.00 g).

Higher test weight might be due to better translocation of photosynthates and bolder seeds [12]. Similar variation in thousand seed weight of different rice genotypes grown under aerobic condition was reported by Shantaet al [8].

### 3.2 Seed Moisture Content (%)

The data on seed moisture content of traditional rice genotypes cultivated under aerobic condition are depicted in Table 1.

Seed moisture content of different genotypes and checks resulted in non significant difference for seed moisture content. Minimum moisture content was recorded in genotype

Bidigekannappa (11.03 %), while maximum moisture content was recorded in genotype Jupodli (12.46 %). Among checks Doddabyranellu recorded minimum moisture content (11.25 %).

### 3.3 Seed Germination (%)

The data pertaining to seed germination of traditional rice genotypes cultivated under aerobic condition are depicted in Table 1.

Seed germination varied significantly among different genotypes and checks produced under aerobic condition. Genotype Neralisale recorded highest seed germination (99.67 %) followed by Jawahar (98.67 %), Chinnaponni (97.67 %), while genotype Ambemohar recorded lowest seed germination (82.33 %). Among checks Doddabyranellu recorded highest seed germination (97.67 %).

The primary goal of every seed production activity is to achieve a high level of germination. Rice seeds should possess at least 80 per cent germination as per minimum seed certification

standards. In this study all the rice genotypes registered germination level of above 80 %. Highest germination percentage might be due to genotype character, water use efficiency and nutrient uptake, this will leads to increase in storage food and it may be utilized during germination and plant developmental stages [11,7]. Similar higher germination percentage of rice genotypes cultivated under aerobic

condition were reported by Madhukar et al [9].

### 3.4 Shoot Length (cm)

The data pertaining to shoot length of traditional rice genotypes cultivated under aerobic condition are depicted in Table 2.

**Table 2. Shoot length, root length, mean seedling length of traditional rice (*Oryza sativa* L.) genotypes cultivated under aerobic condition**

Genotype number	Genotypes	Shoot length (cm)	Root length (cm)	Mean seedling length(cm)
1	KRH-4	9.76	18.18	27.94
2	Doddabyranellu	8.41	15.42	23.83
1	Tvrs mukkanasanna	8.16	15.27	23.43
2	Salemsanna -2	10.38	19.55	29.94
3	Belegamibasmati	5.37	10.64	16.01
4	Kadupeple	9.60	17.04	26.64
5	Jeenugodu	12.70	17.83	30.53
6	Neralisale	10.70	21.70	32.40
7	Shashtikashale	8.08	15.42	23.51
8	Tulasiya	8.00	9.32	17.32
9	Rajamudikempu	7.86	16.22	24.08
10	Mysoorumalligae	10.06	16.31	26.37
11	Doddayasandha	9.63	17.53	27.16
12	HMT-1	12.28	14.74	27.02
13	Nirgulabattha	9.29	14.14	23.44
14	RNR	8.12	13.06	21.19
15	Saraswati -1	7.13	16.47	23.61
16	Salemsanna -1	10.76	13.25	24.02
17	Nirgasamba	10.87	16.43	27.30
18	Pattubattha	11.75	19.38	31.13
19	Mutthinasanna	7.48	10.79	18.27
20	Jupodli	13.24	19.79	33.03
21	Gudabattha- 2	11.97	17.55	29.52
22	Ratnasagara	9.82	18.36	28.19
23	Bangaragandu	15.15	19.74	34.89
24	Navali	9.54	15.21	24.75
25	Bidigekannappa	9.57	16.74	26.31
26	Kalainamik -1	8.44	11.49	19.93
27	Doddabattha	12.24	19.50	31.74
28	Ambemohar	13.12	18.05	31.17
29	MSN 99	7.73	15.62	23.36
30	Imbangidda	9.20	13.81	23.01
31	Mobikar	6.40	10.54	16.94
32	Kagesele	9.81	16.35	26.17
33	Belejadu	9.17	18.24	27.42
34	Kuhadisamba	7.58	15.58	23.20
35	Kundipullan	11.83	20.53	32.36
36	Sannakki	7.27	15.33	22.60
37	Vannekattu	12.03	17.68	29.71
38	AP 21	10.24	15.37	25.61
39	Thogarsi	10.61	13.28	23.89
40	Ranangali	12.36	16.93	29.29

Genotype number	Genotypes	Shoot length (cm)	Root length (cm)	Mean seedling length(cm)
41	Santhethal	6.91	12.17	19.08
42	SLR 3449	10.25	14.57	24.82
43	Basmati	10.14	13.53	23.67
44	Kyasalai	10.59	17.45	28.04
45	Valakkasanna	8.68	17.86	26.54
46	KMP 225	9.04	14.54	23.58
47	Beppigedoddabyra	13.61	22.02	35.63
48	Karibasmati	9.34	16.46	25.80
49	Sannalibattha	7.57	14.45	22.02
50	Chinnaponni	15.06	22.49	37.83
51	Bebbanna	9.13	12.73	21.87
52	GK-1	9.73	18.22	27.95
53	Sannakempakki	7.15	11.06	18.22
54	AP-6	12.17	17.44	29.62
55	Jeerigesanna	7.34	12.25	19.59
56	Jawahar	12.45	15.87	28.33
57	Ratbad	8.61	17.39	26.01
58	Bellineralu	9.05	11.36	20.42
	<b>Mean</b>	9.84	15.94	25.78
	<b>SEm±</b>	0.31	0.40	0.55
	<b>CD (p=0.05)</b>	0.89	1.12	1.56
	<b>CV (%)</b>	5.62	4.35	3.75

Shoot length varied significantly among different genotypes and checks produced under aerobic condition. Genotype Bangaragandu recorded highest shoot length (15.15 cm) followed by Chinnaponni (15.06 cm), while lowest was recorded by genotype Belegamibasmati (5.35 cm). Among checks KRH-4 recorded highest shoot length (9.76 cm).

### 3.5 Root Length (cm)

The data pertaining to root length of traditional rice genotypes cultivated under aerobic condition are depicted in Table 2.

Root length varied significantly among different genotypes and checks produced under aerobic condition. Genotype Chinnaponni recorded highest root length (22.49 cm) followed by Beppigedoddabyra (22.02 cm), while lowest was recorded by genotype Tulasiya (9.32 cm). Among checks KRH-4 recorded highest root length (18.18 cm).

### 3.6 Mean Seedling Length (cm)

The data on mean seedling length of traditional rice genotypes cultivated under aerobic condition are depicted in Table 2.

Mean seedling length varied significantly among different genotypes and checks produced under

aerobic condition. Genotype Chinnaponni recorded highest mean seedling length (37.83 cm) followed by Beppigedoddabyra (35.63 cm), while lowest was recorded by genotype Belegamibasmati (16.01 cm). Among checks KRH-4 recorded highest mean seedling length (27.94 cm).

The higher mean seedling length in Chinnaponni is due to higher shoot and root length.

### 3.7 Mean Seedling Dry Weight (mg)

The data pertaining to mean seedling dry weight of traditional rice genotypes cultivated under aerobic condition are depicted in Table 3.

Mean seedling dry weight varied significantly among different genotypes and checks produced under aerobic condition. Genotype Chinnaponni recorded maximum seedling dry weight (22.53 mg) followed by Bangaragandu (22.32 mg), while genotype Belegamibasmati recorded lowest mean seedling dry weight (5.22 mg). Among checks KRH-4 recorded maximum seedling dry weight (10.63 mg).

### 3.8 Seedling Vigour Index-I

The data pertaining to seedling vigour index-I of traditional rice genotypes cultivated under aerobic condition are depicted in Table 3.

Seedling vigour index-I varied significantly among different genotypes and checks produced under aerobic condition. Genotype Chinnaponni recorded seedling vigour index-I (3693) followed by Bangaragandu (3348), while genotype Belegamibasmati (1564) recorded lowest seedling vigour index-I. Among checks KRH-4 recorded maximum seedling vigour index-I (2445).

The highest seedling vigour index-I is due to highest germination percentage and mean seedling length.

### 3.9 Seedling Vigour Index-II

The data pertaining to seedling vigour index-II of traditional rice genotypes cultivated

under aerobic condition are depicted in Table 3.

Seedling vigour index-II varied significantly among different genotypes and checks produced under aerobic condition. Genotype Chinnaponni recorded seedling vigour index-II (2216) followed by Bangaragandu (2143), while genotype Belegamibasmati recorded lowest seedling vigour index-II (510). Among checks Doddabyranellu recorded maximum seedling vigour index-II (1003).

The highest seedling vigour index-II is due to highest germination percentage and mean seedling dry weight.

**Table 3. Mean seedling dry weight, seedling vigour index-I, mean seedling vigour index-II of traditional rice (*Oryza sativa* L.) genotypes cultivated under aerobic condition**

Genotype number	Genotypes	Mean seedling dry weight (mg)	Seedling vigour index-I	Seedling vigour index-II
1	KRH- 4	10.63	2445	935
2	Doddabyranellu	10.27	2327	1003
1	Tvrs mukkanasanna	10.78	2281	1049
2	Salemsanna -2	13.03	2875	1251
3	Belegamibasmati	5.22	1564	510
4	Kadupeple	7.81	2331	686
5	Jeenugodu	13.53	2972	1317
6	Neralisale	19.81	3229	1974
7	Shastikashale	8.50	2210	799
8	Tulasiya	6.70	1663	643
9	Rajamudikempu	13.74	2191	1250
10	Mysoorumalligae	14.17	2566	1379
11	Doddayasandha	15.97	2582	1516
12	HMT-1	16.26	2585	1558
13	Nirgulabattha	13.53	2257	1303
14	RNR	12.61	2313	1200
15	Saraswati -1	10.05	2211	941
16	Salemsanna -1	9.44	2274	893
17	Nirgasamba	8.93	2657	869
18	Pattubattha	19.48	2873	1799
19	Mutthinasanna	8.20	1748	784
20	Jupodli	19.73	3205	1914
21	Gudabattha- 2	19.04	2883	1859
22	Ratnasagara	17.78	2639	1663
23	Bangaragandu	22.32	3348	2143
24	Navali	16.75	2106	1421
25	Bidigekannappa	15.06	2418	1387
26	Kalainamik -1	8.10	1840	748
27	Doddabattha	17.95	2891	1633
28	Ambemohar	17.20	2560	1415
29	MSN 99	15.93	2212	1506
30	Imbangidda	9.84	2119	904
31	Mobikar	5.83	1579	544



Genotype number	Genotypes	Mean seedling dry weight (mg)	Seedling vigour index-I	Seedling vigour index-II
32	Kagesele	14.42	2505	1379
33	Belejadu	10.13	2557	946
34	Kuhadisamba	10.22	2262	998
35	Kundipullan	18.88	3107	1812
36	Sannakki	12.68	2116	1186
37	Vannekattu	16.80	2884	1630
38	AP 21	13.78	2334	1259
39	Thogarsi	11.93	2325	1161
40	Ranangali	12.34	2589	1090
41	Santhethal	10.51	1844	1016
42	SLR 3449	13.79	2051	1138
43	Basmati	13.38	2044	1154
44	Kyasalai	12.83	2664	1219
45	Valakkasanna	14.86	2541	1421
46	KMP 225	15.74	2264	1511
47	Beppigedoddabyra	21.63	3063	1857
48	Karibasmati	11.39	2426	1071
49	Sannalibattha	9.91	1962	886
50	Chinnaponni	22.53	3693	2216
51	Bebbanna	8.41	1980	762
52	GK-1	16.62	2674	1590
53	Sannakempakki	6.84	1773	667
54	AP-6	17.14	2893	1674
55	Jeerigesanna	8.10	1894	782
56	Jawahar	10.15	2795	1001
57	Ratbad	9.34	2505	900
58	Bellineralu	6.95	1872	636
	<b>Mean</b>	13.09	2426	1229
	<b>SEm±</b>	0.35	80.50	37.54
	<b>CD (p=0.05)</b>	0.99	225	105.13
	<b>CV (%)</b>	4.68	5.74	5.28

### 3.10 Electrical Conductivity (dScm<sup>-1</sup>g<sup>-1</sup>)

The data pertaining to electrical conductivity (dScm<sup>-1</sup> g<sup>-1</sup>) of traditional rice genotypes cultivated under aerobic condition are depicted in Table 4.

Electrical conductivity varied significantly among different genotypes and checks produced under aerobic condition. Genotype Neralisale recorded lower electrical conductivity (3.48 dScm<sup>-1</sup> g<sup>-1</sup>) followed by Jawahar (3.56 dScm<sup>-1</sup>g<sup>-1</sup>), Chinnaponni (3.63 dScm<sup>-1</sup>g<sup>-1</sup>), while genotype Ambemohar recorded higher electrical conductivity (10.48 dScm<sup>-1</sup>g<sup>-1</sup>). Among checks Doddabyranellu recorded lower electrical conductivity (3.83 dScm<sup>-1</sup>g<sup>-1</sup>).

The EC of seed leachate is sensitive index of seed quality, which shows negative association with other seed quality [3, 15, 16, 17]. Electrical conductivity of the seed leachate was greatly influenced by the method of cultivation. This might be due to less number of leachetes ,

increased amount reactive oxygen species and more number of living cells and ion exchange is in dual acceptable condition between plant and soil in case of aerobically developed genotype [11, 19]. Similar results were reported by [9, 10, 18].

### 3.11 Total Dehydrogenase Activity (A<sub>480nm</sub>)

The data pertaining to total dehydrogenase activity of traditional rice genotypes cultivated under aerobic condition are depicted in Table 4.

Total dehydrogenase activity varied significantly among different genotypes and checks produced under aerobic condition. Genotype Neralisale recorded highest total dehydrogenase activity (1.123) followed by Jawahar (1.103), while genotype Ambemohar recorded lowest total dehydrogenase activity (0.283). Among checks Doddabyranellu recorded highest total dehydrogenase activity (0.727).

The highest total dehydrogenase activity may be due to high vigour seeds having more number of active and living cells which is present in the seeds of aerobically grown genotypes and these will contribute to good germination and growth. Similar results of highest total dehydrogenase activity in rice genotypes grown under aerobic condition were reported by Sannagoudar and Murthy [11, 20].

### 3.12 A-Amylase Activity (cm)

The data pertaining to  $\alpha$ -amylase activity of traditional rice genotypes cultivated under aerobic condition are depicted in Table 4.

$\alpha$ -amylase activity varied significantly among different genotypes and checks produced under aerobic condition. Genotype Neralisale recorded highest  $\alpha$ -amylase activity (2.03 cm) followed by Chinnaponni (1.87 cm), while genotype Ambemohar (0.57 cm) recorded lowest  $\alpha$ -amylase activity. Among checks Doddabyranellu recorded highest  $\alpha$ -amylase activity (1.49 cm).

Higher amylase activity indicated that mobilization of reserve food materials and make available for the root and shoot growth hence the cultivars showing higher amylase activity resulted in higher germination [10, 21].

**Table 4. Electrical conductivity ( $\text{dScm}^{-1} \text{g}^{-1}$ ), total dehydrogenase activity ( $A_{480\text{nm}}$ ),  $\alpha$ -amylase activity (cm) of traditional rice (*Oryza sativa* L.) genotypes cultivated under aerobic condition**

Genotype number	Genotypes	Electrical conductivity ( $\text{dScm}^{-1} \text{g}^{-1}$ )	Total dehydrogenase activity ( $A_{480\text{nm}}$ )	$\alpha$ -amylase activity (cm)
1	KRH- 4	5.01	0.389	0.72
2	Doddabyranellu	3.83	0.727	1.49
1	Tvrs mukkanasanna	3.93	0.682	1.50
2	Salemsanna -2	4.04	0.523	1.24
3	Belegamibasmati	3.80	0.711	1.36
4	Kadupeple	6.95	0.366	0.62
5	Jeenugodu	3.74	0.603	1.30
6	Neralisale	3.41	1.123	2.03
7	Shashtikashale	4.31	0.418	1.16
8	Tulasiya	4.01	0.511	1.20
9	Rajamudikempu	4.78	0.376	0.84
10	Mysoorumalligae	3.48	0.634	1.51
11	Doddayasandha	4.08	0.421	1.14
12	HMT-1	4.04	0.475	1.17
13	Nnirgulabattha	4.11	0.504	1.27
14	RNR	4.05	0.467	1.08
15	Saraswati -1	4.25	0.342	1.11
16	Salemsanna -1	4.18	0.431	1.07
17	Nirgasamba	3.91	0.694	1.46
18	Pattubattha	4.45	0.395	1.02
19	Mutthinasanna	3.84	0.456	1.14
20	Jupodli	3.97	0.623	1.35
21	Gudabattha- 2	3.83	0.701	1.66
22	Ratnasagara	4.76	0.322	1.04
23	Bangaragandu	3.79	0.687	1.18
24	Navali	9.24	0.341	0.61
25	Bidigekannappa	4.73	0.384	0.89
26	Kalainamik -1	4.66	0.370	0.86
27	Doddabattha	4.85	0.357	0.82

Genotype number	Genotypes	Electrical conductivity (dScm <sup>-1</sup> g <sup>-1</sup> )	Total dehydrogenase activity (A <sub>480nm</sub> )	α-amylase activity (cm)
28	Ambemohar	10.48	0.283	0.57
29	MSN 99	4.12	0.408	1.09
30	Imbangidda	4.60	0.355	0.84
31	Mobikar	4.27	0.388	0.96
32	Kagesele	3.93	0.526	1.17
33	Belejadu	4.38	0.467	0.94
34	Kuhadisamba	3.71	0.908	1.61
35	Kundipullan	3.80	0.444	1.53
36	Sannakki	4.41	0.407	0.81
37	Vannekattu	4.01	0.697	1.32
38	AP 21	5.17	0.601	0.87
39	Thogarsi	3.95	0.801	1.58
40	Ranangali	4.97	0.397	0.74
41	Santhethal	3.93	0.511	1.28
42	SLR 3449	9.88	0.296	0.61
43	Basmati	8.80	0.326	0.77
44	Kyasalai	4.07	0.478	1.11
45	Valakkasanna	4.09	0.501	1.17
46	KMP 225	3.80	0.483	1.19
47	Beppigedoddabyra	7.91	0.367	0.74
48	Karibasmati	3.95	0.445	1.09
49	Sannalibattha	4.84	0.401	0.77
50	Chinnaponni	3.63	1.014	1.87
51	Bebbanna	4.96	0.415	0.79
52	GK-1	3.83	0.482	1.16
53	Sannakempakki	3.75	0.748	1.43
54	AP-6	3.92	0.881	1.62
55	Jeerigesanna	4.05	0.526	1.57
56	Jawahar	3.56	1.103	1.83
57	Ratbad	4.16	0.503	1.30
58	Bellinalu	4.81	0.377	0.96
	<b>Mean</b>	9.12	0.526	1.15
	<b>SEm±</b>	4.64	0.017	0.037
	<b>CD (p=0.05)</b>	0.42	0.049	0.10
	<b>CV (%)</b>	5.68	5.80	5.58

#### 4. CONCLUSION

From this study it could be concluded that genotypes Chinnaponni and Neralisale are well-suited for aerobic condition, benefiting farmers facing irrigation constraints by ensuring improved seed quality.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Krishnakutty J, Blakeney M, Raju RK, and Siddique KH. Sustainability of traditional rice cultivation in Kerala, India-A Socio-Economic Analysis. Sustainability. 2021; 13(2):980.
2. Sannagoudar MS, Patil RH, Rajanna GA, Ghosh A, Singh AK, Halli HM, Khandibagur V, Kumar S and Kumar RV. Rising temperature coupled with reduced rainfall will adversely affect yield of kharif sorghum

- genotypes. Current Science. 2023;124(8):921.
3. Biradar B, Jayadev HM, Halli HM and Sannagoudar MS. Mapping of soil test-based spatial fertilizer recommendations for paddy and maize using GIS, GPS and STCR approaches in a micro-watershed of Karnataka, India. Current Science. 2023; (00113891):124(10).
  4. Anonymous. International Rules for Seed Testing (ISTA). Seed Science and Technology. 2021;24:307-335.
  5. Abdul Baki AA and Anderson JD. Vigor determination in soybean seed by multiple criteria. Crop Science. 1973; 13(6):630-633.
  6. Kittock DL and Law AG. Relationship of seedling vigour to respiration and tetrazolium chloride reduction by germinating wheat seeds. Agronomy Journal.1968;60:286- 288.
  7. Kalavathi D, Ananthakalaiselvi A and Vijaya J. Storage potential of parental lines and hybrid seeds of rice. Seed Research.1999;27(2):146-148.
  8. Shanta CM, Ramesha YM, Krishnamurthy D, Umesh MR and Bhanuvally M. Growth and yield response of direct seeded rice (*Oryza sativa* L.) to dates of sowing and varieties. IJCS. 2020;8(4):295-299.
  9. Madhukar P, Raju CS, Senguttuvel P and Vanisree S. Evaluation and selection of promising rice (*Oryza sativa* L.) genotypes under augmented block design in aerobic system. International Journal of Pure & Applied Bioscience. 2017;5(4):1358- 1367.
  10. Ranjitha HP, Paramesh R, Shashidhar HE, Amruta N and Kiran SP. Effect of water regimes on seed quality parameters of rice grown under aerobic and wetland conditions. Journal of Applied and Natural Science. 2016; 8(3):1546 – 1550.
  11. Sannagoudar MS and Murthy KK. Growth and Yield of Maize (*Zea mays* L.) as Influenced by Planting Geometry and Nutrient Management in Maize Based Intercropping. Mysore Journal of Agricultural Sciences. 2018;52(2):278-284.
  12. Parashivamurthy, Rajendra Prasad R, Siddaraju S and Lakshmi J. Study on the effect of promising varieties and methods of cultivation on growth, seed yield and quality parameters in rice (*Oryza sativa* L.), Mysore Journal of Agricultural Sciences. 2012;46(3):529-537.
  13. Hibbard RP and Miller JV. Biochemical studies on seed viability, measurement of conductance and reduction. Plant Physiology. 1928;3:335-352.
  14. Baradwal H, Ghosh A, Kumar A, Singh PD, Sannagoudar MS, Ahamad S, Jha P, Singh AK, Bhattacharyya R, Manna MC and Kumar S. Ecological restoration of degraded lands with alternate land use systems improves soil functionality in semiarid tropical India. Land Degradation & Development. 2022; 33(7): 1076-1087.
  15. Sannagoudar MS, Murthy KK, Ghosh A, Singh AK, Gupta G, Halli HM and Kumar RV. Comparative efficacy of leguminous intercrops and weed management practices on nutrient uptake, productivity and profitability of maize based intercropping system. Legume Research. 2021;1(6): 10-18805.
  16. Gupta G, Dhar S, Dass A, Sharma VK, Shukla L, Singh R, Kumar A, Kumar A, Jinger D, Kumar D and Sannagoudar MS. Assessment of bio-inoculants-mediated nutrient management in terms of productivity, profitability and nutrient harvest index of pigeon pea–wheat cropping system in India. Journal of Plant Nutrition. 2020;43(19):2911-2928.
  17. Rajanna GA, Dass A, Singh VK, Choudhary AK, Paramesh V, Babu S, Upadhyay PK, Sannagoudar MS, Ajay BC and Reddy KV. Energy and carbon budgeting in a soybean–wheat system in different tillage, irrigation and fertilizer management practices in South-Asian semi-arid agroecology. European Journal of Agronomy. 2023; 148:126877.
  18. Radha, R. Effect of aerobic cultivation on seed quality, milling and cooking quality in rice, M.Sc., Thesis, University of Agricultural Sciences, Bangalore; 2005.
  19. Sannagoudar MS, Murthy KK, Nagaraju, Rajanna GA, Ghosh A, Singh AK, Gupta G and Kumar RV. Influence of weed management practices in maize (*Zea mays*) based intercropping system. The Indian Journal of Agricultural Sciences. 2021;91(7): 1067-1071.
  20. Sannagoudar MS, Patil RH, Kumar RV, Singh AK, Ghosh A and Halli HM.

- Simulated impacts of rise in temperature on kharif sorghum genotypes in Northern Transitional Zone of Karnataka, India. Cereal Research Communications. 2020;48, 113-120.
21. Gupta G, Dhar S, Kumar A, Choudhary AK, Dass A, Sharma VK, Shukla L, Upadhyay PK, Das A, Jinger D. and Rajpoot SK and Sannagoudar MS. Microbes-mediated integrated nutrient management for improved rhizo-modulation, pigeonpea productivity, and soil bio-fertility in a semi-arid agro-ecology. Frontiers in Microbiology. 2022;13.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/113594>