



# Evaluating Particulate Matter (PM<sub>2.5</sub> and PM<sub>10</sub>) Concentrations in Educational Institutions within South City Corporation, Dhaka, Bangladesh

Ahmad Kamruzzaman Majumder <sup>a,b,++\*</sup>, Tareq <sup>b#</sup>,  
Marziah Rahman <sup>a†</sup>, Rasheduzzaman Majumder <sup>a‡</sup>  
and Kaspia Rahman Tanima <sup>a^</sup>

<sup>a</sup> Center for Atmospheric Pollution Studies (CAPS), Dhaka-1217, Bangladesh.

<sup>b</sup> Department of Environmental Science, Stamford University Bangladesh, Dhaka-1217, Bangladesh.

## Authors' contributions

This work was carried out in collaboration among all authors. Authors AKM and MR conceptualized the study, performed the methodology, did investigation, wrote and prepared the original draft of the manuscript. Authors AKM and RM supervised the study. Authors AKM, MR, KRT and RM wrote, reviewed and edited the manuscript. Author Tareq prepared the initial draft. Authors MR and KRT did data curation and formal analysis. Author MR did data visualization. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ajarr/2024/v18i9746>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122319>

Original Research Article

Received: 26/06/2024  
Accepted: 31/08/2024  
Published: 10/09/2024

<sup>++</sup>Professor and Chairman;

<sup>#</sup>Post Graduate Student;

<sup>†</sup>Scientific Officer;

<sup>‡</sup>Director;

<sup>^</sup>Research Associate;

<sup>\*</sup>Corresponding author: Email: [dk@stamforduniversity.edu.bd](mailto:dk@stamforduniversity.edu.bd);

**Cite as:** Majumder, Ahmad Kamruzzaman, Tareq, Marziah Rahman, Rasheduzzaman Majumder, and Kaspia Rahman Tanima. 2024. "Evaluating Particulate Matter (PM<sub>2.5</sub> and PM<sub>10</sub>) Concentrations in Educational Institutions Within South City Corporation, Dhaka, Bangladesh". *Asian Journal of Advanced Research and Reports* 18 (9):221-32. <https://doi.org/10.9734/ajarr/2024/v18i9746>.

## ABSTRACT

Dhaka is widely recognized as the most polluted city in Bangladesh due to its high levels of particulate matters. Most of the educational buildings in Dhaka South City Corporation are situated alongside the main road. The objective of this study is twofold. First and foremost, it aims to identify the status of air pollution within educational institutions located in the Dhaka South City Corporation. Secondly, it seeks to assess the relationships among different zones and different air quality parameters, specifically focusing on particulate matter levels of  $PM_{2.5}$  and  $PM_{10}$ . This study was conducted in 37 locations of Dhaka metropolitan, by using Aeroqual s500. This research study is found that the average concentrations of  $PM_{2.5}$  and  $PM_{10}$  in all the selected places of the study area were 68.82 and 113.95  $\mu\text{g}/\text{m}^3$  respectively. The average concentration of particulate matter ( $PM_{2.5}$  and  $PM_{10}$ ) in all the zones had been observed to be more than one times higher than the acceptable level of concentration set by the National Ambient Air Quality Standard (NAAQS) levels. It had been measured that the average value of  $PM_{2.5}/PM_{10}$  was 60.41%. Additionally, the study demonstrated that as the P values are less than 0.05, the concentrations of  $PM_{2.5}$  may vary significantly between and among zones. The outcome of this research is arranged in descending order based on the average concentration of  $PM_{2.5}$  which follows as Zone-5 (Chak Bazar, Bangshal, Kotwali) > Zone-2 (Dhanmondi) > Zone-6 (Sutrapur, Gendaria, Jatra Bari) > Zone-3 (Ramna, Shahbagh) > Zone-1 (Hazaribagh) > Zone-4 (Lalbagh, New Market) > Zone-7 (Kadamtali).

**Keywords:** Air pollution; particulate matter; educational institution; Dhaka South City corporation; Bangladesh.

## 1. INTRODUCTION

Air pollution has risen to the status of a major environmental health concern in many nations throughout the globe. Human exposure is the most obvious consequence of pollution, but the hidden cost to society from these contaminants' impacts on people's health and the ecosystem is also significant [1]. Particulate Matter has a harmful effect not only on human health but also on the environment and can raise respiratory problems and global warming, respectively [2]. However, the inventory of air pollutants is declining in Europe and the USA, while in Asia, the situation is worsening in real-time [3]. Bangladesh has the foulest air in South Asian countries and each year 21% of all deaths in the country are caused due to a long-time exposure of high PM contain air [4]. Dhaka, the most populated city in the world and the capital of Bangladesh was identified as the fifth most polluted capital city in the world [5]. The level of air pollution in Dhaka city is now a quite concerning issue, with the AQI concentrations ranging from "Unhealthy" to "Extremely Unhealthy" for two-thirds of the year [6]. The city has grown increasingly crowded during the previous ten years. The land use with the highest average concentrations was the commercial area followed by residential, mixed, industrial, and sensitive area in Dhaka city [7]. The municipal infrastructure is being built, locals are building high-rise residential and commercial

developments to accommodate the city's growing population [8] which helps to increase air pollution. The massive and uncontrolled increase of emissions from industrial activities and transportation specifically, personal vehicles in Dhaka city deteriorates the air quality [9]. These two parts use a great amount of fuel that resulted in significant increases in emissions as well as economic losses [10]. Vulnerable groups like school going children are the major exposures and sufferers of PM because their immune and whole-body systems are under developing stage at this age [11]. In addition to having an adverse effect on children's physical and mental growth, air pollution aggravates respiratory diseases including asthma and seasonal allergic rhinitis (SAR), better known as hay fever, which is most frequent in schoolchildren [12,13]. The proximity of educational institutions to major roads and busy traffic areas exacerbates the exposure of students and staff members to harmful pollutants. The levels of  $PM_{2.5}$  near schools in Dhaka City exceeded the World Health Organization's (WHO) recommended guidelines, posing a significant health risk to the vulnerable population, especially children [14].

### 1.1 Objectives

The objectives of the Study are:

1. To identify the status of air pollution in educational institution of Dhaka South City Corporation, Dhaka.

- To assess the relationship among Zones and parameters ( $PM_{2.5}$  and  $PM_{10}$ ).

- Chak Bazar, Bangshal and Kotwali resided in Zone-(5), after that Sutrapur, Gendaria and Jatra Bari area involves in Zone-(6) and lastly Kadamtali area; Zone-(7).

## 2. METHODOLOGY

### 2.1 Study Area

The study was conducted in Dhaka South city corporation. The monitoring location was nearest to the road with no other primary pollutant sources nearby. We divided the study area into seven zones based on thanas of Dhaka South City Corporation area according to Dhaka City Thana Map of Bangladesh which include in total thirty-six educational institution locations. The zones are divided as following, Hazaribagh area; Zone-(1), Dhanmondi area; Zone-(2), Ramna and Shahbagh area includes in Zone-(3), than Lalbagh and New Market area consisted in Zone-

### 2.2 Data Collection

Aeroqual portable monitors (Model: S500) were employed to collect air quality data. These monitors were positioned horizontally on tripods at a height of approximately 1.5 meters above ground level. The monitors were deliberately positioned alongside the road near the educational institution, away from other major sources of pollution, in order to obtain precise measurements. The Aeroqual monitors were configured to measure and record the average concentrations of particulate matter (PM) for a period of 1 minute. In order to commence




Fig. 1. Study area (Dhaka South City corporation and data collection locations point)



**Fig. 2. Portable air quality monitor**

the monitoring process, the device was activated by pressing and holding the power button until the screen became operational. Prior to commencing data collection, the sensors necessitated a 3-minute warm-up period to eradicate any impurities. A distinct Location ID was allocated to each monitoring site for the purpose of identification.

The sensor autonomously gathered and stored air quality data following the initialization period. Data retrieval was expedited by establishing a connection between the device and a personal computer using a USB cable. The data can be readily downloaded and exported to Excel format using the provided software.

### **2.3 Data Processing & Interpretation of Results**

The collected data were processed using IBM SPSS V20 and MS Excel 2020. These tools were employed for data input, analysis, and visualization. A variety of graphs, tables, diagrams, and box-whisker plots were created in order to comprehend the properties of the data. Descriptive statistics were utilized to evaluate the spread of each land use characteristic, while ANOVA tests were employed to ascertain the statistical significance of the results. The air quality data is displayed in a thorough manner using an array of graphs, charts, and tables.

## **3. RESULTS AND DISCUSSION**

### **3.1 Status of Air Pollution Studies in Educational Institutions Area of South Dhaka City Corporation**

The status of air pollution in educational institution zones area of South Dhaka City Corporation were shown in Fig. 3 (a, b, c, d, e, f, g and h). While there should always be less pollution in sensitive areas, in Dhaka city, there

was a significant amount of pollution in these areas [7]. Study found that in zone 1 (Fig. 3 (a)) out of four places in Hajaribag area, the three highest average concentration of  $PM_{2.5}$  and  $PM_{10}$  were in front of Shahid Sheikh rasel Govt. High School, outside of Hajaribag Model Town High School and Hajaribag Ideal School & College with the concentration of 73 and 146  $\mu g/m^3$ ; 71 and 96  $\mu g/m^3$  and 34 and 103  $\mu g/m^3$  respectively. On the other hand, the least  $PM_{2.5}$  concentration was found in front of Hajaribag Girl's High School (31  $\mu g/m^3$ ) and  $PM_{10}$  concentration was found outside of Hajaribag Model Town High School (96  $\mu g/m^3$ ). It also found that the most polluted place was 1.12 folds higher than NAAQS level and 5.84 and 2.92 folds higher than WHO standard. Where, the Bangladesh NAAQS level is 65  $\mu g/m^3$  set by the Department of Environment (DoE) for concentrations of  $PM_{2.5}$ . Also, the Air Quality Standard (24-hour) set by the WHO for  $PM_{2.5}$  and  $PM_{10}$  are 25 and 50  $\mu g/m^3$  respectively. The primary cause of the severe dust pollution in the sensitive area of Dhaka city is vehicle emission [15] Fig. 3 (b) illustrated that among four places in Dhanmondi area, the three highest average concentration of  $PM_{2.5}$  was in front of Dhaka City College (108  $\mu g/m^3$ ), outside of Daffodil University (101  $\mu g/m^3$ ) and Stamford College (84  $\mu g/m^3$ ). Comparatively less polluted place was in front of Dhaka College (77  $\mu g/m^3$ ). The average concentration of  $PM_{10}$  found higher outside of Daffodil University (206  $\mu g/m^3$ ), Dhaka City College (116  $\mu g/m^3$ ) and Stamford College (105  $\mu g/m^3$ ) respectively and least polluted place was in front of Dhaka College (91  $\mu g/m^3$ ). The study also indicated that the most polluted place were 1.66 and 1.36 times higher than the national standards. The study estimated that  $PM_{2.5}$  was found in  $PM_{10}$  in Zone-2 was 76.69%. The comparison of the average concentration of  $PM_{2.5}$  and  $PM_{10}$  of zone-3 area was demonstrated through Fig. 3 (c). Within seven

places in Ramna and Shahbagh area, the three highest average concentration of  $PM_{2.5}$  was University of Dhaka (DU) area ( $80 \mu\text{g}/\text{m}^3$ ), outside of BTCL Ideal School and College ( $73 \mu\text{g}/\text{m}^3$ ) and Sher-E-Bangla School & College ( $56 \mu\text{g}/\text{m}^3$ ) and the least contaminated place was in front of Ibrahim Medical College ( $32 \mu\text{g}/\text{m}^3$ ). The average concentration of  $PM_{10}$  found higher outside of BTCL Ideal School and College ( $152 \mu\text{g}/\text{m}^3$ ), University of Dhaka (DU) ( $121 \mu\text{g}/\text{m}^3$ ) and in front of Siddeswari Girls College ( $109 \mu\text{g}/\text{m}^3$ ) respectively and comparatively least contaminated place was in front of Ibrahim Medical College ( $52 \mu\text{g}/\text{m}^3$ ). The most polluted place were 1.23 and 1.01 times higher than the NAAQS levels.

The Fig. 3 (d) presented that out of four places of zone-4 in Lalbagh and New Market area, the highest average concentration of  $PM_{2.5}$  were in front of Madrasha Rahmania Hafijia ( $68 \mu\text{g}/\text{m}^3$ ) and outside of Govt. laboratory High School ( $66 \mu\text{g}/\text{m}^3$ ) and comparatively less contaminated place was Dhaka Ideal High School ( $27 \mu\text{g}/\text{m}^3$ ). On the other hand, the average concentration of  $PM_{10}$  found higher outside of Govt. laboratory High School ( $133 \mu\text{g}/\text{m}^3$ ), Madrasha Rahmania Hafijia ( $113 \mu\text{g}/\text{m}^3$ ) and in front of Dhaka Ideal High School ( $70 \mu\text{g}/\text{m}^3$ ) respectively. The least contaminated place was Lalbagh Govt. Model School & College ( $58 \mu\text{g}/\text{m}^3$ ). Study observed that the most polluted place was 1.05 times higher than the NAAQS and 2.72 and 2.66 times higher than WHO standard. The study estimated that the ratio of  $PM_{2.5}/PM_{10}$  was 50.02% in Zone-4.

In Fig. 3 (e) zone-5 indicated that in Chak Bazar, Bangshal and Kotwali area, out of six places the highest average concentration of  $PM_{2.5}$  and  $PM_{10}$  were in front of Haybot Nagar Govt. Primary School ( $232$  and  $232 \mu\text{g}/\text{m}^3$ ), outside of Armanitola Govt. Primary School ( $120$  and  $186 \mu\text{g}/\text{m}^3$ ) and Suritola Model Govt. Primary School ( $117$  and  $119 \mu\text{g}/\text{m}^3$ ) respectively. Moreover, the least contaminated place was outside of Darul Huda Madrasa with  $PM_{2.5}$  and  $PM_{10}$  concentration of  $50.00$  and  $88.00 \mu\text{g}/\text{m}^3$  respectively. However, study found that the most polluted places were 3.60 and 1.55 folds higher than the NAAQS level.

The study estimated that in Zone-5, 73.45%  $PM_{2.5}$  was found into  $PM_{10}$ . Study found that in Sutrapur, Gendaria and Jatra Bari area Zone-6, the maximum average concentration of  $PM_{2.5}$

was in front of Central Women's College ( $91 \mu\text{g}/\text{m}^3$ ), outside of Narindra Govt. High School ( $79 \mu\text{g}/\text{m}^3$ ) and Dhaka Mahanagar Mohila College ( $74 \mu\text{g}/\text{m}^3$ ) and the average concentration of  $PM_{10}$  found higher outside of Narindra Govt. High School ( $151 \mu\text{g}/\text{m}^3$ ), Dhaka Mahanagar Mohila College ( $142 \mu\text{g}/\text{m}^3$ ) and in front of Central Women's College ( $140 \mu\text{g}/\text{m}^3$ ) respectively.

Furthermore, the least concentration of  $PM_{2.5}$  and  $PM_{10}$  was found in front of Salimullah College ( $45$  and  $73 \mu\text{g}/\text{m}^3$ ). In addition to the study found that the most polluted place were 3.60 and 1.55 times higher than the NAAQS level. Study found in zone-7 (Kadamtali area), the most average concentration of  $PM_{2.5}$  was in front of Muradpur Govt. Primary & High School ( $55 \mu\text{g}/\text{m}^3$ ), outside of Bangladesh bank Adarsha High School ( $45 \mu\text{g}/\text{m}^3$ ) and relatively less contaminated place was Syampur Govt. Primary School and in front of Shyampur Ideal High School ( $44 \mu\text{g}/\text{m}^3$ ). However, the average concentration of  $PM_{10}$  found higher outside of Bangladesh bank Adarsha High School ( $143 \mu\text{g}/\text{m}^3$ ), in front of Syampur Govt. Primary School ( $87 \mu\text{g}/\text{m}^3$ ) and Muradpur Govt. Primary & High School ( $71 \mu\text{g}/\text{m}^3$ ) respectively. Comparatively less contaminated places less polluted places were in front of Muradpur Govt. Primary & High School ( $71 \mu\text{g}/\text{m}^3$ ). Moreover, the most polluted place were 2.2 and 2.86 times higher than WHO standard. The study estimated that the ratio of  $PM_{2.5}/PM_{10}$  was 53.63% in Zone-7. Finally Fig 3 (h) illustrates the comparison among of the average concentration of  $PM_{2.5}$  and  $PM_{10}$  of seven zones in Dhaka South City Corporation. The average concentration of  $PM_{2.5}$  was higher in the Chak Bazar, Bangshal and Kotwali area (Zone-5), Dhanmondi area (Zone-2) and Ramna and Shahbagh area (Zone-3) with the values of  $116.00$ ,  $92.50$  and  $69.38 \mu\text{g}/\text{m}^3$  respectively. It was also observed that the average concentration of  $PM_{10}$  was higher in the Chak Bazar, Bangshal and Kotwali area (Zone-5), and Dhanmondi area (Zone-2) with the values of  $154.50$ , and  $129.50 \mu\text{g}/\text{m}^3$  respectively with the highest concentration of  $PM_{2.5}$  and  $PM_{10}$  in the Zone-5. It was also noticed that the concentrations of  $PM_{2.5}$  and  $PM_{10}$  found in the most polluted Zone were 1.78 and 1.03 times higher than the national standards. Moreover, the average concentration of  $PM_{2.5}$  ( $47.00 \mu\text{g}/\text{m}^3$ ) was found to be least in Kadamtali area (Zone-7) and  $PM_{10}$  ( $93.50 \mu\text{g}/\text{m}^3$ ) were found to be least in Lalbagh and New Market area (Zone-4).






Fig. 3. Comparison of particulate matters of educational institutions in Dhaka South City Corporation Area Based on Zones

Children in Dhaka cities experience respiratory issues as a result of air pollution [15]. Children's brain structure has been changed by particulate matter, which has an impact on cognitive performance [16].

### 3.2 Dispersion of the Educational Institutions in Dhaka South City Corporation Area Based on Zones

Table 1 presents the descriptive statistics for  $PM_{2.5}$  and  $PM_{10}$  across seven different zones. For  $PM_{2.5}$ , the highest concentration range was observed in Zone-5 at  $182 \mu\text{g}/\text{m}^3$ , followed by Zone-2 at  $74 \mu\text{g}/\text{m}^3$ , while the lowest was in Zone-7 at  $11 \mu\text{g}/\text{m}^3$ . The minimum concentration of  $PM_{2.5}$  across all zones was  $27 \mu\text{g}/\text{m}^3$  in Zone-4, and the maximum was  $232 \mu\text{g}/\text{m}^3$  in Zone-5. The highest mean  $PM_{2.5}$  concentration was in Zone-5 at  $116.00 \mu\text{g}/\text{m}^3$ , and the lowest was in Zone-7 at  $47.00 \mu\text{g}/\text{m}^3$ . Zone-5 also had the highest standard deviation of  $62.78 \mu\text{g}/\text{m}^3$ , whereas Zone-7 had the lowest at  $5.35 \mu\text{g}/\text{m}^3$ . The coefficient of variation for  $PM_{2.5}$  was highest in Zone-5 at 54.12% and lowest in Zone-7 at 11.39%. For  $PM_{10}$ , Zone-5 exhibited the highest range at  $144 \mu\text{g}/\text{m}^3$ , followed by Zone-2 at  $115 \mu\text{g}/\text{m}^3$ , and the lowest range was in Zone-1 at  $50 \mu\text{g}/\text{m}^3$ . The minimum  $PM_{10}$  concentration was  $52 \mu\text{g}/\text{m}^3$  in Zone-3, with the maximum again in Zone-5 at  $232 \mu\text{g}/\text{m}^3$ . The highest mean  $PM_{10}$  concentration was found in Zone-5 at  $154.50 \mu\text{g}/\text{m}^3$ , and the lowest was in Zone-4 at  $93.50 \mu\text{g}/\text{m}^3$ . The standard deviation was highest in Zone-2 at  $52.02 \mu\text{g}/\text{m}^3$  and lowest in Zone-1 at  $22.20 \mu\text{g}/\text{m}^3$ . The coefficient of variation for  $PM_{10}$  was highest in Zone-2 at 40.17% and lowest in Zone-1 at 19.47%.

The whisker box plot shows the average of  $PM_{2.5}$  and  $PM_{10}$  concentrations in seven zones in Fig. 4. Following whisker box plot of  $PM_{2.5}$  revealed that Zone-5 had the highest dispersion with extremely negative skewed distribution which contradicted with descriptive statistics where Zone-5 showed highest values along with one lower outlier; followed by Zone-4 and Zone-1 with normally skewed values with normally distributed values. This episode was found in front of Dhaka College. Moderate distribution was seen in Zone-2 with normal skewness and Zone-3 with positive skewness. These situations were found upper outlier in front of Haybot Nagar Govt. Primary School and lower outlier found outside of Darul Huda Madrasa in Zone-5. Less moderate distribution was seen in Zone-6 with

negatively skewed values. Zone-7 had the most clustered concentration with extremely positive distributed values.

This whisker box plot of  $PM_{10}$  revealed that the highest concentration that was seen in Zone-5 with positively skewed distribution followed by Zone-2, Zone-4 and Zone-6 with positive to normal skewed values. Moderate distribution was seen in Zone-7 with positively distributed values. Less Moderate distribution was seen in Zone-1 with extremely positive skewness, which was found in front of Shahid Sheikh Rasel Govt. High School. Enforcing stricter vehicle emissions laws and promoting the use of public transportation are essential to addressing the particulate matter problem in the Dhaka Metropolitan Area [7].

### 3.3 Significance Test

Table 2 of Analysis of variance (ANOVA) is performed to find whether the changes in the concentration of all the parameters between and within land uses are significant. The F values were calculated to be 4.052 for  $PM_{2.5}$  and 1.840 for  $PM_{10}$  where P value of  $PM_{2.5}$  and  $PM_{10}$  were 0.004 and 0.125. The following table shows that the concentrations of  $PM_{2.5}$  change significantly between and within in the land uses as the P values are less than 0.05. However,  $PM_{10}$  do not change significantly as the P values are greater than 0.05.

### 3.4 Land Use Based Cluster Analysis

Fig. 5 illustrates the dendrogram plots resulting from cluster analyses of  $PM_{2.5}$  and  $PM_{10}$  concentrations, with Z-score normalization applied. The  $PM_{2.5}$  analysis identified four distinct clusters. The first cluster includes Zone-4, Zone-7, Zone-1, and Zone-3. This cluster merges with the second cluster, which comprises Zone-6, at an approximate distance of 10. Subsequently, the combined group joins the third cluster, which includes Zone-2. Finally, the fourth cluster, consisting of Zone-5, merges with the third cluster to form a single cohesive cluster at an approximate distance of 25.

For  $PM_{10}$  concentrations, three clusters were identified. The first cluster comprises Zone-1, Zone-6, and Zone-2. This cluster merges with the second cluster, which includes Zone-4, Zone-7, and Zone-3, at a distance of approximately 13. The merged clusters then join with the third cluster, which includes Zone-5, forming a unified cluster at an approximate distance of 25.

**Table 1. Descriptive statistics table for PM<sub>2.5</sub> and PM<sub>10</sub>**

S. N.	Land Use	Number of Land use	PM <sub>2.5</sub>				PM <sub>10</sub>			
			Range (µg/m <sup>3</sup> ) (Min-max)	Mean (µg/m <sup>3</sup> )	Std. Deviation (µg/m <sup>3</sup> )	Coefficient of Variation (%)	Range (µg/m <sup>3</sup> ) (Min-max)	Mean (µg/m <sup>3</sup> )	Std. Deviation (µg/m <sup>3</sup> )	Coefficient of Variation (%)
1.	Zone-1 (Hazaribagh)	4	42	52.25	22.85	43.74	50	114.00	22.20	19.47
2.	Zone-2 (Dhanmondi)	4	31	92.50	14.43	15.60	115	129.50	52.02	40.17
3.	Zone-3 (Ramna, Shahbagh)	7	48	56.86	15.75	27.71	100	97.29	34.59	35.56
4.	Zone-4 (Lalbagh, New Market)	4	41	47.75	22.28	46.65	75	93.50	35.37	37.83
5.	Zone-5 (Chak Bazar, Bangshal, Kotwali)	6	182	116	62.78	54.12	144	154.50	51.68	33.45
6.	Zone-6 (Sutrapur, Gendaria, Jatra Bari)	8	46	69.38	13.62	19.63	78	113.63	29.16	25.66
7.	Zone-7 (Kadamtali)	4	11	47.00	5.35	11.39	72	95.25	32.50	34.12






Fig. 4. Whisker-box plot showing the concentration of PM<sub>2.5</sub> and PM<sub>10</sub> in different zones

Table 2. Significance test

		ANOVA				
		Sum of Squares	df	Mean Square	F	Sig.
PM <sub>2.5</sub>	Between Groups	21281.957	6	3546.993	4.052	0.004
	Within Groups	26259.232	30	875.308		
	Total	47541.189	36			
PM <sub>10</sub>	Between Groups	15825.257	6	2637.543	1.840	0.125
	Within Groups	43001.554	30	1433.385		
	Total	58826.811	36			




Fig. 5. Rescaled distance cluster combine for PM<sub>2.5</sub> and PM<sub>10</sub> in different zones

#### 4. CONCLUSION

The study determined that the average concentrations of  $PM_{2.5}$  and  $PM_{10}$  at 37 educational institutions within the Dhaka South City Corporation were  $68.82 \mu\text{g}/\text{m}^3$  and  $113.95 \mu\text{g}/\text{m}^3$ , respectively. The results show variations in  $PM_{2.5}$  levels across different zones, ranked in descending order as follows: Zone-5 ( $116.00 \mu\text{g}/\text{m}^3$ ), Zone-2 ( $92.50 \mu\text{g}/\text{m}^3$ ), Zone-6 ( $69.38 \mu\text{g}/\text{m}^3$ ), Zone-3 ( $56.86 \mu\text{g}/\text{m}^3$ ), Zone-1 ( $52.25 \mu\text{g}/\text{m}^3$ ), Zone-4 ( $47.75 \mu\text{g}/\text{m}^3$ ), and Zone-7 ( $47.00 \mu\text{g}/\text{m}^3$ ). Additionally,  $PM_{2.5}$  and  $PM_{10}$  concentrations were found to be 1.78 and 1.03 times higher than the NAAQS levels of  $65 \mu\text{g}/\text{m}^3$  and  $150 \mu\text{g}/\text{m}^3$ , respectively.

The study also found that  $PM_{2.5}$  and  $PM_{10}$  dispersion varied by land use, with Zone-5 showing the maximum range, followed by Zone-2, while Zone-7 and Zone-1 had the lowest ranges. The coefficient of variation for  $PM_{2.5}$  and  $PM_{10}$  was highest in Zone-5 and Zone-4, respectively, and lowest in Zone-7 and Zone-1. Whisker-box plots revealed that Zone-4 exhibited the highest dispersion of  $PM_{2.5}$  with a normally skewed distribution, whereas Zone-5 had the highest  $PM_{10}$  values with a positively skewed distribution. Priorities the implementation of consistent air quality monitoring systems at educational institutions, particularly in areas with high levels of  $PM_{2.5}$  and  $PM_{10}$  concentrations, to effectively monitor pollution patterns and trends. Implementing more stringent restrictions and improving pollution control systems can effectively decrease the release of particulate matter, especially in heavily polluted areas like Zone-5 and Zone-2. In addition, the implementation of more green spaces in the vicinity of educational institutions can contribute to the absorption of pollutants and enhance the overall air quality. It is crucial to increase awareness among students, faculty, and the community on the health hazards linked to air pollution. This will promote the adoption of practices that reduce exposure, especially during periods of high pollution. Finally, further investigation is required to determine the origins of particulate matter and comprehend the influence of various land uses on air quality. This will facilitate the creation of focused interventions and policies.

#### 5. LIMITATIONS OF THE STUDY

The research's geographical scope, focusing on South City Corporation educational institutions,

limits its generalizability to other Dhaka or Bangladesh areas due to potential variations in particulate matter concentrations. The study need longtime data to capture the seasonal variations in  $PM_{2.5}$  and  $PM_{10}$  concentrations. The study couldn't focus on all health risk factors related to PM exposure as a result, it couldn't explain the overall public health impacts on the studied populations. The study emphasizes the need for more investigation to completely understand particulate matter pollution's effects on health in cities like Dhaka.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### ACKNOWLEDGEMENT

The authors are thankful to the Center for Atmospheric Pollution Studies (CAPS) for supporting data collection and to the Department of Environmental Science, Stamford University Bangladesh, for their technical support.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. World Health Organization (WHO) Results Report; 2020-2021. Available:<https://www.who.int/.../who-results-report-2020-2021>
2. Manisalidis I, Stavropoulou E, Stavropoulos A, Bezirtzoglou E. Environmental and health impacts of air pollution: A review, *Front. Public Health*, Sec. Environmental health and Exposome. 2020;8.
3. Juginović A, Vuković M, Aranza I, Biloš V. Health impacts of air pollution exposure from 1990 to 2019 in 43 European countries, *Sci Rep*. 2021;11:22516.
4. The World Bank Annual Report ;2018. Available:<https://documents.worldbank.org/en/publication>.
5. World Air Quality Index (AQI) Ranking (IQAir) Report; 2023. Available:<https://www.iqair.com/world-air-quality-ranking>

6. Nahian MA, Ahmad T, Jahan I, Chakraborty N, Nahar Q, Streatfield PK. Poor air quality and pregnancy outcome in Dhaka, Bangladesh, The Journal of Climate Change and Health. 2022; 100187.
7. Majumder AK, Islam M, Rahman M, Patoary MNA, Osman AO. Spatial distribution of atmospheric particulate matter (PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub>) in the Dhaka Metropolitan area, Bangladesh. International Journal of Multidisciplinary Research and Development. 2024;11(8): 69-78
8. Dipta OB, Hassan MM, Hasan M. A study of high-rise building construction practice in Bangladesh, AIP Conference Proceedings. 2023;2713(1).
9. Begum BA, Nasirs M, Randal S, Sivertsen B, Hopke PK. Identification and apportionment of sources from air particulate matter at Urban Environments in Bangladesh British Journal of Applied Science & Technology. 2014;4(27):3930-3955.
10. Fattah MA, Zannat MEU, Morshed SR, Roni AT. Modeling the impact of motorized vehicles' activities on emissions and economic losses in a fast-growing developing city, Dhaka, Bangladesh, Journal of Transport & Health. June 2022;25:101377.
11. Currie J, Zivin JSG, Mullins J, Neidell MJ. What do we know about short and long-term effects of early life exposure to pollution? NBER Work. Pap. 2013;6:217-247.
12. Mizen A, Lyons J, Milojevic A, Doherty R, Wilkinson P, Carruthers D, et al. Impact of air pollution on educational attainment for respiratory health treated students: A cross sectional data linkage study Health & Place. 2020;63(102355).
13. Skoner DP. Allergic rhinitis: Definition, epidemiology, pathophysiology, detection, and diagnosis, J. Allergy Clin. Immunol. 2001;108:2-8.
14. Ahmed S, Islam MS, Rahman MM. Assessment of PM<sub>2.5</sub> concentration near schools in Dhaka City, Bangladesh, Environmental Science and Pollution Research. 2020;27(20):25347-25357.
15. Majumde AK, Rahman M, Patoary MNA. A bibliometrics of air pollution studies in Bangladesh from 1995-2020. World Journal of Advanced Engineering Technology and Sciences. 2023;09(01): 228–239.
16. Woo MK, Young ES, Mostofa MG, Golam M, Afroz S, Hasan MOSI, et al. Lead in air in Bangladesh: Exposure in a Rural community with elevated blood lead concentrations among Young Children. Int. J. Environ. Res. Public Health. 2018;15: 1947.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/122319>