



Influence of Genetic and Non-genetic Factors for Growth Traits of Abergelle and Boer Goat Crossbred F1 Kids in Tigray Region, Ethiopia

Mebrahtom Hagazi ^{a*} and Yosef Tadesse ^b

^a *Abergelle Agricultural Research Center, Ethiopia.*

^b *Haramaya University, Ethiopia.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.globalpresshub.com/review-history/1765>

Original Research Article

Received: 14/09/2024

Accepted: 18/11/2024

Published: 30/11/2024

ABSTRACT

The study was undertaken at Abergelle agricultural research center goat breed evaluation and distribution site with the aim of evaluating the influence of genetic and non genetic factors on growth traits of Abergelle and their Boer goat crossbred F1 kids. A total of 787 records on each growth traits and average daily gains collected for a period of 10 years were used for the analysis by considering birth type, season of birth, sex, parity and year of birth as fixed effects. General Linear Model procedures of SAS and paternal half sib analysis was used to analyze the fixed effects and genetic parameters respectively. The study revealed that genetic groups, birth type, season of birth, sex, parity and year of birth and dam weight at kidding significantly ($p < 0.01$) affect birth weight of kids. The overall mean birth weight was 2.29 ± 0.01 and 2.98 ± 0.03 kg for both genetic groups respectively. Single born kids, kids born in the major wet season and male were heavier than their

*Corresponding author: E-mail: mebrae04@gmail.com;

Cite as: Hagazi, Mebrahtom, and Yosef Tadesse. 2024. "Influence of Genetic and Non-Genetic Factors for Growth Traits of Abergelle and Boer Goat Crossbred F1 Kids in Tigray Region, Ethiopia". *Asian Journal of Research and Review in Agriculture* 6 (1):733-49. <https://jagriculture.com/index.php/AJRRR/article/view/147>.

counterparts. The heritability values for birth weight (0.14 ± 0.05 and 0.20 ± 0.06) for the local and the crossbred were moderate, and show significance difference ($P < 0.05$). The genotypic and phenotypic correlations between the studied growth traits showed positive values for both Abergelle and Boer cross kids. Though the growth performance of the crosses in this study are better than the local, it is still below the expectation. This is because of their intensive management requirement and sub optimal adaptability. Therefore, it is advisable to try to bring genetic improvement through within breed selection among the indigenous goat breeds in terms of reducing importation cost, conservation and adaptation. Considering the non genetic factors in the breeding strategies is also appreciated to improve the production and productivity of local goats.

Keywords: Abergelle goat; correlation; genetic parameter; heritability; non genetic effect; repeatability.

1. INTRODUCTION

“Goat (*Capra hircus*), one of the early domesticated livestock species, provide huge benefits in the livelihood of human population, especially to women, children and aged who are often most vulnerable members of the society in terms of under nutrition and poverty. Small ruminants in general, goats in particular are important sources of income, food (milk and meat), nonfood products (manure and skin), means of risk mitigation during crop failure and social and cultural functions” [1,2].

“The number of goats in Ethiopia is projected to be 50.2 million, with indigenous breeds accounting for 99.97% of the total” [3]. “The gross production value average growth rate from the year 2000 to 2016 was 4.5%” [4]. “Increasing population pressure, land scarcity and diminishing production resources are some of the causes for the higher interest of the community for goat husbandry in comparison to other species in the tropics In addition to that goats are relatively tolerant to drought and widely adapted to different climates, they can survive on woody browse and infrequent watering, less competitive to the cultivated land and their fast reproduction rate enables their owners to recover quickly, following a drought than other livestock species” [5].

“Though, goats play great role in the livelihood of the producers and the economy of the country at large, production and productivity of indigenous goats is very low. This may due to different factors such as poor nutrition, prevalence of diseases, lack of appropriate breed and breeding strategies. Thus, intensification of livestock production system by using improved exotic breeds and crossing them with local breeds is an option and advocated for the past many decades in Ethiopia” [6].

“Combining the adaptability (harsh climates, disease resistance, heat tolerance, ability to utilize poor quality feeds) of native goat with the high milk and meat production potential and fast growth rates of temperate goat through cross breeding led to good adaptability, improve fertility, reduce mortality, efficient growth, feed conversion, milk and meat improvement and this has been practiced as a more sustainable way to improve productivity” [7].

A study of Galal et al. [8] on cross breeding indicated that a substantial increase in milk production has been demonstrated at the 50% level of crossbreeding between Saanen and Ethiopian high land goats. In harsher areas, however, there have been adaptation problems for first crosses Saanen x low land goats indicating that lower level of exotic genotype might be more suitable. Other earlier study [9] evaluated and reported that there were no differences between pure breed Adal and quarter bred Saanen does in any of the reproductive traits recorded. However, cross bred does produced more milk than pure bred does. Breed had no effect on birth weight; however, cross bred progenies were heavier than pure bred Adal goats at 3 months and 6 months.

“There has been introduction of highly productive meat breeds of Boer goat in to Ethiopia to produce cross breed with various local goat breeds in the country. Boer goat is considered to be one of the most desirable goat breeds for meat production” [10]. It has gained worldwide recognition for excellent body conformation, fast growing rate, and good carcass quality. According to Deribe and Taye [11] birth and weaning weight Boer cross with Central high land goat was higher than local breed of Abergelle, Central high land goats, Boran, Somali, and rift valley. When the age advanced to six month and yearling the weight was comparable to central

high land goats and little bit higher than Abergelle and Boran Somali.

Crossing of Boer bucks with Abergelle does, one of the genetic improvement programs had been implemented at Abergelle Agricultural Research Center to evaluate the growth performance and associated factors for the crossbred kids. Though the research had been conducted since 2009, the genetic parameters (heritability, repeatability and genetic and phenotypic correlations) and impact of non-genetic factors on crossbred goats was not yet estimated. With the above background the study was undertaken with the objectives of estimating the genetic and non-genetic factors and evaluating growth performance of Abergelle goats and their crosses with Boer.

2. MATERIALS AND METHODS

2.1 Study Area

The study was conducted at Abergelle Agricultural Research station, Tanqua Abergelle district, Central Zone of Tigray. The area found at 110 km from Mekelle (the capital city of the National Regional State) and 893 km from Addis Ababa with a latitude and longitude of 13° 14' 06" N and 38° 58' 50" E respectively. It is categorized as hot to warm sub-moist lowland (SML-4) sub-agro ecological zone of the region with an altitude of 1300-1800 m above sea level. The main rainy season is June to August with mean annual rainfall ranges from 299 to 650 mm and is characterized by low, erratic and variable rainfall. A little amount of rain is found in the short rain season from March to April. The mean annual temperature ranges from 2/8 to 40°C. There are about 154222 (56222 male and 98108 female) goats in the study area [12].

2.2 Breed Description

As [13,14,15,16,17], the key identifying features of Abergelle breed is that it is found in mixed farming and agro pastoral areas with average flock size of 20±16 and flock structure 84% female and 16% male (14% intact and 2% castrated). Facial profile of the goats is 44% straight and 56% concave and almost all males have horn (89% spiral, 11% straight), in which all directed backwards. Ruffs and wattles are present in both sexes. The common coat colors are plain, patchy, and spotted 56, 33 and 11% respectively. The height at wither (cm), mature body weight (kg), chest girth (cm), ear length

(cm) and horn length (cm) are 71.4±3.5, 33.6±5.9, 79.5±2.9, 13.0±0.8 and 37.07±9.1 for males and 65.0±2.8, 28.4±3.5, 71.2±3.8, 12.7±0.8 and 19.6±5.7 for females respectively. The common birth type is single (98.7%) and rare twinning (1.3%).

The Boer breed is known for its good-quality meat and fast growth rate. Boer goats have earned popularity in many parts of the world. These goats are characterized by their large white bodies and brown heads. The brown color extending or covering the neck and part of the chest regions may be observed in a few. However, some may be completely brown or white in color. Boer goats are known for their rapid growth, high fertility, and adaptability. They are sturdy animals and can be taken care of and managed easily and are also capable of surviving droughts, in the absence of any supplementary feed. The does in this breed have strong maternal skills when compared to other goats. A mature buck has a weight of around 110 to 135 kg, while a mature doe is around 90 to 100 kg. The weights of their young ones at birth are around 3 to 4 kg. Also, the male kids are about 0.5 kg heavier than the female kids [10].

2.3 Animal Management

A semi-intensive management system was practiced at the center with the herd grazing and browsing outdoors during the day (7:30 - 11:30 AM and 2:00 - 5:30 PM) and housed in pens at night. The house was constructed as half concrete building walls with mesh wire and corrugated metal sheet roofs. Hay and water were freely available but supplementary feed (composition of 59% wheat bran, 33% "nug" cake, 4% molasses, 3% lime and 1% salt and nutrient content of CP 20% and ME (Mcal/kg) 2.18) was provided depending on the age and physiological status of the animal.

Does and bucks were herded and housed separately except during the mating periods. During the mating period the bucks kept separately in the day and housed together in the night time. Bucks were selected for general health and absence of observable defects (small testes, hocked joint) before randomly assigned to doe groups. In addition, a few bucks (3-4% of the buck population) were kept in reserve and used to replace if there was sudden death and those with poor libido. The pedigree of each buck was checked for close relationships with the does in their respective groups and when such

relationships exist, does were exchanged between groups taking similarity in age and weight into consideration.

Mating groups were assigned twice a year (June and December). This is because of the feed availability following the rain season and 15 to 20 does were assigned to a single buck. The mating period usually lasts for about 42 days (minimum of two oestrous cycles) and controlled single sire mating was practiced. Beginning from date of mating assignment and following the kidding, all necessary information (sire and dam breed and id, birth weight, date of birth, birth type, sex of kid, parity, year, season of birth and dam weight at kidding, etc.) were recorded in the herd book. Then after, data on weaning weight, six month weight, nine month weight, and yearling weight, and others (death and other health problems) were recorded. Finally, those kids of both sexes with better performance were selected for future parent stock based on own and parent performance.

2.4 Data for Statistical Analysis

The data for this study were collected at Abergelle Agricultural Research Center from 2009 to 2018. A total of 787 records (339 crosses and 448 pure) from 21 sire bucks (12 local and 9 pure Boer blood) and 225 does were collected. Each kid was weighed at birth and then monthly up to yearling using a hanging balance that has 200 g precision. The data were tested for normality, homogeneity, duplications and missing values. Then after checking the following growth traits were evaluated; Birth weight (BW), weaning weight (WW) and six month weight (SMW) as well as average daily gain during different growth stages i.e. from birth to weaning (ADG₁), from weaning to six months (ADG₂) and from birth to six months of age (ADG₃) of all animals at the two seasons (wet and dry) and birth years from 2009 to 2018. Growth traits were not considered after six months. This is because the F1 kids were distributed to the community for on farm assessment and thus data were not enough for analysis. Weaning weight and six month weight was adjusted to 90 and 180 days of age, using a linear regression calculated for that cohort. The pre- and post-weaning daily gains were calculated as total gain divided by the number of days [18].

$$ADG_1 \text{ (Birth to weaning)} = [(WW - BW) / 90] * 1000 \text{ g} \quad (1)$$

$$ADG_2 \text{ (Weaning to Six month)} = [(SMW - WW) / 90] * 1000 \text{ g} \quad (2)$$

$$ADG_3 \text{ (Birth to Six month)} = [SMW - BW] / 180] * 1000 \text{ g} \quad (3)$$

The recorded data was classified with the fixed effects of parity, sex, season, year, type of birth, and dam weight at birth. Data was analyzed using PROC GLM procedure of SAS, 2008 for the least-squares means and standard errors of the fixed factors. Genetic parameters (heritability (h²), Repeatability, and genetic and phenotypic correlations) were computed by the method of paternal half-sib analysis using VARCOMP procedures of SAS 2008.

2.5 Heritability

Estimates of heritability for birth, weaning, six month weight and average daily gains for both breeding groups were determined from sire by using the following formula (Becker, 1993): Heritability from sire component:

$$h^2 s = 4 * \sigma^2 s / \sigma^2 p \quad (4)$$

where $\sigma^2 s$ (sire variance) = (mean square model - mean square error) / k

k=number of progenies produced per sire

2.6 Repeatability

Repeatability estimates were made for traits with repeated records and was calculated by using the formula of Fowler and Cohen, (1986) from variances partitioned using single factor (one way) ANOVA, with both “within and across context consistency” and considering the confounding factors that may be especially problematic when attempting to measure repeatability in goats. Environmental contexts like temperature, feed availability, aging, habituation, physiological states, time, source of animals for study and collection bias and biotic factors like parasitic infestation were potential confounding factors that cause the variations in the repeatability values. Because, repeatability introduces time dimension, it was tried to minimize time dependent changes in sources of measurement error.

Repeatability estimates were made for traits with repeated records and was calculated by using the formula of Fowler and Cohen, (1986).

$$\text{Repeatability (R)} = \frac{B - W}{[B + (N - 1) W]} \quad (5)$$

Where, **B** is variance between individuals and **W** variance within individuals. **N** is simply the size of each sample and for variable sample size it is calculated as **N= [M – (∑ n²/M)/ k-1]**

Where **k**, is individuals measured at least twice and **M**, is the total number of measurements which is number of individuals 'k' plus number of records (**R**), ∑ means 'the sum of and **n** is the number of measurements for an individual.

2.7 Genetic and Phenotypic Correlations

The genetic cause of correlation between characters is largely due to pleiotropy, although linkage of genes on the same chromosome is a transient cause of correlation. Genetic (r_G) and phenotypic (r_P) correlations between traits were estimated from variance and covariance components using the following formulae (Becker, 1984).

$$\text{Genetic Correlation: } r_G = \frac{\sigma^2_{s_{xy}}}{\sqrt{\sigma^2_{s_{xx}} \times \sigma^2_{s_{yy}}}} \quad (6)$$

$$\text{Phenotypic Correlations } r_P = \frac{\sigma^2_{P_{xy}}}{\sqrt{\sigma^2_{P_{xx}} \times \sigma^2_{P_{yy}}}} \quad (7)$$

Where: $\sigma^2_{s_{xy}}$ = sire variance component $\sigma^2_{s_{xy}}$ = sire covariance of traits x and y $\sigma^2_{s_{xx}}$ = sire variance of trait x $\sigma^2_{s_{yy}}$ = sire variance of trait y $\sigma^2_{P_{xy}}$ = phenotypic covariance of traits x and y $\sigma^2_{P_{xx}}$ = phenotypic variance of trait x $\sigma^2_{P_{yy}}$ = phenotypic variance of trait y.

2.8 Statistical Models

Least square means analysis was carried out to examine the influence of fixed effects using the General Linear Model (GLM) procedures of SAS (2008). The following models were used to estimate the growth traits;

2.8.1 Model 1

$$Y_{ijklmno} = \mu + BT_i + P_j + X_k + S_l + Y_m + G_n + Wo + e_{ijklmno} \quad (8)$$

Where;

$Y_{ijklmno}$: The record of BW, 6MW, YW, ADG1, ADG2 or ADG3 measured on nth kid born at mth year of birth, ith birth type, jth parity, kth sex and lth season.

μ : Overall mean

BT_i : effect of ith birth type (i, 1=single, 2= twin);
 P_j : effect of jth parity (J, 1= parity1, 2= parity 2, 3= parity 3 and 4= parity 4);
 X_k : effect of kth sex (k, 1= male and 2= female);
 S_l : effect of lth season (l, 1= wet season and 2= dry season);
 Y_m : effect of mth year (m, 2009- 2018);
 G_n = effect of nth genetic group (Abergelle and Abergelle x Boer)
 Wo = effect of oth weight of dam
 $e_{ijklmno}$: Random error particular to the ijklmnoth observation

2.8.2 Model 2

The (co)variance components and genetic parameters (heritabilities, direct genetic parameter correlations) was estimated by fitting an animal model which fits direct additive effect as random effect and the significant fixed effects. The representation of the animal model was:

$$Y = Xb + Z_1a + e \quad (9)$$

Where Y is the vector of records, b is a vector of an overall mean and fixed effects with incidence matrix X; a, is vector of random additive direct genetic, Z₁ incidence matrix of the random effects and e is a vector of random errors.

3. RESULTS AND DISCUSSION

3.1 Growth and Weight Gain Comparison of the Genotypes

The Least square means of all traits shows high significance difference (p<0.001) between the genetic groups (Table 1).

When compared with the Abergelle (AB) kids, Abergelle x Boer (AB x BO) kids had higher birth weights and live weights in the different growth periods, as well as increased Average Daily Gains (ADG) in the pre-weaning and post-weaning periods (P<0.001). This difference between the two genotypes is due to the effect of heterosis exploited from epistatic and dominant gene effect during cross breeding. Heterosis has a positive effect because in the crossbreds many genes are heterozygous that were homozygous in the parent breeds. Alleles with a negative effect are often recessive. In the cross breeds these negative alleles are ruled out and hybrid vigor could be exploited [19]. The results of the present study on the Boer and Abergelle crosses was in agreement with reports on Boer and

Spanish and Boer and Angora crosses which was reported as they had higher dry matter intake, average daily gain than pure Spanish and Angora goats [20]. The birth and weaning weights of the local breeds in this study were in contrast with the results reported by Khadiga et al. [21] on Sudanese Nubian goats (2.93 ± 0.18), Woldu et al. [22] on Arsi-Bale goats (2.8 kg and 8.39 kg), Shenkute [23] on Keffa goats (2.78 and 9 kg), which were traditionally managed and Zeleke (2017) on Somali goats (3.19 and 11.67 kg) and Teddy kids in Pakistan (8.50±2.18). A study of Galal et al. [8] on cross breeding indicated that a substantial increase in milk production has been demonstrated at the 50% level of crossbreeding between Saanen and Ethiopian high land goats. In harsher areas, however, there have been adaptation problems for first crosses Saanen x low land goats indicating that lower level of exotic genotype might be more suitable. Other earlier study [9] evaluated and reported that there were no differences between pure breed Adal and quarter bred Saanen does in any of the reproductive traits recorded. However, cross bred does produced more milk than pure bred does. Breed had no effect on birth weight; however, cross bred progenies were heavier than pure bred Adal goats at 3 months and 6 months.

According to Deribe and Taye [11] birth and weaning weight of Boer crossed with Central high land goat was higher than local breed of Abergelle, Central high land goats, Boran, Somali, and rift valley. But, when the age advanced to six month and yearling the weight was comparable to central high land goats and a little bit higher than Abergelle and Boran Somali.

However, the BW and WW in this study were in agreement with the study Tegenu [24] on Western lowland goats (2.28 kg) and Getachew et al. [25] on Central High land goats (7.17 kg) respectively. The lower BW and WW in the Abergelle goats may be related with the genetic performance and environment on which Abergelle goats is reared.

Weight gains of local breeds in this study were in agreement with the study Tesema et al. [26] conducted on long eared Somali, Central Highland, and Afar goat breeds supplemented with low and high proportion of concentrate with a weight gain of 37 to 44 g/day. In the contrary the result of average daily weight gain in this study was lower than the study Kebede et al. [27] on Arsi Bale goats with average daily body weight gain reported between 63 and 68 g/day. The reasons for these variations may be due to scarcity of feed in terms of quality and quantity, overall management and husbandry practices in the study area and feed conversion efficiency of the breeds.

3.2 Factors Affect Growth Parameters

3.2.1 Birth type

Majority (N=729) of the kids were born as single. No triple birth was observed in this study. All traits except ADG₃ in twin birth show high significance (p<0.001) difference between the genotypes. But there was no significance difference (p>0.05) on weaning weight, six month weight, ADG₂ and ADG₃ for the pure breeds (Table 2).

Table 1. Mean comparison between the genetic groups

Traits	Genetic group	N	Mean ± SEM	CV%	Minimum	Maximum	P value
BW (kg)	Abergelle	448	2.29 ^b ±0.01	10.2	1.2	3.2	P<0.001
	Abergelle x Boer	339	2.98 ^a ± 0.03	16.1	1.5	4.4	
WW (kg)	Abergelle	379	7.12 ^b ± 0.07	19.2	4.2	12	P<0.001
	Abergelle x Boer	309	8.30 ^a ± 0.11	20.5	4	14	
SMW (kg)	Abergelle	343	9.98 ^b ± 0.11	18.1	4.2	14.8	P<0.001
	Abergelle x Boer	278	10.69 ^a ±0.13	19.4	5.2	20	
ADG1 (g)	Abergelle	379	53.21 ^b ±0.86	31.4	17.8	111	P<0.001
	Abergelle x Boer	309	59.57 ^a ±1.06	27.9	16.7	111.1	
ADG2 (g)	Abergelle	343	44.02 ^b ±1.4	27.5	17.8	128.9	P<0.001
	Abergelle x Boer	278	59.15 ^a ±0.79	22.6	11.11	117.8	
ADG3 (g)	Abergelle	343	32.73 ^b ±1.0	23.7	9	74.4	P<0.001
	Abergelle x Boer	278	42.82 ^a ±0.66	24.2	12.2	88.9	

ADG1 = Average Daily Gain from birth to weaning, ADG2= Average Daily Gain from weaning to six month, ADG3= Average Daily Gain from birth to six month, BW = Birth Weight, CV = Coefficient of Variation, SEM = Standard Error of Mean, SMW = Six Month Weight, WW = Weaning Weight, Means on the same column for a given trait with different superscripts (a,b,) are significantly different (P<0.001)

Table 2. Least square means of birth type for the Abergelle goats and their crosses with Boer

Traits	Genetic group	N	Mean \pm SEM		P-value (within)
			Single	Twin	
BW (kg)	AB	448	2.30 \pm 0.01 (n=418)	2.10 \pm 0.02 (n=30)	***
	ABxBO	339	3.03 \pm 0.30 (n=311)	2.43 \pm 0.06 (n=28)	***
p-value (between)			***	***	
WW (kg)	AB	379	7.08 \pm 0.07 (n=353)	7.59 \pm 0.29 (n=26)	NS
	ABxBO	309	8.5 \pm 0.11 (n=288)	6.30 \pm 0.22 (n=21)	***
p-value (between)			***	***	
SMW (kg)	AB	343	9.95 \pm 0.11 (n=322)	10.53 \pm 0.48 (n=21)	NS
	ABxBO	278	10.90 \pm 0.13 (n=263)	8.14 \pm 0.22 (n=15)	***
p-value (between)			***	***	
ADG ₁ (g)	AB	379	52.64 \pm 0.9 (n=353)	61.07 \pm 3.4 (n=26)	**
	ABxBO	309	60.6 \pm 1.1 (n=288)	42.98 \pm 2.14 (n=21)	***
p-value (between)			*	***	
ADG ₂ (g)	AB	343	44.41 \pm 1.4 (n=322)	38.59 \pm 0.4 (n=21)	NS
	ABxBO	278	60.65 \pm 0.8 (n=263)	47.54 \pm 1.7 (n=15)	***
p-value (between)			***	*	
ADG ₃ (g)	AB	343	32.72 \pm 1.1 (n=322)	32.78 \pm 0.99 (n=21)	NS
	ABxBO	278	43.82 \pm 0.7 (n=263)	31.77 \pm 1.09 (n=15)	***
p-value (between)			***	NS	

AB = Abergelle BO = Boer, ADG₁ = Average Daily Gain from birth to weaning, ADG₂= Average Daily Gain from weaning to six month, ADG₃= Average Daily Gain from birth to six month, BW = Birth Weight, N= Total number of observations, n= number of observations separately SEM = Standard Error of Mean, SMW = Six Month Weight, WW = Weaning Weight, , p values are within breed and between breed comparisons. * Significant at $p < 0.05$ ** significant at $p < 0.01$ ***significant at $p < 0.001$ NS= Non Significant

All body weight traits and pre and post weaning daily weight gains of the crossbred kids had shown high significance ($p < 0.001$) variation between single and twin births. The present result was in consistent with that of Tesema et al. [26] studied on Boer and Central Highland and found significant differences between single and twins of cross breeds in all growth traits. The higher BW of single than twins in this study was in compatible with the findings of Petros et al. [28], Deribe and Taye [29], Mohammed et al.

[30]. This may be the fierce competition of twins on the scarce feed (browse plant or supplements) than single born kid. This difference between the single and twin at birth weight is probably due to the intrauterine environment, where a higher availability of nutrients to the single kid, lack of competition as well as more space may facilitate growth. In this regard, Hagan et al. [31] reported that as the number of foetus increases, the number of caruncles attached to each foetus decreases,

thus reducing the feed supply to the foetus and consequently resulted into lower birth weight. In a comprehensive analysis of factors associated with low kid birth weight, Gardner et al. [32] identified that litter size had the greatest influence on increasing or decreasing birth weight, while year of birth, maternal birth weight, maternal nutrition, and maternal body composition at mating were also important.

3.2.2 Season

Season of birth had significant effect ($p < 0.001$) on all growth traits between the genetic groups. But there is no significance difference for the weaning weight; six month weight and the average daily gains within the genetic groups (Table 3).

Kids born in the wet season were heavier than the kids born in the dry season. Generally, under extensive and semi-intensive system of management where animals spend some time outdoors to graze, season has appreciable influence on a number of economic traits of farm animals. In this study, season of birth had significant effect on birth weight. The birth weight of kids born in the rainy season of both genotypes were significantly ($p < 0.001$) heavier than those of the dry season (Table 3). The availability of green and high-quality forage during the late pregnancy period could have contributed to this observation. This result is in conformity with Kebeda et al. [27], and Baiden [33] who were reported heavier birth weight for kids born in the major wet season. Similar reasons were given by Boujenane and El-Hazzab [34], Ahuya et al. [35] for the similar traits studied. According to these reports, influence of season on growth traits can be explained by variability of feeds in different seasons. Kidding season may well affect kids' growth, because the season could affect the climate thereby feed supply and animal health.

3.2.3 Birth year

Year of birth had significant effect ($p < 0.01$) on all growth parameters and average daily gains at different growth stages between the two genotypes.

Maximum and minimum birth weights were obtained in the years 2012, 2013 and 2018 (3.2 ± 0.1 vs 2.7 ± 0.2) respectively for the cross bred F1 kids. Highest weaning and six month weights were found in the birth years of 2009 while lowest weights were recorded in the year

2016. This difference might be due to changes in management and climatic condition, which might have caused nutritional stress that might have resulted in loss of dams' body weight, induced retardation of fetal growth and reduced birth weight and the sample size of the data used in the analysis.

The results in the present research were in agreement with Bedhane et al. [36] who found the effect of year on growth traits of Arsi Bale goats [37] observed the significant variation in the growth rate of Kutchi kids due to year of birth and Thiruvankadan et al. [38] found significant differences associated with the year of kidding in body weights at birth, weight gain and efficiency in weight gain at different stages of growth. But no significant variation was found in Boer and Turkish hair goats because of year of birth [39]. It is worthwhile to mention, that the level of management is bound to vary according to the ability of the farm manager, the system of crop husbandry, the methods and criteria of culling and his overall supervision at the farm, availability of farm resources and their mobilization in different years.

3.2.4 Sex of kids

Out of the total kids (787) born, about 44% were male 56% were female. Sex of kids had highly significant effect ($p < 0.001$) on birth weight of Abergelle kids (Table 5).

The male kids were heavier than females at birth. Similar results were reported by Eknath [40], Mia et al. [41] on the crosses of Turkish indigenous with that of Boer and Sangamneri kids, but contradicted with that of Rensch [42] in Black Bengal goat observed significant effect of sex on all body weight. The superiority of males to female at BW could be due to hormonal differences that enhance growth. In addition, goat belongs to the most dimorphic mammals and other domestic animals, which exists along the life of animals from fertility until adult age. Yusuff et al. [43] described that in many taxa, sexual size dimorphism (SSD) varies with body size and larger species generally exhibits higher male to female body size ratio, which is known as "Ranch's rule". The cause of sexual dimorphism is described by the study of Dadi et al. [44] in that, resource availability may influence the body size of one sex to a greater degree than the other when relative energy expenditure on mating and reproduction is greater for that sex. In addition, ecological factors may select for small body size in both sexes but female body size may be constrained by fecundity selection.

Table 3. Effect of season on growth traits of Abergelle goats and their crosses with Boer

Traits	Genetic group	Season (Mean ±SEM)		P value (within)
		Wet	Dry	
BW	AB	2.31 ± 0.02 (n=332)	2.21±0.03 (n=116)	***
	AB x BO	3.01 ± 0.03 (n=296)	2.78±0.07 (n=43)	**
<i>P</i> value (between)		***	***	
WW	AB	7.10 ± 0.08 (n=282)	7.14±0.14 (n=97)	NS
	AB x BO	8.37 ± 0.12 (n=273)	7.87 ± 0.29 (n=36)	NS
<i>P</i> value (between)		***	***	
SMW	AB	9.90 ± 0.13 (n=257)	10.23 ± 0.23 (n=86)	NS
	AB x BO	10.74 ± 0.14 (n=249)	10.32 ± 0.37 (n=29)	NS
<i>P</i> value (between)		**	NS	
ADG ₁	AB	53.05 ± 0.95 (n=282)	53.69 ± 1.89 (n=97)	**
	AB x BO	59.52 ± 1.15 (n=273)	56.62 ± 2.79 (n=36)	NS
<i>P</i> value (between)		**	*	
ADG ₂	AB	44.20 ± 1.58 (n=257)	43.49 ± 2.77 (n=86)	NS
	AB x BO	59.79 ± 0.84 (n=249)	58.04 ± 2.29 (n=29)	NS
<i>P</i> value (between)		***	***	
ADG ₃	AB	32.66 ± 1.11 (n=257)	32.94 ± 2.10 (n=86)	NS
	AB x BO	42.96 ± 0.71 (n=249)	41.88 ± 1.84 (n=29)	NS
<i>P</i> value (between)		**	**	

AB = Abergelle BO = Boer, ADG₁ = Average Daily Gain from birth to weaning, ADG₂= Average Daily Gain from weaning to six month, ADG₃= Average Daily Gain from birth to six month, BW = Birth Weight, n= number of observations, SEM = Standard Error of Mean, SMW = Six Month Weight, WW = Weaning Weight, p values are within breed and between breed comparisons * Significant at p<0.05 ** significant at p<0.01 ***significant at p<0.001 NS= Non Significant

Table 4. The effect of Birth year on the growth parameters of Abergelle goats and their crosses

Year	Genetic group	N	Traits					
			BW	WW	SMW	ADG ₁	ADG ₂	ADG ₃
2009	AB	14	2.3±0.2 ^b	7.1±0.1 ^b	9.7±0.2 ^b	53.7±2.1 ^b	54.2±2.3 ^b	41.1±1.6 ^b
	ABxBO	71	2.9±0.1 ^a	9.6±0.2 ^a	12.4±0.3 ^a	74.8±2.2 ^a	62.5±1.8 ^a	52.7±1.5 ^a
2010	AB	18	2.3±0.0 ^b	6.8±0.2 ^b	10.1±0.4 ^a	51.0±1.8 ^b	61.6±2.1 ^a	43.7±1.1 ^a
	ABxBO	65	2.9±0.1 ^a	7.6±0.2 ^a	9.9±0.3 ^b	52.3±2.3 ^a	57.9±1.6 ^b	39.2±1.3 ^b
2011	AB	17	2.4±0.0 ^b	7.0±0.0 ^b	9.7±0.3 ^b	51.6±2.3 ^b	55.9±2.2 ^b	40.6±1.7 ^b
	ABxBO	42	3.0±0.1 ^a	8.0±0.3 ^a	10.6±0.4 ^a	55.7±2.9 ^a	62.4±2.2 ^a	42.2±1.7 ^a
2012	AB	24	2.5±0.0 ^b	7.2±0.2 ^b	9.9±0.3 ^b	52.2±2.5 ^b	57.4±2.6 ^b	40.8±1.5 ^b
	ABxBO	48	3.2±0.1 ^a	8.7±0.3 ^a	11.4±0.3 ^a	61.6±2.4 ^a	65.5±2.0 ^b	46.1±1.5 ^a
2013	AB	67	2.3±0.0 ^b	7.2±0.2 ^b	11.3±0.3 ^a	54.4±2.3 ^b	72.8±3.1 ^a	50.2±1.5 ^a
	ABxBO	37	3.2±0.1 ^a	9.1±0.3 ^a	10.5±0.4 ^b	65.4±3.0 ^a	51.4±2.5 ^b	40.5±1.8 ^b
2014	AB	42	2.3±0.1 ^b	6.9±0.2 ^b	9.3±0.4 ^{ns}	51.8±2.3 ^{ns}	52.9±2.8 ^b	39.5±2.1 ^a
	ABxBO	20	3.0±0.1 ^a	7.7±0.3 ^a	9.6±0.4 ^{ns}	51.8±2.7 ^{ns}	54.5±2.8 ^a	36.4±1.7 ^b

Year	Genetic group	N	Traits					
			BW	WW	SMW	ADG1	ADG2	ADG3
2015	AB	61	2.2±0.0 ^b	6.7±0.2 ^b	8.7±0.2 ^b	49.7±1.6 ^a	47.9±1.6 ^b	35.9±1.2 ^{ns}
	ABxBO	14	3.0±0.0 ^a	7.2±0.4 ^a	9.1±0.4 ^a	46.6±3.1 ^b	54.8±3.1 ^a	34.2±1.5 ^{ns}
2016	AB	70	2.3±0.0 ^b	7.1±0.1 ^a	10.9±0.2 ^a	53.8±1.5 ^a	67.3±2.2 ^a	48.1±1.4 ^a
	ABxBO	13	2.8±0.1 ^a	6.5±0.2 ^b	8.8±0.2 ^b	40.8±2.1 ^b	56.8±2.3 ^b	33.0±1.4 ^b
2017	AB	82	2.2±0.0 ^b	7.5±0.2 ^a	9.7±0.3 ^a	58.3±2.2 ^a	48.7±1.6 ^b	41.4±1.4 ^a
	ABxBO	18	3.0±0.1 ^a	7.0±0.4 ^b	9.3±0.5 ^b	44.2±3.5 ^b	58.6±3.1 ^a	34.9±2.4 ^b
2018	AB	53	2.3±0.0 ^b	7.2±0.2	9.2±0.2 ^b	53.9±2.2 ^b	47.8±1.2 ^b	38.3±1.1 ^b
	ABxBO	11	2.7±0.2 ^a	7.7±0.4 ^a	10.5±0.8 ^a	55.4±4.1 ^a	61.5±6.8 ^a	43.6±4.1 ^a

AB= Abergelle, ADG1 = Average Daily Gain birth to weaning, ADG2 = Average Daily Gain from weaning to Six month, ADG3 = Average Daily Gain from birth to six month, BO= Boer, BW = Birth Weight, N = Number of observations, SEM = Standard Error of Mean, SMW = Six Month Weight, Super scripts down column variation between genetic group, WW = Weaning Weight

Table 5. The effect of sex on the growth performance of local and cross breed goats

Traits	Genetic group	N	Sex (Mean ±SEM)		P value (within)
			Male	Female	
BW	AB	219	2.42±0.02	2.16±0.02	***
	AB x BO	150	3.06±0.04	2.91±0.04	**
<i>P value (between) *** **</i>					
WW	AB	181	7.19±0.09	7.06±0.10	NS
	AB x BO	131	8.29±0.15	8.31±0.15	NS
<i>P value (between) *** **</i>					
SMW	AB	159	10.09±0.17	9.89±0.15	NS
	AB x BO	122	10.75±0.19	10.64±0.17	NS
<i>P value (between) ** **</i>					
ADG1	AB	181	52.57±1.16	53.79±1.26	NS
	AB x BO	131	58.21±1.56	59.91±1.46	NS
<i>P value (between) * *</i>					
ADG2	AB	159	42.94±2.07	45.05±1.82	NS
	AB x BO	122	61.20±1.21	58.28±1.03	NS
<i>P value (between) *** **</i>					
ADG3	AB	159	30.92±1.47	34.46±1.32	NS
	AB x BO	122	42.72±1.01	42.90±0.88	NS
<i>P value (between) ** **</i>					

ADG1 = Average Daily Gain birth to weaning, ADG2 = Average Daily Gain from weaning to Six month, ADG3 = Average Daily Gain from birth to six month, BW = Birth Weight, SEM = Standard Error of Mean, SMW = Six Month Weight, WW = Weaning Weight, p values are within breed and between breed comparisons. * Significant at p<0.05 ** significant at p<0.01 ***significant at p<0.001 NS= Non Significant

3.2.5 Parity

Parity of dam in this study had significant effect (p<0.01) on least square mean of the growth parameters of kids between the genotypes. But parity had significant effect only on birth weight of local Abergelle goats (Table 6).

Average BW of Kids of the crosses from fourth parity dams exceed by about 27% (3.14±0.09) than the kids from first parity (2.87±0.05). The

increment in birth weight as parity increase is related with utilization of more feed for production and reproduction than for growth in older age doe and provide more feed for the fetus than young age does [31,37]. Hence, as parity increases mothering ability and milk production increases and this is observed in kid growth. Effect of parity on weight is reported in different studies [45,46]. The result of this study was in line with the results reported by Sharma et al. [47] on West African Dwarf and Red Sokoto goats.

Table 6. Least square means of parity in Abergelle local and crossbred goats

Traits	Genetic group	Parity (Mean \pm SEM)							
		N	1	N	2	N	3	N	4
BW (kg)	AB	105	2.2 \pm 0.0	131	2.3 \pm 0.0	123	2.3 \pm 0.0	89	2.5 \pm 0.0
	AB x BO	78	2.9 \pm 0.1	90	2.9 \pm 0.5	103	3.1 \pm 0.1	68	3.1 \pm 0.1
			***		***		***		***
WW (kg)	AB	90	7.0 \pm 0.1	114	7.1 \pm 0.1	98	7.1 \pm 0.1	77	7.3 \pm 0.2
	AB x BO	71	8.8 \pm 0.2	85	8.1 \pm 0.2	94	8.1 \pm 0.2	59	8.1 \pm 0.3
			***		**		**		**
SMW (kg)	AB	77	10.4 \pm 0.2	101	9.7 \pm 0.2	91	10.0 \pm 0.2	74	10.0 \pm 0.2
	AB x BO	66	11.5 \pm 0.2	74	10.6 \pm 0.2	86	10.3 \pm 0.2	52	10.2 \pm 0.32
			**		**		*		NS
ADG1 (g)	AB	90	54.6 \pm 1.6	114	53.9 \pm 1.5	98	50.7 \pm 2.0	77	53.9 \pm 1.9
	AB x BO	71	65.6 \pm 2.0	85	58.0 \pm 2.0	94	56.0 \pm 1.9	59	54.5 \pm 2.3
			**		*		*		NS
ADG2 (g)	AB	77	44.29 \pm 3.1	101	42.4 \pm 3.1	91	53.93 \pm 1.9	74	47.3 \pm 2.7
	AB x BO	66	61.6 \pm 1.8 ^a	74	59.4 \pm 1.4	86	58.2 \pm 1.5	52	58.3 \pm 2.2 ^a
			**		**		NS		*
ADG3 (g)	AB	77	33.8 \pm 2.2	101	31.9 \pm 1.8	91	53.2 \pm 0.9	74	34.9 \pm 1.9
	AB x BO	66	47.7 \pm 1.3	74	42.5 \pm 1.2	86	39.97 \pm 1.2	52	39.0 \pm 1.5
			*		*		*		NS

AB = Abergelle, BO = Boer, ADG1 = Average Daily Gain birth to weaning, ADG2 = Average Daily Gain from weaning to Six month, ADG3 = Average Daily Gain from birth to six-month, BW = Birth Weight, SEM = Standard Error of Mean, SMW = Six Month Weight, T = Total, WW = Weaning Weight. Superscripts (a,b) down column represent differences between the genetic groups. P values show differences within genotypes

Table 7. The effect of dam weight at kidding on growth traits of local and cross breed goats in the study area

Factor	Abergelle (Mean \pm SEM)						
	Dam weight	BW(kg)	WW(kg)	SMW(kg)	ADG1(g)	ADG2(g)	ADG3(g)
	15-25	2.27 \pm 0.02	7.04 \pm 0.11	10.11 \pm 0.17	52.72 \pm 1.21	60.43 \pm 1.69	43.36 \pm 0.96
	26-35	2.31 \pm 0.02	7.22 \pm 0.11	9.84 \pm 0.15	53.80 \pm 1.41	52.98 \pm 1.23	42.11 \pm 0.87
	Total	2.29 \pm 0.01	7.12 \pm 0.07	9.98 \pm 0.11	53.21 \pm 0.86	44.02 \pm 1.37	32.73 \pm 0.99
	P-value	0.043	0.969	0.864	0.965	0.354	0.602
Abergelle X Boer (Mean \pm SEM)							
	15-25	2.88 \pm 0.04	8.84 \pm 0.18	11.43 \pm 0.22	66.26 \pm 1.84	60.65 \pm 1.29	42.82 \pm 0.66
	26-35	3.04 \pm 0.04	7.98 \pm 0.13	10.25 \pm 0.16	54.95 \pm 1.23	58.93 \pm 0.99	40.04 \pm 0.75
	Total	2.98 \pm 0.03	8.30 \pm 0.11	10.69 \pm 0.13	59.15 \pm 1.06	59.57 \pm 0.79	42.82 \pm 0.66
	P-value	0.049	0.437	0.357	0.442	0.381	0.356

ADG1 = Average Daily Gain from birth to weaning ADG2= Average Daily Gain from weaning to six month, ADG3= Average Daily Gain from birth to six- month weight, BW = Birth Weight, SEM = Standard Error of Mean, SMW = Six Month Weight, WW = Weaning Weight. NS = Non significant * = significant at 0.05

3.2.6 Dams weight at kidding

Differences due to dams' weight at kidding were found to be significant ($p < 0.05$) to kid's body weight at birth (Table 7).

The minimum and maximum body weights of does at kidding were 18 and 34 with an average of 25.78 kg. Similar values were reported by different scholars for Abergelle breeds [11,16,17] Only birth weight of kids was affected by the dam weight. The most important influences on birth

weight in small ruminants are the number of newborns, followed by number of parturitions, body weight and condition of female around the time of parturition, nutrition and gender of newborns [33]. This finding was related with the findings of Eknath [40] on weight of Boer and Turkish hair goat. However, Kharkar et al. [48], Bosso et al. [49] reported non-significant effect of dam's weight at kidding on all growth traits at birth and 3 months of age in India goats and at 6, 9, and 12 months of ages in Jamunapari goats respectively.

3.3 Estimation of Genetic and Phenotypic Parameters

3.3.1 Heritability

Results given in Table 8 showed a range of the heritability and repeatability estimates for body weight from birth to six months of age between the genetic groups. Heritability of the growth traits and pre and post weaning daily gains revealed significance variation ($p < 0.05$) between Abergelle goats and their crosses with Boer (Table 8).

The heritability values for birth weight and ADG2 of the pure breed were moderate, while the weaning weight, six-month weight, ADG1 and ADG3 were low. The heritability value of local breed was in a decreasing trend as it goes from BW up to six-month weight. This may indicate the higher impact of environment and management at later age in comparison to earlier age and it may be due to absence of appropriate husbandry practices in the study area. This study result was different from Bolacali et al. [39] who studied the heritability of body weight tended to increase with increasing age from birth to six months.

The heritability value of birth weight in the present study of both genotypes were lower than the studies of Al-Shorepy et al. [50], Zhang et al. [51] on West African Dwarf goats and Emirati goats reported the heritability estimates of 0.50 and 0.39, respectively. Zhang et al. [51] and Al-Saef [52] also reported the heritability estimates of birth weight in Syrian Damascus goat and Boer goat as 0.41 and 0.30, which are higher than the present study. Lower heritability values (0.09 ± 0.08 , 0.03 ± 0.08 , 0.04 ± 0.08 , 0.02 ± 0.10) were reported on birth, weaning, six month and yearling weights of Arsi Bale goats of Ethiopia respectively [37] which are similar to the present study. The heritability values in this study were lower than Draa, West Africa Dwarf, Ardi, Angora, Black Bengal, Boer, Damascus and Jamnapari goats [19,51,53,54].

This difference in the heritability of traits could be due to genetic and non genetic variations of the breeds and production environment. The difference in estimates may also arise from sample size, method of data analysis and the

model used by each investigator. Heritability of most traits in the current study was low to moderate for both local and crossbred, which implies most variations are due to environmental and management variations.

3.3.2 Repeatability

The repeatability values of birth weight, six-month weight, pre weaning daily gain and daily gain from birth to six month showed significance difference ($p < 0.05$) between the genotypes while, there was non-significant variation ($p > 0.05$) in the post weaning daily gain (ADG2) (Table 8). The maximum (0.26 ± 0.12) and minimum (0.06 ± 0.04) repeatability values were observed on birth weight and six-month weight respectively. The repeatability values of the crosses were agreed with report of Alade et al. [55] of Tanzanian blended goats but they were by far lower than the results of Bedhane et al. [56] repeatability on birth weight, pre weaning weight, weaning weight, post weaning gain and 9 month body weight were 0.61 ± 0.15 , 0.37 ± 0.11 , 0.52 ± 0.12 , 0.24 ± 0.08 and 0.40 ± 0.04 respectively.

The variations in the repeatability values between the studied breeds and among other breeds were due to the continuous physiological changes of the dams and variations on the individual progeny performance. Repeatability value is greater than heritability value since repeatability estimates include the permanent environmental variance in addition to the additive genetic variance component. When repeatability is high, it indicates that a single record of performance on an animal is, on average, a good indicator of that animal's producing ability. When repeatability is low, a single phenotypic value tells very little about producing ability. Lower repeatability estimate for traits could be also due to higher influence of specific environmental effects on a given record that may inflate within animal records variability [57].

3.4 Genetic and Phenotypic Correlation

Genetic correlations of all traits of the cross breeds had high significant difference ($p < 0.01$). High genetic correlations ($p < 0.001$) were observed in the correlation between six month weight and ADG3 in both genetic groups (Table 9).

Table 8. Estimates of heritability for the growth parameters of local and cross breed goat populations

Traits	Genetic group	Heritability (h ² s)	Repeatability
BW	AB	0.14 ^b ±0.01	0.24 ^a ±0.05
	AB x BO	0.20 ^a ±0.06	0.20 ^b ±0.08
WW	AB	0.08 ^b ±0.01	0.19 ^{ns} ±0.06
	AB x BO	0.14 ^a ±0.05	0.17 ^{ns} ±0.04
SMW	AB	0.04 ^b ±0.02	0.14 ^b ±0.03
	AB x BO	0.10 ^a ±0.04	0.26 ^a ±0.12
ADG1	AB	0.11 ^b ±0.03	0.08 ^b ±0.02
	AB x BO	0.16 ^a ±0.07	0.18 ^a ±0.04
ADG2	AB	0.09 ^b ±0.02	0.10 ^{ns} ±0.07
	AB x BO	0.13 ^a ±0.06	0.09 ^{ns} ±0.03
ADG3	AB	0.07 ^{ns} ±0.02	0.24 ^a ±0.05
	AB x BO	0.09 ^{ns} ±0.01	0.20 ^b ±0.08

ADG1 = Average Daily Gain from birth to weaning, ADG2 = Average Daily Gain from weaning to Six month, ADG3= Average Daily Gain from birth to six- month BW = Birth Weight, h²s = Heritability of sire, SMW = Six Month Weight, WW = Weaning Weight p<0.05

Table 9. Genetic (below diagonal) and phenotypic (above diagonal) diagonal (number of observations) correlations of body weights and average daily gains

Genetic group		BW	WW	SMW	ADG1	ADG2	ADG3
Abergelle	BW	448	0.08 ^a	0.01 ^a	0.1 ^a	0.12 ^b	0.12 ^a
	WW	0.12 ^a	379	0.59 ^a	0.98 ^{ns}	0.07 ^a	0.58 ^a
	SMW	0.25 ^{ns}	0.51 ^a	343	0.59 ^a	0.75 ^a	0.99 ^{ns}
	ADG1	0.31 ^{ns}	0.49 ^a	0.41 ^a	379	0.01 ^a	0.60 ^a
	ADG2	0.15 ^a	0.07 ^a	0.75 ^a	0.01 ^a	343	0.73 ^a
	ADG3	0.17 ^a	0.58 ^{ns}	0.29 ^a	0.06 ^a	0.29 ^a	343
Abergelle x Boer	BW	339	0.55 ^b	0.50 ^b	0.32 ^b	0.50 ^b	0.31 ^b
	WW	0.35 ^b	309	0.88 ^b	0.97 ^{ns}	0.33 ^b	0.84 ^b
	SMW	0.20 ^{ns}	0.18 ^b	278	0.85 ^b	0.70 ^b	0.98 ^{ns}
	ADG1	0.32 ^{ns}	0.29 ^b	0.55 ^b	309	0.22 ^b	0.86 ^b
	ADG2	0.50 ^b	0.23 ^b	0.40 ^b	0.20 ^b	278	0.65 ^b
	ADG3	0.31 ^b	0.54 ^{ns}	0.68 ^b	0.56 ^b	0.45 ^b	278

ADG1 = Average Daily Gain from birth to weaning, ADG2 = Average Daily Gain from weaning to Six month, ADG3= Average Daily Gain from birth to six-month BW = Birth Weight, SMW = Six Month Weight, WW = Weaning Weight p<0.05. ^{a,b} Significantly difference between the genetic groups ns = Non significant

Almost all genetic correlations in the cross breeds were ranged from moderate to high and positive while in the pure breeds all (low, medium and high) correlations were observed. The genetic and phenotypic correlations between birth weight and weaning weight, birth weight and six month weight and weaning weight and six month of the pure Abergelle were almost similar to that of Arsi- Bale goats which was numerically reported as 0.70±0.55 and 0.17, 0.64±0.47 and 0.19 and 0.57±0.43 and 0.65 respectively [55]. The results also agreed to Boujenane and El-Hazzab [19], Zhang et al. [51] studied genetic and phenotypic correlation of Draa WAD goats

respectively and reported BW vs. WW, 0.58 and 0.27, BW vs SMW, 0.28 and 0.15, WW vs. SMW, 0.43 and 0.51 and BW vs. WW, 0.74 ± 0.08 and 0.30 for genetic and phenotypic correlations respectively.

The phenotypic correlations between all traits found same trend as the genetic correlations in accordance with previous studies in different goat breeds. Phenotypic correlation between birth weight, one month weight and three months weight of West African dwarf kids were ranging from 0.45 to 0.99 which was within the range of the current study findings [52]. The high genetic

correlation in the cross breeds between the two traits in present study indicated that both traits are under the influence of similar genes and selection for higher birth weight will result in higher weaning weight as a correlated response. Similarly, the phenotypic correlation between the two traits indicated a positive relationship between the traits.

The genotypic and phenotypic correlations between the studied growth traits at different stages showed positive values for both Abergelle and crossbred goats. Similarly, Xiong et al. [58] found positive genotypic and phenotypic correlations between body weights at different ages.

4. CONCLUSION AND RECOMMENDATION

In conclusion, it was determined that the genotype, parity of dam, live weight of dam at kidding, sex, birth type, season and year of birth have a significant influence on the birth weight, live weight and ADG of both genotype kids. Heritability (h^2) were estimated for traits viz. live weight at (birth, 3 month (weaning), 6 month) and average daily gains at different growth stages (ADG1, ADG2 and ADG3) and values were ranged from low to moderate for both genetic groups and this show most variations are not due to genetic rather they are due to environmental and managemental variations.

The genotypic and phenotypic correlations between the studied growth traits at different stages showed positive values for both Abergelle and Boer goats. Almost all genetic correlations in the cross breeds were ranged from moderate to high and positive while in the pure breeds low, medium and high correlations were observed. Generally, an effective breeding program depends on the accuracy of genetic and nongenetic parameter estimation, which includes heritability, repeatability and genetic and phenotypic correlations between traits. So, knowing and incorporating the production and reproduction factors in a breeding program may lead to a great.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENT

Authors would like to express their deepest and sincere gratitude to Abergelle Agricultural Research Center livestock researchers and farm workers for their support and Tigray Agricultural Research Institute for financing this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aklilu H, Assen E. Sheep and goat production and utilization in different agro-ecological zones in Tigray, Ethiopia. *Livest Res Rural Dev*; 2012.
2. Neme Y. Study of productive and reproductive performances and farmers' traits preferences for breeding of small ruminants in Ada Barga and Ejere districts of West Shoa Zone, Oromia, Ethiopia; 2016.
3. Central Statistical Agency (CSA). Report on livestock and livestock characteristics agricultural sample survey. 2020;2.
4. Food and Agriculture Organization (FAO). The state of food and agriculture: Innovation in family farming. Rome: FAO; 2019.
5. Jembere T. Genetic parameters, productivity indices and breeding plans for designing community-based goat breeding programs in Ethiopia [doctoral dissertation]. Haramaya University, Haramaya, Ethiopia; 2016.
6. Hagos A, Gizaw S, Urge M. Begait goat production systems and breeding practices in Western Tigray, North Ethiopia; 2017.
7. Effa K, Tadelle ZW, Haile A. Genetic and environmental trends in term dairy cattle genetic improvement programmes in the central tropical highlands of Ethiopia. *J Cell Anim Biol*. 2011;5.
8. Galal ESE, Afifi EA, EL-Kimaray S, Ahmed IA, Shawar AF. Lamb survival as affected by inbreeding and crossbreeding. *J Agric Sci*. 1981;96(1):1–5.
9. Awgichew K, Gipson TA. Overview of genotype program activities. *Proc ESGPIP Mid-term Conf*, Hawassa, Ethiopia; 2009.
10. Lu CD. Milk feeding and weaning of goat kids. *Small Rumin Res*; 2001.
11. Deribe B, Taye M. Evaluation of growth performance of Abergelle goats under

- traditional management systems in Sekota district, Ethiopia. *Pak J Biol Sci.* 2013; 16(14).
12. Bureau of Agriculture and Rural Development (BOARD). Livestock population in Tanqua Abergelle woreda; 2020.
 13. Food and Agricultural Research Management for Africa (FARM Africa). Goat types of Ethiopia and Eritrea: Physical description and management systems. London: FARM Africa; 1996.
 14. Abegaz S, Mwai AK, Gebreyesus G, Haile A, Rischkowsky B, Gizaw S, et al. Review of goat research and development projects in Ethiopia. ILRI Project Report. Nairobi, Kenya: International Livestock Research Institute; 2014.
 15. Hassen H, Michael BM, Rischkowsky B, Tibbo M. Phenotypic characterization of Ethiopian indigenous goat populations; 2012.
 16. Alemu A. Evaluation of Abergelle and Central Highland goat breeds as an input for designing community-based breeding programs [MSc thesis]. Haramaya University, Dire Dawa, Ethiopia; 2015.
 17. Brihanae M. Morphological characterization of goat population and performance evaluation of Abergelle goat under community-based breeding program in Tigray and Amhara regional states, Ethiopia [MSc thesis]. Bahir Dar University; 2017.
 18. Mandal A, Pant KP, Nandy PK, Rout PK, Roy R. Genetic analysis of growth traits in Muzaffarnagari sheep. *Trop Anim Health Prod.* 2003;35.
 19. Boujenane I, El-Hazzab A. Genetic parameters for direct and maternal effects on body weights of Draa goats. *Small Rumin Res.* 2012;80(1–3).
 20. Cameron MR, Luo J, Sahlu T, Hart SP, Coleman SW, Goetsch AL. Growth and slaughter traits of Boer x Spanish, Boer x Angora, and Spanish goats consuming a concentrate-based diet. *J Anim Sci.* 2001;79.
 21. Khadiga ME, Khair ABM, Lutfi AMA. Estimates of phenotypic and genetic parameters of growth traits in the Sudanese Nubian goat. *Res J Anim Vet Sci;* 2008.
 22. Woldu T, Dadi H, Guru M, Gelashe D. Productivity of Arsi Bale goat types under farmers' management condition: A case of Arsi Negelle. In: Degefa T, Feyissa F, editors. Proceedings of the 13th Annual Conference of the Ethiopian Society of Animal Production (ESAP); 2004.
 23. Shenkute B. Production and marketing systems of small ruminants in Gomma district of Jimma Zone, western Ethiopia [MSc thesis]. Hawassa University, Hawassa, Ethiopia; 2009.
 24. Tegenu T. Characterization of goat production systems and on-farm evaluation of the growth performance of grazing goats supplemented with different protein sources in Metema woreda, Amhara region, Ethiopia [MSc thesis]. Haramaya University, Haramaya, Ethiopia; 2009.
 25. Getachew T, Lemma S, Gizaw S. Growth and reproductive performance of Central Highland goats in North Shoa and South Wollo. Proceedings of the First Annual Conference on Completed Livestock Research Activity; 2006.
 26. Tesema Z, Tilahun M, Deribe B, Lakew M, Belayneh N, Zegeye A, et al. Effect of non-genetic factors on pre-weaning growth, survivability and prolificacy of Central Highland x Boer crossbred goats in North Eastern Ethiopia. *Livest Res Rural Dev.* 2017;29.
 27. Kebede T, Haile A, Dadi H, Alemu T. Genetic and phenotypic parameter estimates for reproduction traits in indigenous Arsi-Bale goats. *Trop Anim Health Prod;* 2012.
 28. Petros A, Aragaw K, Shilima B. Pre-weaning kid mortality in Adami Tulu Jedokombolcha district, Mid Rift Valley, Ethiopia. *J Vet Med Anim Health.* 2013;6.
 29. Deribe B, Taye M. Evaluation of growth performance of Abergelle goats under traditional management systems in Sekota district, Ethiopia. *Pak J Biol Sci.* 2013;16.
 30. Mohammed KM, Kamal EL-Den MA, Dahmouh AY. Heritability and variance components estimates for growth traits in Saudi Ardi goat and Damascus goat and their crosses; 2018.
 31. Hagan JK, Apori SO, Bosompem M, Ankobea G. Morphological characteristics of indigenous goats in the coastal savannah and forest eco-zones of Ghana. *J Anim Feed Res.* 2014;4.
 32. Gardner DS, Buttery PJ, Daniel Z, Symonds ME. Factors affecting birth

- weight in sheep: Maternal environment; 2007.
33. Baiden RY. Birth weight, birth type and pre-weaning survivability of West African Dwarf goats raised in the Dangme West district of the Greater Accra region of Ghana. *Trop Anim Health Prod.* 2007;39:141–7.
 34. Boujenane I, El-Hazzab A. Genetic parameters for direct and maternal effects on body weights of Draa goats. *Small Rumin Res;* 2008.
 35. Ahuya CO, Ojango JMK, Mosi RO, Peacock CP, Okeyo AM. Performance of Toggenburg dairy goats in smallholder production systems of the eastern highlands of Kenya. *Small Rumin Res.* 2009;83.
 36. Bedhane M, Haile A, Dadi H, Alemu T. Estimates of genetic and phenotypic parameters for growth traits in Arsi-Bale goat in Ethiopia. *J Anim Sci.* 2013;3.
 37. Yadav JS, Yadav MC, Khan BU. Non-genetic factors affecting growth of Kutchi goats. *Indian J Anim Sci.* 2008;78.
 38. Thiruvankadan AK, Panneerselvam S, Kandasamy N. Reproductive performance of Kanni Adu goats under field conditions. *Indian J Anim Sci.* 2000;70.
 39. Bolacali M, Öztürk Y, Yilmaz O, Küçük M, Karsli MA. Effect of genotype and non-genetic factors on growth traits and survival rates in Turkish indigenous hair goats and their first crosses with Boer bucks. *Indian J Anim Res.* 2017;51(6).
 40. Eknath TA. Effect of various genetic and non-genetic factors affecting growth performance of Sangamneri kids [MSc thesis]. Maharashtra Animal and Fishery Sciences University, Nagpur, India; 2015.
 41. Mia MM, Khandoker MA, Husain SS, Faruque MO, Notter DR, Haque MN. Genetic evaluation of growth traits of Black Bengal goat. *Iran J Appl Anim Sci.* 2013;3(4).
 42. Rensch B. Evolution above the species level. New York: Columbia University Press; 1960.
 43. Yusuff AT, Sola-Ojo FE, Ojo V, Alli OI, Fayeye TR. Heritability estimates for body weight in pure and crossbred Nigerian goats. *J Anim Prod Res;* 2015.
 44. Dadi H, Gameda D, Shelima B, Fayera T, Tadesse T, Woldu T, et al. Non-genetic factors influencing post-weaning growth and reproductive performances of Arsi-Bale goats. *Livest Res Rural Dev.* 2008;20(7).
 45. Jimenez-Badillo MR, Rodrigues S, Sanudo C, Teixeira A. Non-genetic factors affecting live weight and daily gain weight in Serrana Transmontano kids. *Small Rumin Res.* 2009;84.
 46. Bemji MN, Osinowo OA, Ozoje MO, Adebambo OA, Aina ABJ. Live weight changes during lactation and its relationship with milk off-take and yield in West African Dwarf and Red Sokoto goats intensively managed within the humid zone of Nigeria. *Niger J Anim Prod.* 2006;33(1).
 47. Sharma MC, Pathodiya OP, Tailor SP, Mishra S. Pre-weaning growth traits of Sirohi kids under farmer's management system. *Indian J Small Rumin.* 2010;16.
 48. Kharkar K, Kuralkar SV, Kuralkar P, Bankar PS, Chopade MM, Hadole KA. Factors affecting body weight and morphometric characters of Berari goats. *Indian J Small Rumin;* 2014.
 49. Bosso NA, Cisse MF, Van der Waaij EH, Fall A, Van Arendonk JAM. Genetic and phenotypic parameters of body weight in West African Dwarf goat and Djallonke sheep. *Small Rumin Res;* 2007.
 50. Al-Shorepy SA, Alhadrami GA, Abdulwahab K. Genetic and phenotypic parameters for early growth traits in Emirati goat. *Small Rumin Res.* 2002;45.
 51. Zhang CY, Zhang Y, Xu DQ, Li X, Su J, Yang LG. Genetic and phenotypic parameter estimates for growth traits in Boer goat. *Livest Sci;* 2009.
 52. Al-Saef. Genetic and phenotypic parameters of body weights in Saudi Arabi goat and their crosses with Syrian Damascus goat. *Small Rumin Res;* 2013.
 53. Ballal KME, Ahmed MKA, Musa LMA. Estimates of phenotypic and genetic parameters of growth traits in the Sudanese Nubian goat. *Res J Anim Vet Sci.* 2008;3:9–14.
 54. Das SM, Rege JEO, Shibre M. Phenotypic and genetic parameters of growth traits of blended goats at Malya, Tanzania; 2013.
 55. Alade NK, Dilala MA, Abdulyekeen AO. Phenotypic and genetic parameter estimates of litter size and body weights in goats. *Int J Sci Nat.* 2010;1(2).
 56. Bedhane M, Haile A, Dadi H, Alemu T. Estimates of genetic and phenotypic

- parameters for growth traits in Arsi-Bale goat in Ethiopia. J Anim Sci. 2013;3: 439–48.
57. Rashidi A, Sheikahmadi M, Rostamzadeh J, Shrestha JNB. Genetic and phenotypic parameter estimates of body weight at different ages and yearling fleece weight in Markhoz goats. Asian-Aust J Anim Sci. 2008;21(10).
58. Xiong Q, Chai J, Li XF, Suo XJ, Chen MX. Two tagSNPs in the prolactin receptor gene are associated with growth and litter traits in Boer and Macheng Black crossbred goats. Livest Sci; 2016.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://prh.globalpresshub.com/review-history/1765>