



Fungal Organisms Associated with Post-harvest Rot of Frafra Potatoes [*Solenostemon rotundifolius* (Poir.)] in Bongo-Soe, Upper East Region, Ghana

S. Apuri^{1*}, C. Kwoseh², E. A. Seweh³ and I. O. Tutu⁴

¹*Department of Ecological Agriculture, School of Applied Science and Art, Bolgatanga Polytechnic, Bolgatanga, Ghana.*

²*Department of Crop and Soil Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.*

³*Department of Agricultural Engineering, School of Engineering, Bolgatanga Polytechnic, Bolgatanga, Ghana.*

⁴*Crops Research Institute, Fumesua, Kumasi, Ghana.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJA/2019/v11i330061

Editor(s):

(1) Dr. Afroz Alam, Associate Professor, Coordinator UG (Botany) and PG (Plant Science) Department of Bioscience and Biotechnology, Banasthali University (Accredited A grade by NAAC) P.O. Banasthali Vidyapith- 304 022 (Rajasthan), India.

(2) Prof. Atilgan Atilgan, Department of Farm Structures and Irrigation, Faculty of Agriculture, Suleyman Demirel University, Turkey.

Reviewers:

(1) G. Anaukwu, Chikodili, Nnamdi Azikiwe University, Nigeria.

(2) Joaquin Guillermo Ramirez Gil, Universidad Nacional de Colombia at Medellin, Colombia.

(3) A. Ogunlade, Clement, Adeleke University, Nigeria.

(4) Soraya Gabriela Kiriachek, Universidad Nacional de La Pampa, Argentina.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/42921>

Short Research Article

Received 14 September 2018

Accepted 23 November 2018

Published 26 July 2019

ABSTRACT

Aims: The study aims to identifying the microorganisms associated with post-harvest rot of frafra potatoes in Bongo-soe, Upper east region of Ghana.

Place and Duration of Study: Department of Horticulture and the Pathology laboratory of the Faculty of Agriculture, Kwame Nkrumah University of Science and Technology (KNUST), Kumasi, Ghana. The *Solenostemon rotundifolius* tubers were stored at the Horticulture Department laboratory while rot identification was carried out at the Pathology laboratory of the Faculty of Agriculture. The Tubers were stored from 2nd November, 2012 to 22nd March 2013.

*Corresponding author: E-mail: samuelapuri@gmail.com;

Methodology: Four hundred (400) tubers of black cultivar and four hundred (400) tubers of a brown cultivar of *Solenostemon rotundifolius* tubers showing visible signs of rot during the storage were collected. Pieces of diseased tissues from the margin of the necrotic collected and immersed in 10% commercial bleach solution for sterilisation, for one minute. These were then blotted dry and plated on Potato Dextrose Agar PDA. The plates were sealed with a cellotape until growth occurred.

Results: The microorganisms identified to be responsible for causing rot in *Solenostemon rotundifolius* tubers were six in number. *Colletotrichum gloeosporioides* was identified to be responsible for 30.76% of rots observed, followed by *Aspergillus niger*, 23.07%, *Curvularia lunata*, 19.23%, *Aspergillus flavus*, 11.54%, *Trichoderma sp* and *Penicillium sp* both recorded 7.70% of rots observed. The percentage incidence of *Aspergillus niger* (15.38%), *Curvularia lunata* (11.54%) and *Aspergillus flavus* (7.69%) was higher in the black cultivar as compared with the brown cultivar which had percentage incidence of 7.69%, 7.69% and 3.85% respectively. Also, the percentage incidence of *Colletotrichum gloeosporioides* (15.38%) and *Penicillium sp* (3.85%) was the same in both the black and brown cultivars of *Solenostemon rotundifolius* tubers used in this study.

Conclusion: The activities of the damaging microorganisms can be reduced by controlling mechanical injury during harvesting, transportation and storage of *Solenostemon rotundifolius* tubers should be prevented or reduced because they pave the way for tuber infection by the rot causing microorganisms.

Keywords: Fungal organisms; post-harvest rot; frafra potatoes; *Solenostemon rotundifolius*.

1. INTRODUCTION

Solenostemon rotundifolius (Frafra potato) belongs to the family Lamiaceae (Labiatae) [1]. Frafra potatoes are herbaceous perennials normally cultivated as annuals [2]. In Ghana, the crop is mainly cultivated in the guinea and Sudan savannah agro-ecological zones [3]. The areas of cultivation mainly cover the Upper East and Upper West regions of Ghana. However, the crop can also be cultivated in the moist semi deciduous forest ecology of Ghana [3]. *Solenostemon rotundifolius* is an important food security crop [4,2]. Apart from cultivation of frafra potatoes for consumption, the crop also has medicinal, industrial and sociocultural importance in areas of its cultivation in Ghana [5,6,7].

In spite of its importance, frafra potato cultivation appears to be declining in Ghana due to the numerous problems encountered by farmers during cultivation. These include post-harvest loss during storage [8,9] which contributes to the immensely to spoilage (rotting) of tubers in storage [6] Sharma, 2016.

Postharvest spoilage (rot) is a major contributor to losses of all root and tuber crops via microbial infestation of healthy tubers which reduces the table quality of such tubers and renders them unappealing to consumers [10,11].

Roots and tubers have been subjected to several diseases by several post-harvest rot causing

organisms including viruses, fungi, bacteria, nematodes etc. For instance in yam (*Dioscoreae Sp*), the microorganisms commonly associated with storage rots include: *Sclerotium rolfsii*, *Colletotrichum gloeosporioides*, *Aspergillus spp*, *Botryoidiplodia theobromea*, *Curvularia verruculosa*, *Rhizoctonia solani*, *Penicillium spp* and *Fusarium moniliforme* [12,13]. Also in Sweet potatoes (*Ipomoea batatas*), pathogens commonly associated with storage rots include: *Ceratocystis fimbriata* (a fungus that causes Black rot), *Fusarium* (a fungus that causes Fusarium surface rot and Fusarium root rot), *Rhizopus stolonifer* (a fungus that causes Rhizopus soft), *Diplodia gossypina* (a fungus that causes Java black rot), *Erwinia chrysanthemi* (a bacteria that causes Bacterial soft rot, also known as bacterial stem and root rot), *Monilochaetes infuscans* (a fungus that causes scurf, also known as soil-stain), *Macrophomina phaseoli* (a fungus that causes Charcoal rot) [14]. However, for *Solenostemon rotundifolius*, no published research work has been cited on the microorganisms associated with post-harvest rot until recently. Mohammed et al. [15] indicates that the following microorganisms are associated with post-harvest rot of *Solenostemon rotundifolius* in Nigeria; *Aspergillus niger*, *Fusarium oxysporum*, *Penicillium expansum* and *Rhizopus stolonifer*. However, no research work has been cited on the microorganisms associated with post-harvest rot of *Solenostemon rotundifolius* in Ghana. This research work, therefore, has the overall objective of identifying the microorganisms associated with post-harvest

rot of frafra potatoes in Bongo-soe, Upper east region of Ghana.

2. MATERIALS AND METHODS

2.1 Geographical Location of the Experimental Site

The research work was carried out at the laboratory of the Department of Horticulture and the Pathology laboratory of the Faculty of Agriculture, Kwame Nkrumah University of science and Technology (KNUST), Kumasi, Ghana.

The *Solenostemon rotundifolius* tubers were stored at the Horticulture Department laboratory while rot identification was carried out at the Pathology laboratory of the Faculty of Agriculture.

2.2 Procurement of Frafra Potato (*Solenostemon rotundifolius*) Tubers and Preparation

The black and brown cultivars of frafra potato tubers used for this research were all obtained from a single farm in Bongo-soe, in the Bongo district of the Upper East region of Ghana. The farm was monitored from planting to harvest, fresh tubers were subsequently obtained on the day of harvest for experimentations. A total of eight hundred samples of frafra potato tubers were used for the studies. This was made up of four hundred black cultivars and four hundred brown cultivars of wounded and non-wounded tubers, the samples were visually inspected for any sign of rot and defective samples were discarded. The tubers were stored in sixteen medium sized plastic bowls which contained fifty tubers each.

2.3 Identification of Rot Pathogens

The Tubers were stored for 141 days, from 2nd November, 2012 to 22nd March 2013. Tubers showing visible signs of rot were taken to the Pathology laboratory of the Faculty of Agriculture for identification of the possible microorganisms responsible for the rot. This was done fortnightly.

The procedure involved taking pieces of diseased tissues (2-5 mm square) from the margin of the necrotic lesion using a sterile scapel blade and surface sterilising of the necrotic tissue segments done by immersing the

tissue segments in 10% commercial bleach solution in a beaker for one minute. Tissue segments were taken out of the sterilant solution in intervals of 10 seconds and blotted dry with a clean paper towel. This ensured that at least some of the segments were exposed to the sterilant for the appropriate period of time. The surface sterilised tissue segments were then plated on Potato Dextrose Agar (PDA) medium, using a pair of forceps which was flamed sterilised periodically. The plates were sealed with a sellotape to prevent desiccation and contamination and incubated at room temperature until growth occurred. The plates were then observed daily for fungal growth from the tissues. The characteristics of the spore and mycelium were studied using a compound microscope. The characteristics observed were used in identifying the rot organisms according to standards described by Mathur and Kongsdal [16] and Barnett and Hunter [17].

3. RESULTS AND DISCUSSIONS

Five genera of microorganisms were identified to be associated with post-harvest rot of stored frafra potato tubers. These are: *Curvularia* sp., *Colletotrichum* sp., *Aspergillus* sp., *Penicillium* sp. and *Trichoderma* sp.

The microorganisms identified to be responsible for causing Post-harvest rot in *Solenostemon rotundifolius* tubers were six in number. *Colletotrichum gloeosporioides* was responsible for 30.76% of rots observed, *Aspergillus niger*, 23.07%, *Curvularia lunata*, 19.23%, *Aspergillus flavus*, 11.54%, *Trichoderma* sp. and *Penicillium* sp. both recorded 7.70% of rots observed.

The percentage incidence of *Aspergillus niger* (15.38%), *Curvularia lunata* (11.54%) and *Aspergillus flavus* (7.69%) was higher in the black cultivar as compared with the brown cultivar which had percentage incidence of 7.69, 7.69 and 3.85% respectively. Also, the percentage incidence of *Colletotrichum gloeosporioides* (15.38%) and *Penicillium* sp (3.85%) was the same in both the black and brown cultivars of *Solenostemon rotundifolius* tubers used in this study. *Trichoderma* sp. caused rot in only tubers of the brown cultivar with percentage incidence of 7.07%. In all, tubers of the black cultivar had more incidence of rot causing microorganisms (53.84%) compared with tubers of the brown cultivar which had a total incidence of 46.16%.

The identification of *Aspergillus niger* and *Penicillium* sp. confirmed the observations made by Mohammed et al. [15]. This is the first time *Colletotrichum gloeosporioides*, *Curvularia lunata*, *Aspergillus flavus*, and *Trichoderma* sp. have been found to cause post-harvest rot in *Solenostemon rotundifolius* tubers. *Curvularia lunata*, *Colletotrichum gloeosporioides*, *Aspergillus niger*, *Aspergillus flavus*, *Penicillium* sp. and *Trichoderma* sp. have been found to be associated with rot in other roots and tuber crops such as Yam [12,13] sweet potato [18], cassava [19]. These microorganisms identified are wound pathogens that infect mechanically injured tubers and have been found to infect tubers in the field and later manifest in storage [18,20]. These microorganisms can also infect tubers through diseased foliage, mother tubers or roots [21].

The skin of *Solenostemon rotundifolius* tubers is easily damaged [15] and farmers make matters worse with poor practices that encourage mechanical injury on tubers during harvesting, transportation and even storage. This therefore explains the presence of these microorganisms in the *Solenostemon rotundifolius* (frafra potatoes) tubers used in this study.

4. CONCLUSION

The findings of this study have shown that *Colletotrichum gloeosporioides*, *Aspergillus flavus*, *Aspergillus niger*, *Curvularia lunata*, *Penicillium* sp. and *Trichoderma* sp. are responsible for the post-harvest rot of *Solenostemon rotundifolius* tubers in Bongo-soe, Upper East Region of Ghana.

5. RECOMMENDATION

Mechanical injury during harvesting, transportation and storage of *Solenostemon rotundifolius* tubers should be prevented or reduced since mechanical injuries can pave the way for tuber infection by the rot causing microorganisms identified in this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tindall HD. Vegetables in the tropics. Macmillan Press, London, United Kingdom. Washington, D.C: IFPRI. 1983;533.
2. National Research Council, NRC. Native potatoes. Lost Crops of Africa: Vegetables. National Academies Press. 2006;II:268-285. [ISBN 978-0-309-10333-6] Available: http://books.nap.edu/openbook.php?record_id=11763&page=269.
3. Opoku-Agyeman MO, Bennett-Lartey SO, Vodouhe RS, Osei C, Quarcoo E, Boateng SK, Osekere EA. Morphological characterization of frafra potato (*Solenostemon rotundifolius*) germplasm from the savannah regions of Ghana. Plant genetic resources and food security in West and Central Africa. Regional Conference, Ibadan, Nigeria. 2004;116-123.
4. Burkill HM. The useful plants of west tropical Africa. J-L. Royal Botanic Gardens, Kew. 1995;3.
5. Apaboh RR. Assessment of the performance of some frafra potato (*Coleus dysentericus* – Baker) Accessions in Nyankpala area of Ghana. A Dissertation submitted to the Faculty of Agriculture, UDS, in partial fulfillment of the requirements for the award of BSc Agric. Technology; 1997.
6. Tetteh JP, Guo I. Problems of Frafra potato (*Solenostemon rotundifolius* Poir.) production in Ghana. Ghana J. Agric. Sci. 1997;30:107-113
7. Umar AM. Review on the nutritional value, cultivation and utilization potential of some minor and under-utilized indigenous root and tuber crops in Nigeria. International Journal of Advanced Research. 2016;4: 1298-1303.
8. Tindall Sharma HK. Tropical roots and tubers: Production, Processing and Technology. John Wiley & Sons; 2016.
9. Vowotor A, Kwame, Mensah-Bonsu, Akwasi, Mutungi, Christopher, Affognon, Hippolyte. Postharvest losses in Africa – Analytical review and synthesis: The case of Ghana; 2012.
10. Amusa NA. Concentric leaf spot of yam (*Dioscorea* sp.) in South-western Nigeria. Mycopathologia. 1999;148:33-36.
11. Anjorin TS, Nwokocho OV, Sanni AD. Morphological characteristics and incidence of diseases on white yam (*Dioscorea rotundata* L. Poir) tubers in Abuja, Nigeria. Nat Sci. 2014;12(7):58-65.
12. Nwankiti AO, Okpala EU. Anthracnose of water yam in Nigeria. In: Proceedings of

- the 6th Triennial Symposium of the International Society for Tropical Root Crops, Peru; 1983.
13. Green KR. Studies on the epidemiology of yam anthracnose. PhD Thesis, University of Reading; 1994.
 14. Sikora EJ, James MD. Field and storage diseases of sweet potatoes; 1995. Available:<http://www.aces.edu/pubs/docs/A/ANR-0917/> (Accessed on 04/06/2013)
 15. Mohammed A, Ishaku BC, Basiri B. Identification and control of Fungi associated with the post-harvest rot of *Solenostemon rotundifolius* (Poir) J.K. Morton in Adamawa State of Nigeria. Journal of Biology, Agriculture and Healthcare. 2013;3(5). [ISSN 2224-3208 (Paper)] [ISSN 2225- 093X (Online)].
 16. Marthur SB, Kongsdal O. Common laboratory seed health testing methods for detecting fungi, 2nd Edition. International Seed Testing Association. Switzerland; 2003.
 17. Barnett HL, Hunter BB. Illustrated genera of imperfect fungi. 3rd Edition. Burgess Publishing Company; 1972.
 18. Rees D, van Oirschot Q, Kapinga R. (eds) Sweetpotato post-harvest assessment: experiences from East Africa, Natural Resource Burkill s Institute, Chatham, UK. 2013;548(2):51-66. ISBN: 0 85954
 19. Messiga AJNA, Mwangi M, Bandyopadhyay R, Nolte C. The status of fungal tuber rots as a constraint to cassava production in the Pouma district of Cameroon. Paper presented at the proceedings of the 9th Triennial Symposium of the International Society for Tropical Root Crops - Africa Branch, held from 31st October – 5th November 2004, at Whitesands Hotel, Mombasa, Kenya; 2004.
 20. Okigbo RN, Agbata CA, Echezona CE. Effects of leaf extracts of *Azadirachta indica* and *Chromolaena Odorata* on post-harvest spoilage fungi of yams in storage. Current Research Journal of Biological Sciences. 2010;2(1):9-12.
 21. Nmeka IA, Okigbo RN. Control of yam tuber rot with leaf extracts of *Xylopi aethiopica* and *Zingiber officinale*. African Journal of Biotechnology. 2005;4(8):804-807.

© 2019 Apuri et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/42921>