



Impact of Introduced Tomato Cultivars and Chemical Fungicides on the Occurrence and Intensity of Late Blight in Western Cameroon

Dietchou Paolo Vincent^{a*}, Keuete Kamdoum Elie^b and Tsopmbeng Numbo Gaston^a

^a Department of Plant Biology, Faculty of Science, University of Dschang, P.O. Box 67, Dschang, Cameroon.

^b Department of Plant Science, Faculty of Science, University of Buea, P.O. Box 63, Buea, Cameroon.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2023/v45i92177

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102039>

Original Research Article

Received: 30/04/2023

Accepted: 06/07/2023

Published: 21/07/2023

ABSTRACT

Late blight, caused by *Phytophthora infestans*, is the main constraint to tomato production and exportation in Cameroon. The objective of this study was to evaluate the impact of chemical and tomato varieties in the control of tomato Late blight in the field for the improvement of tomato production in the west region of Cameroon. Seedlings of 7 tomato varieties produced in the nursery were transplanted into pits each containing 100 g of fowl droppings and 30 g of the nematicide Carbofuran 5%. From the 21st day after transplanting, the different fungicides were applied every four days. Data were collected on the incidence, severity of the disease at different stages of plant

*Corresponding author: E-mail: dpaolovincen@yahoo.fr;

growth and development and on yields. The results showed that the AVTO1219 and AVTO1311 varieties were less sensitive to Late blight. The control plants of AVTO1219 variety showed incidences of 7.7, 11.7 and 16.7% and severities of 1.7, 14.9 and 17.2% in the vegetative, flowering and fruiting phases respectively. Control plants of CLN1462A, CLN1464A, CLN1464B, RIOGRANDE 2 and RIOGRANDE + varieties showed incidence of 100% from the flowering phase. Control plants of varieties AVTO1213 (17.29 t ha⁻¹) and AVTO1311 (11.33 t ha⁻¹) produced the highest yields. Bonsoin was the more effective fungicide and AVTO1311 was the more effective variety to manage disease. However, the combination of Bonsoin with the AVTO1219 and AVTO1311 varieties was most effective in controlling late blight of tomatoes in the field. These results suggest that to improve tomato production, Late blight resistant varieties should be used and effective fungicides applied as treatments.

Keywords: Tomatoes cultivars; fungicides; occurrence; intensity; late blight.

1. INTRODUCTION

Tomato (*Solanum lycopersicum* L.) is an important vegetable crop grown around the world and is the second next to potato and it belongs to the family Solanaceae. It originated from tropical Mexico to Peru [1]. Tomato is a source of minerals, vitamins, lycopene and health benefits in reduce cancer and heart disease [2,3]. Economically, it is the fourth most important crop in the world after rice, wheat, and soybean [4]. In Cameroon, the production of tomato increases and become a source of cash income for producers, and retains its importance for daily household consumption.

The production of the crop is attrition by several biotic and abiotic factors. Among biotic constraints, fungal diseases are major factors affecting production and productivity as well as the quality of the crop. Many fungal diseases like late blight, *Fusarium* wilt, are attacking this crop, and can cause 15-95% crops loss both in lowland and highland areas of the tropics [5,6]. Late blight of tomato, caused by *Phytophthora infestans*, is one of the most serious diseases of this crop. It is present all over the world where tomato cultivation exists and even remains the main limiting factor for tomato production worldwide [7]. Late blight can also cause damage to other wild and cultivated Solanaceae, especially *Solanum tuberosum*, *S. melongena*, *S. torvum* and *S. nigrum* [8,9,10].

In Cameroon and in the West Region in particular, tomato farmers have noted this through disease outbreaks, once in the field, this disease increased the difficulties of maintaining Late blight-free tomato plots [11]. This disease is responsible for damages, affecting mainly the aerial parts of the plant (leaves, stems, flower or fruits). Generally, lesions on the leaves first

appear as irregular dark spots that widen as new lesions develop [12]. On the upper surface of the attacked leaves, a lighter green halo often surrounds the necrotic area, on the lower surface, a milky white ring sporulation develops around the lesions, if the humidity is favourable for the development of the pathogen. Stem lesions are elongated, greyish-brown to black in colour and often encircle the stem [13]. These lesions are often found in the axils of bracts and at the apex of stems [14,4]. On the floral clusters (or inflorescences), the disease causes browning and then the fall of many flowers.

On the fruits, Late Blight is characterized by irregular brown patches with a tan and shiny appearance. Thus, the fruit has a bumpy appearance with an interior corky texture and a brown colour and may also incur the appearance of white fuzz on the infected parts [15,1]. Infected unripe green fruits turn brown on the surface. These lesions develop and become enlarged brown and leathery in appearance. Hence, only those partially attacked will mature [8,7]. It can cause up to 90% of crop losses in cool and wet weather conditions, most prevalent during the rainy season and cause yield losses of up to 100% and fruit losses up to 30-60% [16,10].

Management of late blight requires aggressive measures that include combined use of cultural, scouting, sanitation, and most importantly the combination of host plant resistance with application of fungicides. Many growers also use different fungicides like Mancozeb, Agrolaxyl, Metalaxy, copper, phosphorus acid and Ridomil for the control of late blight of tomato [14,17,18]. Among many alternative measures available, use of resistant varieties has been the most important and a cost-effective approach for the management of the disease [6].

Thus, the objective of this study was to evaluate the effects of fungicides and tomato varieties on controlling Late Blight in tomatoes.

2. MATERIALS AND METHODS

2.1 Description of the Experimental Site

The field experiment was conducted under the main rainy season in the West Region of Cameroon, Ndé Division, precisely in the Bangangté Sub-division. This region is the main tomato production basin [19]. Bangangté is located between latitude 5°5' and 6°5' North and between longitude 10°5' and 11°5' East. It has an average altitude of 1432 m [20] and belongs to the agro-ecological zone of the Western Highlands of Cameroon, which is characterized by a humid tropical climate with two seasons, a short dry season and a long rainy season. Temperatures vary between 14° and 28°C with an annual average of 22°C. Rainfall varies between 1400 and 2500 mm per year. The soil is fertile and has a sandy-clay texture; with a slightly acidic pH [21].

2.2 Planting Materials and Fungicides

The experiment included two factors: tomato varieties and fungicides. Tomato varieties AVTO1219, AVTO1311, CLN1462A, CLN1464A and CLN1464B (exotic) and RIOGRANDE+ and RIOGRANDE2 (local) were obtained from World Vegetable Center of Yaounde. The Three chemical fungicides, one systemic Bonsoin (Chlorothalonil 30% and Cymoxanil 6%) and two contact: Mancostar (Mancozeb 80 WP) and Plantineb (Plantneb 80 WP), were used. These fungicides were purchased from a phyto-sanitary center in the town of Bangangté.

2.3 Experimental Design and Treatment

Tomato seedlings of 12 to 15 cm in height, previously produced in the nursery, were transplanted inside pits 50 cm apart and containing 100 g of fowl droppings as a background fertilizer [22] and 30 g Carbofuran 5% (nematicide) [19]. These pits were made on 5 m long and 1 m wide ridges. Seven days after transplanting (DAT), Yara Mila (12-11-18) and




Fig. 1. Location map of the study area

Urea (46% N) fertilizers were applied at the manufacturer's rate as starter fertilizer. Plant maintenance was carried out whenever necessary through weeding, hoeing and ridging. At 37 (DAT), Yara Mila (12-11-18) and potassium sulphate (50% K₂O -12% S) were applied at a concentration of 240 kg ha⁻¹ [22,23]. Staking of the plants was done using bamboos before fruiting i.e. 40 DAT. The insecticide Tamega was applied at 7 DAT; as well as once a week from the fruiting period until the ripening of tomato fruits at the concentration of 1 l/ha. From the 21st DAT, phyto-sanitary treatments were made every four days with the different fungicides at the manufacturer's concentration (2.5 kg ha⁻¹) to control Late Blight [1]. The experiment was laid out in a randomized complete block design, split-plot arrangement with 24 treatments replicated thrice.

2.4 Data Collection

2.4.1 Disease incidence and severity

Data on the disease incidence and severity of Late blight of tomato were collected at three stages of tomato plant development. These developmental stages were: vegetative phase (38 DAT), flowering phase (50 DAT) and fruiting phase (62 DAT). Disease incidence was recorded by counting of plants that showing visible symptoms of late blight in the central rows and the data were expressed as a percentage of the total assessed plants. The disease incidence was calculated with the following formula [24]:

$$\text{Disease incidence (\%)} = \frac{\text{Number of infected tomato plants}}{\text{Total number of tomato plants examined}} \times 100$$

Disease severity was assessed on the central two rows and recorded at different stages of development. Ten plants were selected randomly from each experimental plot, and then five leaves of each plant were used to determine the disease severity. Severity of late blight was recorded on the basis of 0-5 rating scales as described by De Putter et al. [25] where scale 0 = no symptoms, 1 = Nearly 10 % of the leaves of the whole plant are infected, 2 = Nearly 25% of the leaves of the whole plant are infected, 3 = Nearly 50% of the leaves of the whole plant are infected, 4 = Nearly 75 % of the leaves of the whole plant are infected and 5 = All the leaves of the plant are infected and death of the plant. severity (S) was calculated by the following formula:

$$\text{Disease severity (\%)} = \frac{\sum(Xi \times ni)}{N \cdot Z} \times 100$$

Where:

Xi = Severity i of disease in tomato plant
 ni = Number of tomato plants with severity i
 N = Total number of examined tomato plant
 Z = Highest score of the scale, 5

2.4.2 Area under disease progressive curve and disease progress curve (AUDPC)

The effect of varieties and chemical fungicides combinations on disease severity data was integrated into area under disease progress curve (AUDPC), as described by Campbell and Madden [26].

$$\text{AUDPC} = \sum_{i=1}^n 0.5 (x_{i+1} + x_i) (t_{i+1} - t_i)$$

Where, n is the total number of assessments, 'ti' is the time of the ith assessment in days from the first assessment date; 'xi' is percentage of disease severity at ith assessment. AUDPC is expressed in percent-days because the severity (x) was expressed in percent and time (t) in days. AUDPC was expressed in percent-days because the severity (x) was expressed in percent and time (t) in days. The rates of disease progress in time was determined by recording the severity of Late Blight at 4 days' interval right from the appearance of the first disease symptoms (28 DAT) till the maturity of the crop in the different treatments.

2.4.3 Assessment of yield

At maturity, tomato fruits were harvested from the central three rows on each plot of each treatment. Additionally, the weights of marketable yield of tomato fruits per plot were recorded. Fruits yield per plot was converted into yield of tons per hectare.

2.4.4 Data analysis

Data on disease parameters (disease incidence, disease severity and AUDPC) and yield component were subjected to analysis of variance (ANOVA). The means were separated through the test of Student Newman-Keuls (SNK) at a 5%. This was done using SPSS software version 21.0.

3. RESULTS

3.1 Effect of Tomato Varieties and Chemical Fungicides on Late Blight Incidence

Late blight of tomato was present at all stages of development of the plant. Incidence (in %) of this disease varied according to the fungicide applied, the tomato variety used, as well as the developmental stage of the plant (Table 1). At the vegetative phase (38 DAT), the lowest incidence with Bonsoin was obtained with the AVTO1219 variety (3.3%) and the highest with the RIOGRANDE+ variety (15.7%).

Statistical analyses showed that the incidence obtained with Bonsoin on the AVTO1219 (3.3%) and AVTO1311 (7.3%) varieties were significantly identical and lower than those obtained on the other varieties. Meanwhile Plantneb, CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and RIOGRANDE2 varieties had significantly identical and higher incidences than AVTO1219 (7.3%) and AVTO1311 (9.3%) varieties. On the other hand, within the same variety, no significant difference was observed between plants of the AVTO1219 (7.7%) and AVTO1311 (8.7%) varieties sprayed with the different fungicides according to Student Newman-Keuls test at 5% probability threshold. With the RIOGRANDE+ variety, the highest incidence was observed in Plantneb (50.7%) and the lowest in Bonsoin (15.7%) and Mancozeb (22.3%).

For the untreated plants, the smallest incidences were obtained with the AVTO1219 (7.7%) and AVTO1311 (8.7%) varieties and the highest with the CLN1462A (61.6%) and RIOGRANDE2 (62.7%) varieties respectively.

In the flowering phase, incidence (in %) of Late blight showed significant difference between the different treatments. Two groups were observed; One group consisted of the AVTO1219 and AVTO1311 varieties, characterized by the smallest incidences. These incidences ranged from 5.3 to 14.3%. The second group composed of the CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and RIOGRANDE2 varieties, characterized by larger incidences ranging from 16 to 56.8% (Table 2).

The varieties AVTO1219 and AVTO1311 showed significantly identical incidences between fungicides. These incidences varied from 5.3% to

9.0% for the AVTO1219 variety and from 7.3% to 14.3% for the AVTO1311 variety. With the CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and RIOGRANDE2 varieties, Student Newman-Keuls test at 5% threshold showed that these different varieties could be grouped into two. The first group made of plants sprayed with Mancozeb and Plantneb, characterized by higher incidences that varied from 40.1 to 56.8%. The second group of plants sprayed with Bonsoin characterized by lower incidences, ranging from 16 to 22.7%.

With the control plants, all plants of CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and RIOGRANDE2 varieties were affected by the disease (100%). The varieties AVTO1219 (11.7%) and AVTO1311 (20.3%) showed the lowest incidence.

According to Table 3, during fruiting, plants sprayed with Mancozeb showed the highest incidences with CLN1462A (70.3%) and RIOGRANDE 2 (63.1%) varieties and lowest incidences with AVTO1311 (12.7%) and AVTO1219 (9.3%) varieties. Bonsoin the CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and RIOGRANDE2 varieties showed significantly identical and higher incidences than the AVTO1219 (6.7%) and AVTO1311 (7.3%) varieties. These incidences ranged from 18.5% to 26.7%. With Plantneb, the highest incidence was observed on the CLN1464B variety (76.7%) followed by CLN1464A (69.3%) and RIOGRANDE2 (65.3%) varieties. The AVTO1219 variety showed the lowest incidence (12.7%).

Within the same variety CLN1462A, the incidences were 70.3, 57.3 and 26.7% for Mancozeb, Plantneb and Bonsoin sprayed plants respectively. With the CLN1464B variety, the highest incidence was obtained on the Mancozeb fungicide (76.7%) and the lowest on plants sprayed with the fungicide Bonsoin (18.7%). The Mancozeb (12.7%) and Plantneb (17.7%) fungicide sprayed plants of the AVTO1311 variety showed significantly similar and higher incidences than the Bonsoin sprayed plants (7.3%) according to the Student Newman-Keuls test at 5% threshold. Plants of the RIOGRANDE2 variety sprayed with Bonsoin showed the lowest incidence (20%) and the plants of the same variety sprayed with Mancozeb showed the highest incidence (63.1%).

The plants of the varieties CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and

RIOGRANDE2 showed significantly identical and highest incidence of 100%. While the AVTO1219 and AVTO1311 varieties showed the lowest incidences of 16.7% and 24.3% respectively.

3.2 Effects of Chemical Fungicides and Tomato Varieties on the Severity of Late Blight

Severity (in %) varied according to the fungicide applied, the tomato variety and the development stage of the plant (Table 4, 5 and 6). During the vegetative phase (Table 4), with plants sprayed with Plantneb, the highest severity was observed on plants of the RIOGRANDE+ variety (17%) and the smallest on the CLN1464B variety (5.8%) followed by the AVTO1219 (2.3%) and AVTO1311 (1.7%) varieties. With Bonsoin fungicide, the CLN1462A, CLN1464A, CLN1464B, RIO GRANDE + and RIO GRANDE 2 varieties showed severities between 2.4 and 5.3%. These different severities were higher than those of the AVTO1219 (1.3%) and AVTO1311 (1%) varieties.

Within the AVTO1219 and AVTO1311 varieties, the severities of the sprayed plants regardless of the fungicide applied were significantly identical according to Student Newman-Keuls test at 5% probability threshold. With the CLN1462A variety, plants sprayed with Mancozeb (6.9%) and Plantneb (7.2%), showed similar and higher severities than plants sprayed with Bonsoin (3.9%). Plants sprayed with Bonsoin (5.3%), Plantneb (5.8%) and Mancozeb (6.8%) had the lowest severity.

With the control plants the highest severities were obtained on the CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and RIOGRANDE2 varieties. These severities ranged from 8.9% to 15.8%. The AVTO1219 variety showed the lowest severity (1.9%) and CLN1464A variety showed the highest severity (15.8%).

During flowering, Bonsoin sprayed plants showed varying degrees of severity (Table 5). The highest severities were observed on the RIOGRANDE+ variety (20.4%) followed by the

Table 1. Incidence of Late blight on different tomato in the vegetative phase (38 DAT) with respect to the chemical fungicides and the tomato varieties used

Varieties	Treatments			
	Bonsoin	Mancozeb	Plantineb	Control
AVTO1219	3.3 ± 5.8 ^{ba}	1.7 ± 2.9 ^{ca}	7.3 ± 8.7 ^{ba}	7.7 ± 7.6 ^{ba}
AVTO1311	7.3 ± 6.4 ^{abA}	6.7 ± 7.6 ^{bcA}	9.3 ± 8.1 ^{ba}	8.7 ± 5.5 ^{ba}
CLN1462A	13.0 ± 7.0 ^{aC}	26.5 ± 8.5 ^{abC}	38.7 ± 19.5 ^{aAB}	61.6 ± 10.4 ^{aA}
CLN1464A	17.0 ± 3.6 ^{ab}	22.7 ± 7.0 ^{abB}	31.7 ± 4.1 ^{ab}	56.3 ± 13.3 ^{aA}
CLN1464B	9.0 ± 5.2 ^{abC}	22.3 ± 13.8 ^{abBC}	32.2 ± 15.4 ^{ab}	54.3 ± 6.8 ^{aA}
RIOGRANDE+	15.7 ± 2.0 ^{ab}	29.4 ± 9.0 ^{ab}	50.7 ± 8.1 ^{aA}	51.0 ± 18.7 ^{aA}
RIOGRANDE2	12.7 ± 2.5 ^{abC}	23.7 ± 12.4 ^{abC}	45.6 ± 4.6 ^{ab}	62.7 ± 7.6 ^{aA}

a,b,c: means assigned the same lower case letter in the same column; as well as A,B,C: means assigned the same upper case letter in the same row are not significantly different according to Student Newman-Keuls test at 5%

Table 2. Incidence of Late blight on different tomato in the flowering phase (50 DAT) with respect to the chemical fungicides and the tomato varieties used

Varieties	Treatments			
	Bonsoin	Mancozeb	Plantineb	Control
AVTO1219	6.0 ± 8.7 ^{ba}	5.3 ± 9.5 ^{ba}	9.0 ± 8.5 ^{ba}	11.7 ± 7.6 ^{ca}
AVTO1311	7.3 ± 6.4 ^{ba}	12.7 ± 11.5 ^{ba}	14.3 ± 4.0 ^{ba}	20.3 ± 2.5 ^{ba}
CLN1462A	22.7 ± 8.7 ^{aC}	54.0 ± 18.1 ^{ab}	53.0 ± 15.1 ^{ab}	100 ± 0.0 ^{aA}
CLN1464A	19.3 ± 1.2 ^{aC}	44.7 ± 4.1 ^{ab}	45.9 ± 17.1 ^{ab}	100 ± 0.0 ^{aA}
CLN1464B	16.0 ± 5.3 ^{abC}	40.1 ± 11.5 ^{ab}	56.8 ± 16.7 ^{ab}	100 ± 0.0 ^{aA}
RIOGRANDE+	18.7 ± 4.0 ^{aC}	49.7 ± 8.1 ^{ab}	54.0 ± 12.1 ^{ab}	100 ± 0.0 ^{aA}
RIOGRANDE2	20.0 ± 5.0 ^{aC}	44.0 ± 9.8 ^{ab}	51.3 ± 11.8 ^{ab}	100 ± 0.0 ^{aA}

a,b,c: means assigned the same lower case letter in the same column; as well as A,B,C: means assigned the same upper case letter in the same row are not significantly different according to Student Newman-Keuls test at 5%

Table 3. Incidence of Late blight on different tomato in fruiting phase (62 DAT). With respect to the chemical fungicides and the tomato varieties used

Varieties	Treatments			
	Bonsoin	Mancozeb	Plantineb	Control
AVTO1219	6.7 ± 7.6 ^{bA}	9.3 ± 9.0 ^{CA}	12.7 ± 11.0 ^{bA}	16.7 ± 5.8 ^{CA}
AVTO1311	7.3 ± 6.4 ^{bb}	12.7 ± 11.0 ^{CAb}	17.7 ± 2.5 ^{bAb}	24.3 ± 5.1 ^{bA}
CLN1462A	26.7 ± 2.9 ^{aD}	70.3 ± 2.1 ^{ab}	57.3 ± 11.4 ^{aC}	100 ± 0.0 ^{aA}
CLN1464A	19.3 ± 1.2 ^{aD}	55.0 ± 7.9 ^{bC}	69.3 ± 5.5 ^{ab}	100 ± 0.0 ^{aA}
CLN1464B	18.7 ± 1.2 ^{aD}	52.3 ± 4.9 ^{bC}	76.7 ± 7.6 ^{ab}	100 ± 0.0 ^{aA}
RIOGRANDE+	18.5 ± 4.5 ^{aC}	55.2 ± 8.7 ^{bb}	65.3 ± 5.7 ^{ab}	100 ± 0.0 ^{aA}
RIOGRANDE2	20.0 ± 5.0 ^{aC}	63.1 ± 4.0 ^{abb}	61.7 ± 6.4 ^{ab}	100 ± 0.0 ^{aA}

a,b,c: means assigned the same lower case letter in the same column; as well as A,B,C: means assigned the same upper case letter in the same row are not significantly different according to Student Newman-Keuls test at 5%

Table 4. Severity of Late blight on different tomato in vegetative phase (38 DAT) with respect to the chemical fungicides and the tomato varieties used

Varieties	Treatments			
	Bonsoin	Mancozeb	Plantineb	Control
AVTO1219	1.3 ± 2.3 ^{bA}	1.7 ± 2.9 ^{bA}	2.3 ± 2.1 ^{CA}	1.7 ± 1.5 ^{CA}
AVTO1311	1.0 ± 1.0 ^{bA}	2.0 ± 3.5 ^{bA}	1.7 ± 1.5 ^{CA}	5.0 ± 3.0 ^{bA}
CLN1462A	3.9 ± 1.8 ^{abb}	6.9 ± 1.5 ^{abA}	7.2 ± 3.1 ^{bA}	8.9 ± 2.8 ^{abCA}
CLN1464A	2.4 ± 1.4 ^{abb}	7.3 ± 1.5 ^{abb}	7.4 ± 0.9 ^{bb}	15.8 ± 5.0 ^{aA}
CLN1464B	5.3 ± 1.6 ^{ab}	6.8 ± 3.1 ^{abb}	5.8 ± 0.5 ^{bb}	14.3 ± 4.1 ^{aA}
RIOGRANDE+	3.5 ± 0.9 ^{abb}	6.6 ± 1.7 ^{abAB}	8.8 ± 2.3 ^{bAB}	12.8 ± 7.7 ^{abA}
RIOGRANDE2	2.9 ± 1.5 ^{abb}	13.5 ± 10.6 ^{aAB}	17.0 ± 1.8 ^{aA}	12.9 ± 4.9 ^{abAB}

a,b,c: means assigned the same lower case letter in the same column; as well as A,B,C: means assigned the same upper case letter in the same row are not significantly different according to Student Newman-Keuls test at 5%

varieties RIOGRANDE2 (16.8%) and CLN1462A (14.6%) and the lowest on the AVTO1219 and AVTO1311 varieties where each had a severity of 1.5%. With the Mancozeb sprayed plants, the 5% threshold Student Newman-Keuls test showed the existence of two homogeneous groups. The group composed of the AVTO1219 and AVTO1311 varieties, whose peculiarity is the presence of the smallest severities of 3.3%.

The other group is made up of the CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and RIOGRANDE2 varieties, characterized by the highest severities (ranging from 22.6% to 35.8% respectively). Within the AVTO1311 variety, the highest severity was observed in the Plantneb sprayed plants (7.8%) and the lowest in plants sprayed with Bonsoin (1.5%). Plants sprayed with the fungicide—Mancozeb (3.3%) had intermediate severities.

Control plants showed higher severities than the plants sprayed with the different fungicides. The highest severities were obtained on the CLN1462A, CLN1464A, CLN1464B, RIO

GRANDE + and RIO GRANDE 2 varieties. These severities ranged from 14.9% to 69.6%. The AVTO1219 variety showed the lowest severity (14.9%) and Rio Grande 2 variety showed the highest severity (69.6%).

During fruiting (Table 6), plants sprayed with Plantneb showed the highest severities on the RIOGRANDE+ (39.9%) and CLN1462B (38.4%) varieties while the lowest severities were obtained with the AVTO1219 (5%) and AVTO1311 (7.8%) varieties. With the fungicide Mancozeb, the CLN1462A, CLN1464A, CLN1464B, RIOGRANDE+ and RIOGRANDE2 varieties had significantly similar and higher severities than the AVTO1219 (4.3%) and AVTO1311 (3.6%) varieties. These severities ranged from 30.1% to 41.9%. Within the RIOGRANDE+ and AVTO1311 varieties, the lowest severities were obtained with Bonsoin sprayed plants. These severities were 1.5% and 23.1% for the AVTO1311 and RIOGRANDE+ varieties respectively. For the AVTO1219, CLN1462A, CLN1464A, CLN1464B and RIOGRANDE2 varieties, the control plants

showed the highest severity and the plants sprayed with the different fungicides the lowest.

The control plants showed that the RIOGRANDE+ (98.5%) and RIO GRANDE 2 (91.5%) varieties had the highest severities and the AVTO1219 (17.2%) and AVTO1311 (15.1%) varieties the lowest (Table 6).

3.3 Area under Disease Progress Curve (AUDPC)

Area under disease progress curve (AUDPC) was highly affected by the tomato varieties. The severity of Late Blight was seen to increase with time. The smallest severity was observed on the 28th DAT and the highest severity was observed on the 62nd DAT whatever the variety (Fig. 2). The AVTO1311 and AVTO1219 varieties were the most sensitive to the disease compared to other varieties, showing the smallest AUDPC whatever the number of DAT. This AUDPC at 62 DAT was 62.2% and 68.9% respectively for

AVTO1311 and AVTO1219 varieties. The RIOGRANDE+ and CLN1464B varieties with the AUDPC of 393.8% and 387.1% respectively were highly sensitive to the disease.

The disease progress curves of tomato late blight (severity versus day after transplanting) for each chemical fungicide were presented separately.

Figs. 3, 4 and 5 shows the sum of severity of Late Blight increases with time. The smallest severity was observed on the 28th DAT and the highest severity was observed on the 62nd DAT whatever the variety. The CLN1464B and AVTO1311 varieties were the least sensitive to the disease compared to other varieties, showing the smallest AUDPC whatever the number of DAT. This AUDPC at 62 DAT was 63.5% and 6% respectively for CLN1464A and AVTO1311 varieties. The RIOGRANDE + and CLN1464A varieties with the AUDPC of 92.3% and 79.3% respectively were highly sensitive to the disease.

Table 5. Severity of Late blight on the different tomato in the flowering phase (50 DAT) with respect to the chemical fungicides and the tomato varieties used

Varieties	Treatments			
	Bonsoin	Mancozeb	Plantineb	Control
AVTO1219	1.5 ± 2.6 ^{CB}	3.3 ± 2.9 ^{BB}	5.0 ± 4.5 ^{CB}	14.9 ± 3.0 ^{BA}
AVTO1311	1.5 ± 2.6 ^{CC}	3.3 ± 2.9 ^{BBC}	7.8 ± 3.3 ^{CB}	15.1 ± 2.6 ^{BA}
CLN1462A	14.6 ± 9.7 ^{abB}	25.6 ± 14.5 ^{ab}	24.9 ± 10.1 ^{abB}	50.6 ± 18.5 ^{AA}
CLN1464A	10.6 ± 3.9 ^{BB}	22.6 ± 12.1 ^{ab}	33.3 ± 18.5 ^{abB}	62.3 ± 10.5 ^{AA}
CLN1464B	11.0 ± 1.1 ^{BB}	25.4 ± 7.2 ^{AB}	29.3 ± 27.7 ^{abAB}	59.2 ± 26.1 ^{AA}
RIOGRANDE+	20.4 ± 4.8 ^{abB}	35.8 ± 5.7 ^{ab}	36.4 ± 7.2 ^{ab}	64.1 ± 16.2 ^{AA}
RIOGRANDE2	16.8 ± 4.6 ^{abB}	33.3 ± 12.1 ^{ab}	34.3 ± 21.6 ^{abB}	69.6 ± 18.0 ^{AA}

a,b,c: means assigned the same lower case letter in the same column; as well as A,B,C: means assigned the same upper case letter in the same row are not significantly different according to Student Newman-Keuls test at 5%

Table 6. Severity of Late blight on the different tomato in fruiting phase (62 DAT) with respect to the chemical fungicides and the tomato varieties used

Varieties	Treatments			
	Bonsoin	Mancozeb	Plantineb	Control
AVTO1219	1.5 ± 2.6 ^{BB}	4.3 ± 2.9 ^{BB}	5.0 ± 4.5 ^{CB}	17.2 ± 1.5 ^{CA}
AVTO1311	1.5 ± 2.6 ^{BC}	3.6 ± 2.9 ^{BBC}	7.8 ± 3.3 ^{CB}	15.1 ± 2.6 ^{CA}
CLN1462A	19.8 ± 13.7 ^{ab}	30.7 ± 16.7 ^{ab}	26.5 ± 9.7 ^{BB}	85.4 ± 9.5 ^{BA}
CLN1464A	15.9 ± 1.5 ^{ab}	30.1 ± 6.8 ^{ab}	35.5 ± 17.4 ^{abB}	90.8 ± 8 ^{abA}
CLN1464B	16.8 ± 2 ^{ab}	38.7 ± 12.2 ^{ab}	38.4 ± 29.9 ^{ab}	96.8 ± 2.8 ^{abA}
RIOGRANDE+	23.1 ± 6.3 ^{aC}	41.9 ± 6.6 ^{ab}	39.9 ± 7.5 ^{ab}	98.5 ± 2.6 ^{AA}
RIOGRANDE2	19.2 ± 2.4 ^{ab}	36.2 ± 10 ^{ab}	37.3 ± 18.7 ^{abB}	91.5 ± 11.2 ^{abA}

a,b,c: means assigned the same lower case letter in the same column; as well as A,B,C: means assigned the same upper case letter in the same row are not significantly different according to Student Newman-Keuls test at 5%




Fig. 2. Area under the disease progression curve of different tomato varieties




Fig. 3. Area under the disease progression curve of different tomato plant treated by bonson

Chemical fungicide Mancozeb significantly reduced disease severity compared to the control (Fig. 4). The tomato varieties AVTO1219 and AVTO1311 showed the lowest AUDPC at 62 DAT. The AUDPC was 13.1%. Meanwhile the RIOGRANDE+ and CLN1464A varieties with the AUDPC of 170.8% and 160% respectively were highly sensitive to the disease at 62 DAT.

The AUDPC of tomato plants treated with Plantineb varied according to tomato variety and DAT (Fig. 5). AVTO1311 and AVTO1219 varieties were the least sensitive to the disease compared

to other varieties, having the smallest AUDPC of 31.1% and 20.1% respectively at 62 DAT. Meanwhile the CLN1464B and RIOGRANDE+ varieties with the AUDPC of 190.3% and 178.8% respectively were highly sensitive to the disease at 62 DAT.

3.4 Effects of Chemical Fungicides and Tomato Varieties on the Yields

All fungicides resulted in higher yields than the controls. These yields varied according to tomato variety and fungicide applied (Table 7). With the

fungicides Bonsoin and Mancozeb, the varieties CLN1462A and CLN1464A showed significantly higher yields than the other varieties. These yields ranged from 46.37 t ha⁻¹ to 53.99 t ha⁻¹. The RIOGRANDE2 variety showed the lowest yields. Yields were 35.7 and 30.56 t ha⁻¹ for Bonsoin and Mancozeb fungicides respectively. Of the plants treated with Plantineb, CLN464A showed the highest yield (45.33 t ha⁻¹) and AVTO1311 the lowest (23.59 t ha⁻¹).

As for the yield of the control plants, the variety AVTO1219 showed the highest yield (17.29 t ha⁻¹), followed by AVTO1311 (11.33 t ha⁻¹) and

CLN464A (3.27 t ha⁻¹). CLN464B and RIOGRANDE+, with yields of 1.87 and 1.83 t ha⁻¹ respectively, were the lowest yielding varieties.

4. DISCUSSION

Late blight caused by *Phytophthora infestans*, is a major constraint to tomato production. The incidence and severity of Late blight varied according to the chemical fungicide applied, the tomato variety used and the developmental phases. All fungicides have a negative effect on the development of Late blight by inducing lower incidences and severities than those observed in




Fig. 4. Area under the disease progression curve of different tomato plant treated by Mancozeb




Fig. 5. Area under the disease progression curve of different tomato plant treated by Plantineb

Table 7. Effects of tomato varieties and chemical fungicides on the yields

Varieties	Treatments			
	Control	Bonsoin	Mancozeb	Plantineb
AVTO1219	17.29 ± 2.45 ^{aC}	40.57 ± 1.67 ^{bA}	37.90 ± 5.81 ^{bcA}	25.67 ± 3.73 ^{cb}
AVTO1311	11.33 ± 3.89 ^{bc}	35.90 ± 2.21 ^{cA}	39.60 ± 3.54 ^{bcA}	23.59 ± 1.78 ^{cb}
CLN1462A	2.37 ± 1.04 ^{cdD}	53.99 ± 0.73 ^{aA}	47.99 ± 2.81 ^{aB}	37.26 ± 0.79 ^{bc}
CLN1464A	3.27 ± 0.62 ^{cC}	50.65 ± 2.05 ^{aA}	46.37 ± 2.42 ^{aAB}	45.33 ± 1.67 ^{aB}
CLN1464B	1.87 ± 0.42 ^{dC}	35.80 ± 1.84 ^{cA}	38.26 ± 1.75 ^{cA}	29.37 ± 1.39 ^{cb}
RIOGRANDE+	1.83 ± 0.07 ^{dD}	38.96 ± 1.93 ^{cAB}	41.99 ± 0.87 ^{bA}	35.76 ± 0.58 ^{bb}
RIOGRANDE2	2.66 ± 1.03 ^{cdD}	35.70 ± 3.67 ^{cA}	30.56 ± 2.45 ^{dB}	27.57 ± 1.06 ^{cc}

a, b, c: means assigned the same lower case letter in the same column; as well as A, B, C: means assigned the same upper case letter in the same row are not significantly different according to Student Newman-Keuls test at 5%

control plants. these results could be due to the fact that the fungicides used were contain of active substances that may have inhibited the development of *Phytophthora infestans* the causal agent of the disease. These results are similar to those of the works of Randriantsalama et al. [14], who showed that some fungicides such as Mancozeb, Bonsoin, and Plantneb have an inhibitory effect on the development of *P. infestans*, causal agent of Late Blight in Solanaceae. The finding of this study is in line with work of Getachew [6], reported that 68.85% infestation from protected plot and 90.97% infestation from unprotected plot. Similarly, Ashenafi et al. [8] research result, the maximum (91.5%) disease incidence was showed from the unsprayed control of susceptible potato variety (Jalene). Of all the fungicides, Bonsoin was the most effective in the management of the disease the disease. In line with Hagos et al. [4], who found that frequently applied fungicides by far reduced disease severity as compared to the less frequently sprayed fungicides and unsprayed plots of tomato. This result could be explained by the fact that not all fungicides used in this study had the same active substance and may have behaved differently towards *P. infestans*. In addition, Bonsoin could contain an active substance to which *P. infestans* would have be more sensitive to, compared to the other fungicides [8].

From one phase of development to another, the incidence and severity of the disease increased. Thus, the highest incidence and severity was obtained in the fruiting phase. This could be due to the contamination of new plants over time. These results corroborate those of Kesksse et al. [1], who showed that as soon as blight appeared in the first tomato plants, the disease spread through contamination of the other plants. Incidence and severity of Late blight varied according to the tomato variety. Varieties

AVTO1219 and AVTO1311 showed the lowest incidences and severities. While RIO GRANDE+ and RIO GRANDE 2 had the highest. These results could be related to the fact that the varieties AVTO1219 and AVTO1311 are less sensitive to Late Blight, caused by *P. infestans*. Hence, the low incidences and severities observed in these two tomato varieties AVTO1219 and AVTO1311. While local variety RIO GRANDE+ were more sensitive to the disease and had the highest incidence and severity in the fruiting phase. This difference of results could be explained by the fact that the AVTO1219 and AVTO1311 varieties have a genetic material that permits them to be less sensitive to the disease. This would not be the case for the RIO GRANDE+ (local variety). Ghislain et al. [27], showed that Late blight was strongly present on all the local tomato varieties they grew. This difference in results could be explained by the fact that none of the potato varieties used were less sensitive to *P. infestans*. However, some of the tomato varieties used in this study were reported to be less sensitive to *P. infestans*. According to Forbes et al. [28] and Nowakowska et al. [29], there are tomato varieties that are genetically resistant to *P. infestans*, causal agent of Late Blight in Solanaceae.

The result of AUDPC showed that, the chemical fungicide Bonsoin considerably reduced the AUDPC of all the varieties studied. This would be due to the dual contact and systemic action of this fungicide which would have inhibited the development of *P. infestans*. These results corroborate with those of Getachew (2017), Daniel et al. [30] and Arafa, et al. [31] working on tomato and potato and reported that, low AUDPC value was obtained from protected tomatoes varieties with four times sprayed Mancozed and high value from unprotected plot of tomatoes varieties.

The yields obtained with the plants treated with the different fungicides were higher than those of the control plants. This could be explained by the fact that the fungicides applied to the plants of the different tomato varieties would have significantly reduced the incidence of fungal diseases. Hence the high yields obtained. These results corroborate with de Getachew [6] and Daniel et al. [30] who reported that fungicide-treated tomato plants produced higher yields than the control plants. The varieties AVTO1219 and AVTO1113, even without receiving fungicide treatments, showed higher yields compared to the other control varieties. These results would be due to the fact that these two tomato varieties are less susceptible to Late Blight. This is because they showed the lowest prevalence's and severities, compared to the other varieties where these two parameters were high. As a result, the disease caused less damage to them; hence the high yields they produced. The results are similar to those of Forbes et al. [28] and Nowakowska et al. [29], who showed that Late Blight-resistant tomato varieties produced higher yields than those not resistant to the disease.

5. CONCLUSION

This study showed that some chemical fungicides such as Bonsoin and tomato varieties AVTO1219 and AVTO1311 could be used to manage the Late Blight of tomato cause by *Phytophthora infestans*. However, RIO GRANDE 2 and RIO GRANDE+, were highly attacked by the disease. Infact when these two tomato varieties were treated with the chemical fungicide, these chemical fungicides help to reduce the incidence and severity of the Late Blight. Genetic control by using less susceptible varieties to late blight is the best way to manage tomato disease. The combination of these varieties with the most effective fungicide would help to increase yields in the region.

ACKNOWLEDGEMENTS

The authors would like to thank the World Vegetable Center, Yaoundé for providing us the tomato seed varieties needed for this work. The staff of the Research Unit and Applied Botany of the University of Dschang for their accompaniment in the realization of this work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Keskse D, Hailu N, Belete E. Integrated Management of Tomato Late Blight (*Phytophthora infestans* Mont.) of Tomato (*Lycopersicon esculentum* MILL.) at Ataye and Shewarobit Districts, Eastern Amhara, Ethiopia. *Journal of Plant Sciences*. 2019; 7(2):63-71.
2. Farheen J, Mansoor S, Abideen Z. Application of salicylic acid improved growth, photosynthesis pigments and oxidative stability in mungbean seedlings at salt stress. *Pak. J. Bot.* 2018;50:901-912.
3. Micah AE, Chen CS, Zlavog BS, Hashimi G, Chapin A, Dieleman JL. Trends and drivers of government health spending in sub-Saharan Africa, 1995–2015. *BMJ Glob Health*. 2018;4:1-10. DOI:10.1136/bmjgh-2018-001159
4. Hagos G, Meles K, Tsehaye H. Effect of varieties and fungicide application frequencies on Late Blight (*Phytophthora infestans*) Disease Development and Fruit Yield of Tomato in North Western Tigray, Ethiopia. *Journal of Plant Pathology and Microbiology*. 2020;11(12): 1-6.
5. Ephrem G. Disease management practice on potato (*Solanum tuberosum* L.) in Ethiopia. *World J Agric Res*. 2015;3(1):34–42. Available:<https://doi.org/10.12691/wjar-3-1-8>
6. Getachew G. Integrated management of tomato late blight (*Phytophthora infestans* (Mont.) de Bary) through host plant resistance and reduced frequency of fungicide application in gamogofa zone, southern Ethiopia. MSc. Thesis College of Agriculture, Haremya University, Haremya, Ethiopia. 2017;59.
7. Bastin LA. Étude de l'interaction entre *Phytophthora infestans* et *Solanum tuberosum*. Faculté des bioingénieurs, Université catholique de Louvain. Prom.: Legrève, Anne ; Colau, Gil ; 2020. Available:<http://hdl.handle.net/2078.1/thesis:27158>
8. Ashenafi MY, Thangavel S, Alemu L, Bekele K. Evaluation of potato cultivars and fungicides for the management of Late blight (*Phytophthora infestans* (Mont) de Bary) in Holleta, West Showa, Ethiopia. *International Journal of Life Sciences*. 2017;5(2):161-179.

9. Mideros MF, Turissini DA, Guayazán N, Ibarra-Avila H, Danies G, Cárdenas M, et al. *Phytophthora betacei*, a new species within *Phytophthora* clade 1c causing late blight on *Solanum betaceum* in Colombia. *Persoonia-Molecular Phylogeny and Evolution of Fungi*. 2018;41(1):39-55.
10. Kassaw A, Abera M, Belete E. The Response of Potato Late Blight to Potato varieties and Fungicide Spraying Frequencies at Meket, Ethiopia. *Cogent Food and Agriculture*. 2021;7(4):1-17.
11. Goufo P, Fontem DA, Ngnokam D. Evaluation of plant extracts for tomato Late blight control in Cameroon. *New Zealand Journal of Crop and Horticultural Science*. 2010;38(2):171-176.
12. Fontem DA, Olanya MO, Tsopmbeng GR, Owona M. Pathogenicity and metalaxyl sensitivity of *Phytophthora infestans* isolates obtained from garden huckleberry, potato and tomato in Cameroon. *Crop Protection*. 2005;24:449-456.
13. Aghofack-Nguemezi J, Tsopmbeng NG, Ndille NC. Influence of calcium and magnesium based fertilizers on fungal diseases, plant growth parameters and fruit quality of three varieties of tomato (*Solanum lycopersicum*). *Journal of Science and Technology*. 2014;34(1):9-20.
Available:<http://dx.doi.org/10.4314/just.v34i1.2>
14. Randriantsalama AR, Randrianaivoarivony JM, Ramalanjaona VL. L'utilisation de la lutte chimique et de la résistance variétale contre le mildiou de la pomme de terre à Madagascar. *African Crop Sciences Journal*. 2014;22(2):959-968.
15. Peerzada SH, Mushtaq A, Dar GH, Chattoo MA, Bhat KA. Evaluation of cultivars of potato (*Solanum tuberosum*) for disease reaction against Late blight pathogen (*Phytophthora infestans* (Mont) de Bary). *International Journal of Current Microbiology and Applied Science*. 2013;2(9):153-159.
16. Savazzini F, Galletti S. Phenotypic and genotypic characterisation of Italian *Phytophthora infestans* isoLate. *Phytopathologia Mediterranea*. 2015; 54(4):524-530.
17. Alharby HF, Metwali EMR, Fuller MP, Aldhebiani AY. The alteration of mRNA expression of SOD and GPX genes, and proteins in tomato (*Lycopersicon esculentum* Mill) under stress of NaCl and/or ZnO nanoparticles. *Saudi J. Biol. Sci*. 2016;23:773-781.
18. Njoroge AW, Andersson B, Lees AK, Mutai C, Forbes GA, Yuen JE, Pelle R. Genotyping of *Phytophthora infestans* in eastern Africa reveals a dominating invasive European lineage. *Phytopathology*. 2019;109(4):670-680.
19. Aghofack-Nguemezi J, Passannet AS, Tatchago V. Effets des extraits ou de la poudre de *Spirulina platensis* et *Jatropha curcas* sur la croissance et le développement de la tomate. *Journal of Applied Biosciences*. 2015;90(2):8413-8420.
20. PNDP (Programme National de Développement Participatif). Plan communal de développement de Bangangté. 2015;34.
21. CTFC (Centre Technique de la Forêt Communale). Rapport d'enquête socio-economique des villages riverains à la réserve forestière ; 2013.
Available:www.foretcommunale-cameroun.org
22. Fondio L, Djidji A, N'gbesso M, Kone D. Evaluation de neuf variétés de tomate (*Solanum lycopersicum* L.) par rapport au flétrissement bactérien et à la productivité dans le Sud de la Côte d'Ivoire. *International Journal of Biological and Chemical Sciences*. 2013;7(3):1078-1086.
23. Mekonen S, Tadesse T. Effect of Varieties and Fungicides on Potato Late Blight (*Phytophthora infestans*, (Mont.) de Bary) Management. *Agrotechnology*. 2018;7(2):1-4.
24. Camara M, Mbaye A, Samba N, Gueye T, Noba K, Cilas C. Etude de la productivité et de la sensibilité de diverses variétés de tomate (*Lycopersicon esculentum* Mill.) à la virose du jaunissement et de l'enroulement en cuillère des feuilles au Sénégal. *International Journal of Biological and Chemical Sciences*. 2013;7(6):2504-2512.
25. De Putter H, Schepers H, Maerere A, Tilya MS, Mamiro DP, Kejo D, et al. Late blight (*Phytophthora infestans*) control in Tomato (*Solanum lycopersicon*) in Tanzania. *Proceeding of 7th Euro Blight Workshop*; May 12–15; York, UK. 2019;115-126.
26. Campbell CL, Madden LV. *Introduction to plant disease epidemiology*. John Wiley and sons, New York, USA ; 1990.
27. Ghislain M, Byarugaba AA, Magembe E, Njoroge A, Rivera C, Román ML, et al.

- Stacking three late blight resistance genes from wild species directly into African highland potato varieties confers complete field resistance to local blight races. *Plant Biotechnology Journal*. 2018;17(3):1119-1129.
28. Forbes G, Pérez W, Andrade PJ. Field assessment of resistance in potato to *Phytophthora infestans*. International Potato Center (CIP), Lima (Peru). 2014 ;1-35.
29. Nowakowska M, Nowicki M, Kłosinska U, Maciorowski R, Kozik E. Appraisal of artificial screening techniques of tomato to accurately reflect field performance of the Late blight resistance. *PLoS ONE*. 2014;9(10):5-13.
30. Daniel K, Negash H, Eshetu B. Integrated Management of Tomato Late Blight (*Phytophthora infestans* (Mont.) of Tomato (*Lycopersicon esculentum* MILL.) at Ataye and Shewarobit Districts, Eastern Amhara, Ethiopia. *Journal of Plant Sciences*. 2019;7(3):63-71.
31. Arafa RA, Kamel SM, Taher DI, Solberg S, Rakha MT. Leaf Extracts from Resistant Wild Tomato Can Be Used to Control Late Blight (*Phytophthora infestans*) in the Cultivated Tomato. *Plants*. 2022;11(2): 1824-1834.

© 2023 Dietchou et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/102039>