



On the 2-Rainbow Domination Number Over the γ -set of Some Classes of Graphs

Isagani S. Cabahug, Jr. ^{a*}

^aDepartment of Mathematics, College of Arts and Sciences, Central Mindanao University, Musuan, Maramag, Bukidnon, Philippines.

Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/ARJOM/2023/v19i9712

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102674>

Received: 13/05/2023

Accepted: 15/07/2023

Published: 21/07/2023

Original Research Article

Abstract

A 2-rainbow domination function (2-RDF) is a function $f : V(G) \rightarrow 2^{[2]}$ on a vertex set $V(G)$ of a graph G , such that for any vertex v with $f(v) = \emptyset$, we have $\bigcup_{u \in N(v)} f(u) = [2]$. The weight of f is $\|f\| = \sum_{v \in V(G)} |f(v)|$. The minimum value of $\|f\|$ is the 2-rainbow domination number denoted by $\gamma_{r2}(G)$. In this study, we determine the 2-rainbow domination number of some special classes of graphs over its corresponding γ -set, specifically, path, cycle, fan, and wheel graphs.

Keywords: Domination; rainbow dominating function; rainbow domination number.

2020 Mathematics Subject Classification: 05C35.

*Corresponding author: E-mail: isaganicabahugjr@cmu.edu.ph;

Asian Res. J. Math., vol. 19, no. 9, pp. 212-217, 2023

1 Introduction

One of the enthralling branches in graph theory is the domination in graphs that is presently effectively in excess of several investigations that have been referred to and presented since this has published in [1]. Several authors have defined and analyzed various varieties of domination. One of such is the concept of rainbow domination by Bresar et al [2]. It determines the k -rainbow domination number of a graph G denoted by $\gamma_{rk}(G)$, which is the minimum weight of a k -rainbow dominating function [3]-[5]. The k -rainbow dominating function (k RDF) is a mapping from $V(G)$ to the power set of $\{1, 2, \dots, k\}$. Prior to the release of the concept of k -rainbow domination, Bresar and Sumenjak in [6], introduced the 2-rainbow domination in graphs for which exact values of 2-rainbow domination number of several classes of graphs can be found, and it is shown that for the generalized Petersen graphs $GP(n, k)$ this number is between $\ast \frac{4n}{5}$ and n . Further study on the 2-rainbow domination in the Petersen graph is also investigated in [7]. In this study we investigate the concept of 2-rainbow domination over a γ -set of some classes of graphs.

2 Preliminary Notes

This section contains some of the fundamental concepts necessary for the understanding of the study. Related definitions that are not in this study can be found in [8], [9], [10], [11], [6], and [2].

Definition 2.1. [1] If S and T are any two nonempty subsets of $V(G)$, we say S dominates T if $T \subseteq N[S]$. If S dominates $V(G)$ then S is a dominating set of G . The domination number of G , denoted by $\gamma(G)$, is the minimum cardinality of dominating set of G . A set whose cardinality is the domination number of a graph G is called γ -set of G .

Definition 2.2. [2] Let $[k] = \{1, 2, 3, \dots, k\}$, where the elements of $[k]$ are called colors. Given a graph G and a positive integer k , a k -rainbow domination function (or k -RDF) is function $f : V(G) \rightarrow 2^{[k]}$, such that for any vertex v with $f(v) = \emptyset$, we have $\bigcup_{u \in N(v)} f(u) = [k]$. Let $\|f\| = \sum_{v \in V(G)} |f(v)|$; we refer to $\|f\|$ as the weight of f . The k -rainbow domination number, $\gamma_{rk}(G)$, of graph G , is the minimum value of $\|f\|$.

3 Main Results

This section deals about the 2-rainbow dominating number of special classes of graphs over γ -set of some classes of graphs, specifically, path, cycle, fan, and wheel.

Theorem 3.1. *If P_n is a path graph of order $n \geq 3$, then*

$$\gamma_{r2}(P_n) = \begin{cases} \frac{2n}{3}, & \text{if } n \equiv 0 \pmod{3} \\ \frac{2n-2}{3}, & \text{if } n \equiv 1 \pmod{3} \\ \frac{2n-1}{3}, & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

Proof: Let $P_n = \{v_1, v_2, \dots, v_n\}$ with $n \geq 3$ and let $\emptyset \neq S \subseteq V(P_n)$ be a γ -set of P_n . Consider the following cases:

Case 1: $n \equiv 0 \pmod{3}$

Let $V(G) = S \cup S_1 \cup S_2$ such that $S = \{v_2, v_5, \dots, v_{n-2}\}$, $S_1 = \{v_1, v_4, \dots, v_{n-3}\}$ and $S_2 = \{v_3, v_6, \dots, v_n\}$. Clearly, $|S_1| = |S_2| = \frac{n}{3}$. Define a function $f : V(P_n) \rightarrow 2^{[2]}$ by

$$f(v) = \begin{cases} \emptyset, & \text{if } v \in S \\ \{1\}, & \text{if } v \in S_1 \\ \{2\}, & \text{if } v \in S_2 \end{cases}$$

Then f is a 2-RDF of G with

$$\|f\| = \sum_{v \in V(P_n)} |f(v)| = |S| + |S_1| + |S_2| = 0 + \frac{n}{3} + \frac{n}{3} + \frac{2n}{3}.$$

Now, to show that $\|f\|$ is the minimum weight of P_n , suppose $\|f\|$ is not the minimum weight of P_n . Then there exists a function $g : V(P_n) \rightarrow 2^{[2]}$ such that $\|g\| < \|f\|$. Since S is a γ -set of P_n , then $g(v) = \emptyset$ for each $v \in S$. Now, since $\|g\| < \|f\|$, then there exist $u, w \in V(G) \setminus S$ such that $g(u) = \{1\} = g(w)$, a contradiction. Thus, $\|f\| = \frac{2n}{3}$ is the minimum weight of P_n . Hence, $\gamma_{r2}(P_n) = \frac{2n}{3}$.

Case 2: $n \equiv 1 \pmod{3}$

With the same assumption as case 1. Let $S = \{v_1, v_4, \dots, v_n\}$, $S_1 = \{v_2, v_5, \dots, v_{n-2}\}$ and $S_2 = \{v_3, v_6, \dots, v_{n-1}\}$. Clearly, $|S_1| = \frac{n-1}{3}$ and $|S_2| = \frac{n-1}{3}$. Define a function $f : V(P_n) \rightarrow 2^{[2]}$ by

$$f(v) = \begin{cases} \emptyset, & \text{if } v \in S \\ \{1\}, & \text{if } v \in S_1 \\ \{2\}, & \text{if } v \in S_2 \end{cases}$$

Then f is a 2-RDF of G with

$$\|f\| = \sum_{v \in V(P_n)} |f(v)| = |S| + |S_1| + |S_2| = 0 + \frac{n-1}{3} + \frac{n-1}{3} + \frac{2n-2}{3}.$$

To show that $\|f\|$ is the minimum weight of P_n . On the contrary, suppose $\|f\|$ is not the minimum weight of P_n . Then there exists a function $g : V(P_n) \rightarrow 2^{[2]}$ such that $\|g\| < \|f\|$. Since S is a γ -set of P_n , then $g(v) = \emptyset$ for all $v \in S$. Now, since $\|g\| < \|f\|$, then there exist $u, w \in V(G) \setminus S$ such that $g(u) = \{1\} = g(w)$, a contradiction. Thus, $\|f\| = \frac{2n-2}{3}$ is the minimum weight of P_n . Hence, $\gamma_{r2}(P_n) = \frac{2n-2}{3}$.

Case 3: $n \equiv 2 \pmod{3}$

Choose $S = \{v_1, v_2, \dots, v_{n-1}\}$, $S_1 = \{v_2, v_5, \dots, v_n\}$ and $S_2 = \{v_3, v_6, \dots, v_{n-2}\}$. Clearly, $|S_1| = \frac{n+1}{3}$ and $|S_2| = \frac{n-2}{3}$. Define a function $f : V(P_n) \rightarrow 2^{[2]}$ by

$$f(v) = \begin{cases} \emptyset, & \text{if } v \in S \\ \{1\}, & \text{if } v \in S_1 \\ \{2\}, & \text{if } v \in S_2 \end{cases}$$

Then f is a 2-RDF of G with

$$\|f\| = \sum_{v \in V(P_n)} |f(v)| = |S| + |S_1| + |S_2| = 0 + \frac{n+1}{3} + \frac{n-2}{3} + \frac{2n-1}{3}.$$

To show that $\|f\|$ is the minimum weight of P_n . On the contrary, suppose $\|f\|$ is not the minimum weight of P_n . Then there exists a function $g : V(P_n) \rightarrow 2^{[2]}$ such that $\|g\| < \|f\|$. Since S is a γ -set of P_n , then $g(v) = \emptyset$ for all $v \in S$. Now, since $\|g\| < \|f\|$, then there exist $u, w \in V(G) \setminus S$ such that $g(u) = \{1\} = g(w)$, a contradiction. Thus, $\|f\| = \frac{2n-1}{3}$ is the minimum weight of P_n . Hence, $\gamma_{r2}(P_n) = \frac{2n-1}{3}$. \square

Corollary 3.2. *If C_n is a cycle graph of order $n \geq 3$, then*

$$\gamma_{r2}(C) = \gamma_{r2}(P_n) = \begin{cases} \frac{2n}{3}, & \text{if } n \equiv 0 \pmod{3} \\ \frac{2n-2}{3}, & \text{if } n \equiv 1 \pmod{3} \\ \frac{2n-1}{3}, & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

Proof:

This immediately follows from the Theorem 3.1 by constructing the same S, S_1 and S_2 for all cases of n . \square

Theorem 3.3. *Let F_n be a fan graph of order $n \geq 3$. Then $\gamma_{r2}(F_n) = n$.*

Proof: Let $F_n = \{v_1, v_2, \dots, v_n\} \cup \{v_0\}$ where $\deg(v_0) = n$ and let $\emptyset \neq S \subseteq V(G)$ be a γ -set of F_n . Choose $S = \{v_0\}$ and $S_1 = \{v_1, v_2, \dots, v_n\}$. Define the function $f : V(F_n) \rightarrow 2^{[2]}$ by

$$f(v_i) = \begin{cases} \emptyset, & \text{if } i = 0 \\ \{1\}, & \text{if } i \text{ is odd} \\ \{2\}, & \text{if } i \text{ is even} \end{cases}$$

Then f is a 2-RDf of G with

$$\|f\| = \sum_{v \in V(F_n)} |f(v_i)| = |S| + |S_1| + |S_2| = 0 + \frac{n}{2} + \frac{n}{2} = n.$$

Clearly, $\|f\|$ is the minimum weight of F_n . Thus, $\gamma_{r2}(F_n) = n$. \square

Theorem 3.4. *If W_n is a wheel graph with $n \geq 3$, then*

$$\gamma_{r2}(W_n) = \begin{cases} n, & \text{if } n \equiv 0 \pmod{2} \\ n + 1, & \text{if } n \equiv 1 \pmod{2} \end{cases}$$

Proof:

Let $W_n = \{v_1, v_2, \dots, v_n\} \cup \{v_0\}$ with $\deg(v_0) = n$ and let $\emptyset \neq S \subseteq V(G)$ be a γ -set of W_n . Consider the following cases:

Case 1: $n \equiv 0 \pmod{2}$

Choose $S = \{v_0\}$ and $S_1 = \{v_1, v_2, \dots, v_n\}$. Define a function $f : V(W_n) \rightarrow 2^{[2]}$ by

$$f(v_i) = \begin{cases} \emptyset, & \text{if } i = 0 \\ \{1\}, & \text{if } i \text{ is odd} \\ \{2\}, & \text{if } i \text{ is even} \end{cases}$$

Then f is a 2-RDf of G with

$$\|f\| = \sum_{v \in V(W_n)} |f(v_i)| = |S| + |S_1| + |S_2| = 0 + \frac{n}{2} + \frac{n}{2} = n.$$

Clearly, $\|f\|$ is the minimum weight of W_n . Thus, $\gamma_{r2}(W_n) = n$. \square

Case 2: $n \equiv 1 \pmod{2}$

Choose $S = \{v_0\}$ and $S_1 = \{v_1, v_2, \dots, v_n\}$. Define a function $f : V(W_n) \rightarrow 2^{[2]}$ by

$$f(v_i) = \begin{cases} \emptyset, & \text{if } i = 0 \\ \{1\}, & \text{if } i \neq n \text{ and } i \text{ is odd} \\ \{2\}, & \text{if } i \text{ is even} \\ \{1, 2\}, & \text{if } i = n. \end{cases}$$

Then f is a 2-RDf of G with

$$\|f\| = \sum_{v \in V(W_n)} |f(v_i)| = |S| + |S_1| + |S_2| = 0 + \frac{n}{2} + \frac{n}{2} + 1 = n + 1.$$

Clearly, $\|f\|$ is the minimum weight of W_n . Thus, $\gamma_{r2}(W_n) = n + 1$. □

4 Conclusion

In this article, we focus on the concept of 2-rainbow dominating number within specific classes of graphs, namely path, cycle, fan, and wheel graphs, over its γ -set for which exact values are generated. Further research and analysis in this area can contribute to the development of graph theory and its applications in various fields.

Acknowledgment

The author would like to thank the anonymous referees for helpful and valuable comments.

Competing Interests

Author has declared that no competing interests exist.

References

- [1] Haynes TW, Hedetniemi ST, Henning MA. (Eds.). Topics in domination in graphs. Cham: Springer. 2020;64.
- [2] Brešar B, Henning MA, Rall DF. Rainbow domination in graphs. Taiwanese Journal of Mathematics. 2008;12(1):213-225.
- [3] Li X, Shi Y. Rainbow connection in 3-connected graphs. Graphs and Combinatorics. 2013;29(5):1471-1475.
- [4] Chang GJ, Wu J, Zhu X. Rainbow domination on trees. Discrete Applied Mathematics. 2010;158(1):8-12.
- [5] Haynes TW, Hedetniemi ST, Slater PJ. Domination in graphs (Advanced Topics). Marcel Dekker Publications. New York; 1998.
- [6] Bresar B, Sumenjak TK. On the 2-rainbow domination in graphs. Discrete Applied Mathematics. 2007;155(17):2394-2400.
- [7] Xu G. 2-rainbow domination in generalized Petersen graphs $P(n, 3)$. Discrete Applied Mathematics. 2009;157(11):2570-2573.

- [8] Tan KSR, Jr., I. S. C. Safe sets in some graph families. Asian Research Journal of Mathematics. 2022;18(9):1–7.
DOI:<https://doi.org/10.9734/arjom/2022/v18i930399>
- [9] Mangubat, DP, Jr., I. S. C. On the restrained cost effective sets of some special classes of graphs. Asian Research Journal of Mathematics. 2022;18(8):22–34.
DOI:<https://doi.org/10.9734/arjom/2022/v18i830395>
- [10] Dinorog MG, Jr., I. S. C. Rings domination number of some mycielski graphs. Asian Research Journal of Mathematics. 2022;18(12):16–26.
DOI:<https://doi.org/10.9734/arjom/2022/v18i12621>
- [11] Ruaya KKB, Jr., ISC, Eballe RG. Another look of rings domination in ladder graph. Asian Research Journal of Mathematics. 2022;18(12):27–33.
DOI:<https://doi.org/10.9734/arjom/2022/v18i12622>

© 2023 Cabahug, Jr.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/102674>