



## **Estimation of Lung Function Test among Healthy and Blow out Whistle Beginners**

**Padmalochini Sudharsan<sup>a</sup>, R. Gayatri Devi<sup>b†</sup> and A. Jothi Priya<sup>b†</sup>**

<sup>a</sup> Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai- 600077, India.

<sup>b</sup> Department of Physiology, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Science, Saveetha University, Chennai- 600077, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Author PS performed the data verification, manuscript drafting. Author RGD managed the literature search, data collection, analysis and wrote first draft of the manuscript. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i59B34404

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/80607>

**Original Research Article**

**Received 15 November 2021**

**Accepted 17 December 2021**

**Published 18 December 2021**

## **ABSTRACT**

**Introduction:** Pulmonary function test is a non-invasive test. It is measured by the spirometer and it gives an idea about the lung volume, lung capacities, rates of flow and gas exchange. Breathing exercise can increase lung function. The commonly performed breathing exercises are blowing bubbles, blow out whistle, pinwheels, Hoberman spheres etc. The aim of this study is to estimate lung function among healthy and blow out whistle beginners.

**Materials and methods:** 60 Dental students consisting of 2 groups were involved in this study. Group 1: Control individuals (30). Group 2 individuals: blow out whistle beginners (30). Standardised RMS Helios spirometer was used. Age of 18-22, Healthy individuals and non-smokers were included in this study and smokers, alcoholic individuals, respiratory disorder individuals and post- COVID individuals were excluded from the study. Data was collected and analysed using paired 't' tests. Significance is considered at P<0.05 level.

**Results:** In this study FVC, FEV1, FEV3 were found to be significant and FEV1/FVC, FEF 25-75, PEFr were found to be insignificant. The post test values of FVC and FEV1 are comparatively higher than the pre test values. The pre test values of FEF 25-75 and FEV3 are comparatively higher than the post test values.

<sup>†</sup>Assistant Professor;

\*Corresponding author: E-mail: [gayatridevi@saveetha.com](mailto:gayatridevi@saveetha.com);

**Conclusion:** When this blow out exercise is performed there was a significant difference between healthy individuals and blow out whistle beginners. Significant increase in lung function is found. This recreational activity increases lung function which can be a preventive for many pulmonary diseases.

*Keywords: Blow out whistle; lung function; spirometer; breathing exercise; Innovative techniques.*

## 1. INTRODUCTION

Pulmonary function test is a non invasive test which review lung function. It gives an idea about lung function, lung volume, lung capacities, rates of flow and gas exchange. Pulmonary function test is a crucial tool within diagnosis assessment. It helps in management of respiratory diseases in adults as well the older children [1]. The ability of this test to be performed in preschoolers helped to assess first onset of respiratory illness and cystic fibrosis. It also aid in assessment of lung function and permit the evaluation of the therapeutic within the course of pulmonary diseases [2-5].

Previous studies about the investigations of the consequences of breathing exercises have revealed improvements in respiratory muscles and respiratory function that activated trunk control and significantly affected everyday mobility. Determined that raising intra-abdominal pressure during expiration through transverse abdominis exercises had a stable impact on pulmonary function, also on posture control and stabilization. Hodges and Richardson [1999] also reported that when movement began within the distal segments of the upper and lower extremities, the diaphragm increased intra-abdominal pressure, thereby establishing greater trunk stability [6]. Other previous studies stated that there's an in depth correlation between breathing activity and postural control. A number of studies have also been conducted to spot methods of improving respiratory function through typical exercises like diaphragm breathing, and pursed-lip breathing. Exercise using dumbbells or weight training tools has generated unexpected results due to difficulties in accurate motion measurement and therefore the risk of injury [7]. Accordingly, previous studies have emphasized the necessity to research the consequences and therefore the safety of heavy equipment in resistance exercises and claimed the necessity for alternative exercise tools. In contrast, elastic bands are considered suitable equipment for the elderly since it they're relatively safe and straightforward to use while providing similar

physical strength effects to resistance exercises using existing weight training tools. Moreover, elastic bands facilitate muscle use across the whole body and are portable and economical [8]. Although previous studies have reported improvements in ability using elastic bands, none have evaluated the consequences of a rubber band exercise program with breathing techniques on pulmonary function in female seniors. Rubber band exercise with breathing techniques on elderly health have proven improvements in pulmonary function by developing a self-directed exercise program that female seniors [9-12].

Pulmonary function tests (PFTS) are a crucial tool within the investigation and monitoring of patients with respiratory pathology. They supply important information concerning the massive and little airways, the pulmonary parenchyma and therefore the size and integrity of the pulmonary animal tissue. Although they are doing not provide a diagnosis intrinsically, different patterns of abnormalities are seen in various respiratory diseases which helps to determine a diagnosis [13]. We describe the indications for performing PFTS, describe abnormal results and correlate these with underlying pathology [14,15]. Previous studies evident that students practicing sudharshana kriya yoga showed significantly better FVC, FEV1 and PEFR values supporting the beneficial role of yogic breathing exercises in adolescent school children .

Pre-schoolers used to play with this blow whistle, unknowingly this is becoming a pulmonary exercise which becomes prevention for many pulmonary diseases such as asthma, bronchitis etc. The ultimate aim of the study is to estimate lung function tests among healthy individuals and blow out whistle beginners. This research is needed to assess the effect of simple breathing exercise on lung volume capacity.

## 2. MATERIALS AND METHODS

The study setting is simple Random sampling. This study is not an invasive procedure; it is safe and quick for most of the people. This can be

considered as pros of the study. The cons of this study are, this test makes people feel dizzy, nauseating and there is a risk that they may faint. Investigator and principal Investigator are the people involved in this study. The sample taken from the student population, Saveetha Dental College. Sample size is 60 and each group consists of 30 individuals. Group 1: control individuals (30) and Group 2: blow out whistle beginners (30). To minimize sampling bias simple Random sampling is used. Internal validity: standardized spirometer is used and external validity: Instrument is standardized before the measurement taken. Standardised RMS Helios spirometer was used. Age of 18-22, Healthy individuals and non-smokers were included in this study and smokers, alcoholic individuals, respiratory disorder individuals and post- COVID individuals were excluded from the study. Statistical Analysis of the data is performed in SPSS software, paired 't' tests is used.

### 3. RESULTS

The post test values of FVC are comparatively higher than the pre test values, the post test values of FEV1 are comparatively higher than the pre test values, The pre test values of FEF 25-75 are comparatively higher than the post test values, the pre test values of PEFR are comparatively higher than the post test values and the post test values of FEV3 are comparatively higher than the pre test values. The parameters FVC, FEV1 and FEV3 were found to be significant (paired t test, p value 0.009, 0.014 and 0.009). The parameters FEF 25-75 was found to be insignificant (paired t test, p value 0.408). The pre test value of FEV1/FVC is comparatively higher than post test values. FEV1/FVC was statistically insignificant (paired t test, p value: 0.408) [Table1].

From Table 1 we can infer that as the parameters FVC, FEV1 and FEV3 were found to be significant (Paired t test, p values 0.009,

0.014 and 0.009) and the parameters, FEF 25-75, PEFR were found to be insignificant (paired t test, p values 0.408 and 0.425). Student t test p value < 0.05 (significant).

### 4. DISCUSSION

Previous studies showed that FVC, FEV1 were significant and FEV1/FVC was significant which was similar to our findings. They found significant differences in FVC, FEV1, BBS, FRT, 10MWT, TUGT in the experimental group. The study has revealed that the pulmonary functions are improved while undergoing aerobics training. During regular aerobics, it raises cardiorespiratory efficiency. It enhances good pulmonary function. Due to these exercises higher lung volumes, high flow rates are achieved [16]. The aerobic trainees after their training period compared their own values with their previous value obtained before their training period [17]. When FVC was found to be normal, FEV1/FVC was low, which indicates simple obstruction and when FVC is low and FEV1/FVC is normal it indicates non-specific abnormality [13,18]. In the present study, the post test value for FVC, FEV1 and FEF 25-75 are comparatively higher than pre test value, the pre test value for PEFR is comparatively higher than post test value. The parameters FVC, FEV1 and FEV3 were found to be significant with p values 0.009, 0.014 and 0.009 and the parameters, FEF 25-75, PEFR were found to be insignificant with p values 0.408 and 0.425 (significance P<0.05).

The study showed that the FEF 25-75, PEFR were significant and FEV3 was insignificant. In the reference study, a paired sample T test was performed and while comparing PEF levels within each group it neither exhibited significant difference before and after the experiment [9,19,20]. FBE and BBE increase pulmonary function is often improved without utilization of specialist equipment. Which emphasizes the necessity of events of varied training methods like balloons. Its importance in group program

**Table 1. Represents lung function parameters before and after breathing exercise (mean ± SD)**

Parameters	Pre Mean±SD	Post Mean±SD	Student 't' test p value <0.05(significant)
FVC	1.53 ± 0.47549	2.269 ± 0.588	0.009
FEV1	1.503 ± 0.432	2.117 ± 0.541	0.014
FEV1/FVC	98.87 ± 3.573	94.12 ± 9.69	0.140
FEF 25-75	3.669 ± 1.282	3.288 ± 1.003	0.408
PEFR	4.839 ± 1.351	4.458 ± 1.408	0.425
FEV3	1.53 ± 0.475	2.269 ± 0.588	0.009

development including recreational factors, extends and helps in the function of respiratory muscles [21-24]. PEFR value is reduced which indicates Asthma and early airway obstruction. If the value of FEF 25-75 is reduced it can be a small airways obstruction [17,25-28]. It can even occur in asthma patients and the value of a normal PEFR may be an indicator of asthma control [29-33]. The sample size were less and time duration is very less for this study. In future study sample size must be increased and time duration must be extended. This study can be extended in different types of breathing exercises. This study should involve more participants in the future and the duration of this study must be increased.

## 5. CONCLUSION

Within the limits of the study, when this blow out exercise is performed there was a significant difference between healthy individuals and blow out whistle beginners. There was a significant increase in lung function. This recreational activity increases lung function which can be a preventive for many pulmonary diseases such as asthma, COPD (chronic obstructive pulmonary diseases), lung cancer, pneumonia etc.

## SOURCE OF FUNDING

The present project was funded by

- Saveetha Institute of Medical and Technical Sciences
- Saveetha Dental College and Hospital
- Saveetha University
- Arora multispeciality Dental Hospital

## CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

This study is approved by SRB and preserved by author (s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Stocks J. Pulmonary function tests in infants and preschool children \*

- \*pulmonary function tests in infants and preschool children, by Janet Stocks: The contributor retains the copyright to her original artwork. There are no other exceptions to the copyright [Internet]. Kendig's Disorders of the Respiratory Tract in Children. 2006;129-67. Available:<http://dx.doi.org/10.1016/b978-0-7216-3695-5.50013-4>
2. Moya Olivares A, Villarroel Del Pino L, Fierro Tolosa L, Foncea Fierro C, Caussade Larraín S. Spirometric values in healthy preschool children. *Rev Chil Pediatr.* 2019;90(1):69-77.
3. Barabadi H, Mojab F, Vahidi H, Marashi B, Talank N, Hosseini O, et al. Green synthesis, characterization, antibacterial and biofilm inhibitory activity of silver nanoparticles compared to commercial silver nanoparticles [Internet]. *Inorganic Chemistry Communications.* 2021;129:108647. Available:<http://dx.doi.org/10.1016/j.inoche.2021.108647>
4. Gowhari Shabgah A, Ezzatifar F, Aravindhan S, Olegovna Zekiy A, Ahmadi M, Gheibihayat SM, et al. Shedding more light on the role of Midkine in hepatocellular carcinoma: New perspectives on diagnosis and therapy. *IUBMB Life.* 2021;73(4):659-69.
5. Kamath SM, Manjunath Kamath S, Jaison D, Rao SK, Sridhar K, Kasthuri N, et al. In vitro augmentation of chondrogenesis by Epigallocatechin gallate in primary Human chondrocytes - Sustained release model for cartilage regeneration [Internet]. *Journal of Drug Delivery Science and Technology.* 2020;60:101992. Available:<http://dx.doi.org/10.1016/j.jddst.2020.101992>
6. Anaesthetic and respiratory equipment. Spirometers intended for the measurement of time forced expired volumes in humans [Internet]. Available:<http://dx.doi.org/10.3403/30157349u>
7. Hegewald M, Gallo H, Linares O, Jensen R, Morris A. Accuracy of spirometers used in primary care [Internet]. 9.1 Respiratory Function Technologists/Scientists; 2015. Available:<http://dx.doi.org/10.1183/13993003.congress-2015.pa1577>
8. Ali BA, Abro YM, Javed NH, Islam MS. Standardization of Different Spirometers [Internet]. Vol. 53, *Respiration.* 1988;58-63.

- Available:<http://dx.doi.org/10.1159/000195397>
9. Kim K, Han JW, Kim YM. Effects of elastic band resistance exercises with breathing techniques on pulmonary function in female seniors. *J Exerc Rehabil.* 2019 Jun;15(3):419–23.
  10. Clarizia G, Bernardo P. *Diverse Applications of Organic-Inorganic Nanocomposites: Emerging Research and Opportunities: Emerging Research and Opportunities.* IGI Global. 2019;237.
  11. Egbuna C, Mishra AP, Goyal MR. *Preparation of Phytopharmaceuticals for the Management of Disorders: The Development of Nutraceuticals and Traditional Medicine.* Academic Press. 2020;574.
  12. Ezhilarasan D. Critical role of estrogen in the progression of chronic liver diseases. *Hepatobiliary Pancreat Dis Int.* 2020 Oct;19(5):429–34.
  13. R GD, Gayatri DR, Sethu G. Evaluation of adenoids by oronasal and nasal spirometry [Internet]. *Asian Journal of Pharmaceutical and Clinical Research.* 2018;11:272. Available:<http://dx.doi.org/10.22159/ajpcr.2018.v11i10.27365>
  14. Lee EJ, Lee SY, In KH, Yoo SH, Choi EJ, Oh YW, et al. Routine pulmonary function test can estimate the extent of tuberculous destroyed lung. *Scientific World Journal.* 2012 May 1;2012:835031.
  15. Morrison MD. Nasal Spirometry: A Volumetric Study of Nasal Air Flow Capacity [Internet]. *Archives of Otolaryngology - Head and Neck Surgery.* 1969;90:636–40. Available:<http://dx.doi.org/10.1001/archotol.1969.00770030638021>
  16. Nandhini T, Devi G. A comparative study of peak Expiratory Flow Rate in acute and chronic periodontitis. *J Evol Med Dent Sci.* 2020 Nov 2;9(44):3294–9.
  17. Lee D-K, Jeong H-J, Lee J-S. Effect of respiratory exercise on pulmonary function, balance, and gait in patients with chronic stroke. *J Phys Therapy Sci.* 2018 Aug;30(8):984–7.
  18. Dempsey TM, Scanlon PD. Pulmonary Function Tests for the Generalist: A Brief Review [Internet]. *Mayo Clinic Proceedings.* 2018;93:763–71. Available:<http://dx.doi.org/10.1016/j.mayocp.2018.04.009>
  19. Bharath B, Perinbam K, Devanesan S, AISalhi MS, Saravanan M. Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT–29 colon cancer cells [Internet]. *Journal of Molecular Structure.* 2021;1235:130229. Available:<http://dx.doi.org/10.1016/j.molstruc.2021.130229>
  20. J PC, Marimuthu T, C K, Devadoss P, Kumar SM. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study. *Clin Implant Dent Relat Res.* 2018 Aug;20(4):531–4.
  21. Ferryani M, S NTR, Biben V, Arisanti F, Tobing IMS, Pradanasari R. Effects of Incentive Spirometry Breathing Exercise on Lung Function, C-Reactive Protein Level and Lipid Ratio in Individuals with Chronic Spinal Cord Injury [Internet]. *Proceedings of the 11th National Congress and the 18th Annual Scientific Meeting of Indonesian Physical Medicine and Rehabilitation Association;* 2019. Available:<http://dx.doi.org/10.5220/0009088402200226>
  22. Vivekanandhan K, Shanmugam P, Barabadi H, Arumugam V, Daniel Raj Daniel Paul Raj D, Sivasubramanian M, et al. Emerging Therapeutic Approaches to Combat COVID-19: Present Status and Future Perspectives. *Front Mol Biosci.* 2021 Mar 8;8:604447.
  23. Wadhwa R, Paudel KR, Chin LH, Hon CM, Madheswaran T, Gupta G, et al. Anti-inflammatory and anticancer activities of Naringenin-loaded liquid crystalline nanoparticles in vitro [Internet]. *Journal of Food Biochemistry.* 2021;45. Available:<http://dx.doi.org/10.1111/jfbc.13572>
  24. Wahab PUA, Abdul Wahab PU, Madhulaxmi M, Senthilnathan P, Muthusekhar MR, Vohra Y, et al. Scalpel Versus Diathermy in Wound Healing After Mucosal Incisions: A Split-Mouth Study [Internet]. *Journal of Oral and Maxillofacial Surgery.* 2018;76:1160–4. Available:<http://dx.doi.org/10.1016/j.joms.2017.12.020>
  25. Santhakumar P, Roy A, Mohanraj KG, Jayaraman S, Durairaj R. Ethanolic extract of capparid fruit ameliorates methotrexate-induced hepatotoxicity by Activating Nrf2/HO-1 and PPAR $\gamma$  Mediated Pathways [Internet]. *Indian Journal of Pharmaceutical Education and Research.* 2021;55:s265–74.

- Available:<http://dx.doi.org/10.5530/ijper.55.1s.59>
26. Saraswathi I, Saikarthik J, Senthil Kumar K, Srinivasan KM, Ardhanaari M, Gunapriya R. Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: a prospective longitudinal study [Internet]. PeerJ. 2020;8:e10164. Available:<http://dx.doi.org/10.7717/peerj.10164>
  27. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma [Internet]. Journal of Oral Pathology & Medicine. 2019;48:299–306. Available:<http://dx.doi.org/10.1111/jop.12835>
  28. Tahmasebi S, Qasim MT, Krivenkova MV, Zekiy AO, Thangavelu L, Aravindhana S, et al. The effects of oxygen–ozone therapy on regulatory T-cell responses in multiple sclerosis patients [Internet]. Cell Biology International. 2021;45:1498–509. Available:<http://dx.doi.org/10.1002/cbin.11589>
  29. Nambi G, Kamal W, Es S, Joshi S, Trivedi P. Spinal manipulation plus laser therapy versus laser therapy alone in the treatment of chronic non-specific low back pain: a randomized controlled study. Eur J Phys Rehabil Med. 2018 Dec;54(6):880–9.
  30. Mudigonda SK, Murugan S, Velavan K, Thulasiraman S, Krishna Kumar Raja VB. Non-suturing microvascular anastomosis in maxillofacial reconstruction- a comparative study. J Craniomaxillofac Surg. 2020 Jun;48(6):599–606.
  31. Prakash AKS, Devaraj E. Cytotoxic potentials of *S. cumini* methanolic seed kernel extract in human hepatoma HepG2 cells [Internet]. Environmental Toxicology. 2019;34:1313–9. Available:<http://dx.doi.org/10.1002/tox.22832>
  32. Rajakumari R, Volova T, Oluwafemi OS, Rajesh Kumar S, Thomas S, Kalarikkal N. Grape seed extract-soluplus dispersion and its antioxidant activity. Drug Dev Ind Pharm. 2020 Aug;46(8):1219–29.
  33. R H, Hannah R, Ramani P, Ramanathan A, Jancy MR, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology. 2020;130:306–12. Available:<http://dx.doi.org/10.1016/j.o000.2020.06.021>

© 2021 Sudharsan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/80607>