



Population Dynamics of Defoliator Pests of (*Arachis hypogea* L.) in Different Staggered Sowings

N. Priyanka ^{a*}, O. Shaila ^a, M. Anuradha ^a, V. Divya Rani ^a
and M. Rajashekhar ^a

^a Department of Entomology, College of Agriculture, Professor Jayashankar Telangana State Agricultural University (PJTSAU), Rajendranagar, Hyderabad-500030, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113387

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108770>

Original Research Article

Received: 25/08/2023

Accepted: 01/11/2023

Published: 04/11/2023

ABSTRACT

The present study was conducted to assess the incidence of defoliator pests of groundnut in three different staggered sowings. The experiment was laid with three treatments and seven replications which was carried out at RARS (Regional Agricultural Research Station), Palem, Nagarkurnool district, Telangana state which is having hot and dry climate. The data has been correlated with various weather parameters, including Maximum temperature, Maximum temperature, RH-I and RH-II. The variety K-6 was grown in three staggered sowings (D1: 1st Oct; D2: 15th Oct; D3: 1st Nov). There was a less population of leaf miner during D1 the correlation studies revealed that tobacco caterpillar showed a non-significant negative correlation with the Tmax °C ($r = -0.233$) and a significant positive correlation with Tmin °C ($r = 0.370$) and RH- II ($r = 0.640$). whereas gram caterpillar showed a positive significant correlation to Tmax ($r = 0.256$) and Tmin ($r = 0.097$) and a negative significant correlation towards RH- I ($r = -0.186$) and RH- II ($r = -0.501$).

*Corresponding author: E-mail: priyankaneeli89@gmail.com;

Keywords: *Spodoptera litura*; *Helicoverpa armigera*; groundnut; staggered sowing.

1. INTRODUCTION

Groundnut (*Arachis hypogea* L.), an annual legume crop, is a member of the Leguminosae or Fabaceae family of legumes. Like other legumes, it has symbiotic nitrogen-fixing nodules in its roots. Additionally, it is a common oilseed crop in India and other tropical and subtropical countries Kandakoor et al. [1]. The major groundnut-producing countries are China, India, Nigeria, USA, Taiwan, Indonesia, Ghana, Argentina and Brazil. China is the world's largest producer of groundnut, followed by India (6.70 M t). In India, approximately 4.76 l ha were planted in 2021, with Karnataka leading the way with 1.32 l ha, followed by Telangana with 1.32 l ha (0.87 lakh ha) www. Agricoop.gov.in [2]. Insect pests play an important role in reducing the yield and productivity of groundnut crop where 52 species of insects affect the groundnut crop. Singh et al. [3]. The altered climatic circumstances enhance thrips and defoliator activity, which peaks during the flowering and pod-formation stages Naresh et al. [4] and results in significant crop losses of 24 to 92%, 16 to 42%, 17 to 40%, and 9 to 22%, respectively Amin, [5]. The larval stage of *Spodoptera litura* Devaki et al. [6] is the only damaging stage, which at first aggressively scrapes the leaf tissue, feeds in a cluster, and quickly skeletonizes the leaves. The larvae of *Helicoverpa armigera* feed on foliage, flowers and leaf buds and make symmetrical holes or cuttings on the leaves Bajja et al. [7] Farmers are continuously spraying a few chemicals with high doses which may lead to resistance. The population dynamics and yield losses of pests are significantly influenced by weather parameters Naresh et al. [8]. So, the present study was formulated to know the incidence of defoliator pests at different dates of sowings. And also, the study emphasizes the incidence of different weather parameters on the pest population.

2. MATERIALS AND METHODS

The present work was carried out at Regional Agricultural Research Station Palem, Nagarkurnool district, Telangana state during Rabi, 2021-22. The experimental site is situated at 16 ° 51' N Latitude and 78 ° 25' E Longitude and 478m of average altitude above the mean sea level. Where the experiment was laid in a 5 x 5 m² patch with three treatments and seven replications in the design of Randomized Block

Design (RBD). The experiment was laid with the K-6 groundnut variety in three staggered sowings (D1: 1.10.2021, D2: 15.10.2021, and D3: 1.11.2021) at every 15 days interval with a spacing of 22.5 cm row to row and 10 cm plant to plant. Regular hand weeding was done to keep the crop free of weeds.

The weekly meteorological data on temperature maximum (Tmax °C), temperature minimum (Tmin °C), relative humidity morning (RH- I %), and evening relative humidity (RH- II %) during the respective Standard Meteorological Weeks (SMW) were recorded from a meteorological observatory farm, RARS, Palem during rabi, 2021-22.

3. RESULTS AND DISCUSSION

3.1 Tobacco caterpillar *Spodoptera litura*

The population dynamics on the incidence of *S. litura* revealed that the per cent damage by *S. litura* in D1 was started during the 42nd SMW with 0.5% infestation and it continued up to 51st SMW with 0.2% (Fig 2). There was a peak increase in the population during the 48th SMW with 11.2% damage where in the temperature maximum was 28.9°C, the temperature minimum was 18.7°C, morning relative humidity was 92.3%, and evening relative humidity was 92.7%, respectively. Then the infestation slowly started to decline at 51st SMW with a 0.2% per cent infestation. In case of D2, the population started at the 45th SMW with an infestation of 2.1% and it continued up to the 2nd SMW. The highest infestation (42.6%) was noticed during the 50th SMW where the temperature maximum, temperature minimum, morning relative humidity, and evening relative humidity were found to be 29.7°C, 17.3°C, 82.4%, and 65.3%, respectively. And then it started decline from 2nd SMW with an infestation of 0.4%. *S. litura* in D3 was initially appeared during the 48th SMW with a 2.1% infestation and lasted until the 3rd SMW and it continuously increased and reached the peak of 21.2% infestation at the 51st SMW where the temperature maximum was 29.3°C, temperature minimum was 17.3°C, morning relative humidity was 80.7%, and evening relative humidity was 62.0%, respectively. However, in the 3rd SMW it began to fall with a 0.2% damage (Table 1).

Table. 1 Seasonal incidence of *S. litura* in relation to the weather parameters

| SMW | Temperature (°C) | | Relative Humidity (%) | | % Damage by <i>S. litura</i> | | |
|-----|------------------|------|-----------------------|--------|------------------------------|------|------|
| | Tmax | Tmin | RH- I | RH- II | D1 | D2 | D3 |
| 42 | 32.0 | 21.0 | 91.1 | 90.0 | 0.5 | 00 | 00 |
| 43 | 31.6 | 19.2 | 87.9 | 60.4 | 1.2 | 00 | 00 |
| 44 | 31.2 | 20.2 | 85.1 | 48.0 | 2.0 | 00 | 00 |
| 45 | 29.8 | 19.6 | 87.6 | 69.7 | 3.2 | 2.1 | 00 |
| 46 | 29.1 | 19.1 | 84.1 | 69.4 | 9.1 | 6.7 | 00 |
| 47 | 29.1 | 18.1 | 85.4 | 84.4 | 9.2 | 22.5 | 00 |
| 48 | 28.9 | 18.7 | 92.3 | 92.7 | 11.2 | 28.5 | 2.1 |
| 49 | 29.7 | 18.2 | 87.6 | 79.9 | 8.5 | 39.2 | 5.2 |
| 50 | 29.7 | 17.3 | 82.4 | 65.3 | 2.2 | 42.6 | 6.7 |
| 51 | 29.3 | 17.3 | 80.7 | 62.0 | 0.2 | 23.1 | 21.2 |
| 52 | 28.2 | 15.0 | 69.0 | 42.3 | 00 | 11.4 | 20.2 |
| 1 | 29.1 | 15.1 | 66.1 | 30.9 | 00 | 4.1 | 10.4 |
| 2 | 29.4 | 15.5 | 75.9 | 52.7 | 00 | 0.4 | 3.2 |
| 3 | 29.5 | 16.8 | 76.4 | 70.9 | 00 | 00 | 0.2 |
| 4 | 29.2 | 16.3 | 77.3 | 55.3 | 00 | 00 | 00 |

SMW- Standard Meteorological Week, Tmax- Temperature maximum, Tmin- Temperature minimum, RH- I - Morning relative humidity morning, RH- II - Evening relative humidity




Fig. 1. Incidence of tobacco caterpillar in relation to weather parameters at different dates of sowings

Table. 2 Correlation coefficient of *S. litura* in relation to the weather parameters

| Dates of sowing | Tmax(°C) | Tmin(°C) | RH- I % | RH- II % |
|-----------------|----------------------|----------------------|----------------------|----------------------|
| D1 | -0.233 ^{NS} | 0.370* | 0.572 ^{NS} | 0.640* |
| D2 | -0.323 ^{NS} | -0.053 ^{NS} | 0.245 ^{NS} | 0.376 ^{NS} |
| D3 | -0.461 ^{NS} | -0.553* | -0.533 ^{NS} | -0.430 ^{NS} |

*Significant at 5% level

**Significant at 1% level

NS- Non significant




Fig. 2. Incidence of gram caterpillar in relation to weather parameters at different dates of sowings

Yadav et al. [9] found that the population of *S. litura* population on groundnut started from the 36th SMW with 0.27 larvae/plant and it reached a maximum during the 41st SMW with 1.07 larvae/plant. According to Ahir et al. [10] the incidence of *S. litura* on groundnut reached a peak during the 40th and 41st SMW with a population of 1.20 and 1.40 larvae/plant, respectively. Arpit et al. [11] observed the incidence of tobacco caterpillar from the first week of the august 33rd SMW and it continued up to the October last week 44th SMW on groundnut. Kumar et al. [12] observed the maximum incidence of *S. litura* during the second fortnight of October on soybean. Pazhanisamy et al. [13] who reported that the incidence of *S. litura* was noticed from 36th SMW and achieved a peak during the 41st SMW with a population of 4.54 larvae per meter row and noticed 68.4 % per cent damage to the leaflets followed by 42nd SMW 50.2 % and 40th SMW 49.5 % during 2010. The maximum larval population occurred during the 40th SMW with 5.92 larvae/mrl 72.5% infestation of leaflets followed by 41st (60.4%) and 39th SMW (54.50%) in 2011 *kharif* on groundnut. According to Monobrullah et al. [14] the population of *S. litura* has noticed from 25th SMW with 3.5 larvae/plant and it increased tremendously

to 4.3 larvae/plant during the 43rd SMW on tomato.

The correlation studies revealed during D1 that the *S. litura* larval population showed a non-significant negative correlation with the Tmax °C ($r = -0.233$) and a significant positive correlation with Tmin °C ($r = 0.370$) and RH- II ($r = 0.640$). While a non-significant positive correlation was occurred with RH- I ($r = 0.572$). Whereas in D2 there was a non-significant positive correlation between RH- I ($r = 0.245$) and RH- II ($r = 0.376$) and a non-significant negative correlation at Tmin °C ($r = -0.053$) and Tmax °C ($r = -0.323$). Similarly, the D3 crop showed a negative non-significant correlation with Tmax °C ($r = -0.461$) and RH- II ($r = -0.430$) and a negative significant correlation with RH- I ($r = -0.533$) and Tmin °C ($r = -0.553$) (Table 2).

Nadaf and Kulkarni [15] observed that there was a positive and significant correlation of *S. litura* with the temperature minimum. According to Pazhanisamy et al. [13] reported that there was a negative correlation between the temperature maximum during the *kharif* 2010 and 2011 on *S. litura* of groundnut. Harsih et al. [16] found that a positive association between *S. litura* population and the relative humidity on groundnut.

Table 3. Multiple regression equation of *S. litura* in relation to the weather parameters

| Different dates of sowings | Regression equation | R ² |
|----------------------------|--|----------------|
| D1 | Y= 59.25+(-3.436) Tmax + (0.8394) Tmin + (0.3790) RH-I + (0.0185) RH-II | 44.7 |
| D2 | Y= 7.603 + (-4.865) Tmax + (-7.235) Tmin + (2.438) RH-I +(-0.0266) RH-II | 75.1 |
| D3 | Y= 33.74 + (-1.7492) Tmax + (-0.8314) Tmin + (-0.0658) RH-I + (0.1088) RH-II | 58.7 |

Y: incidence of *S. litura*, Tmax: Temperature maximum, Tmin: Temperature minimum, RH- I : Morning relative humidity morning, RH- II : Evening relative humidity

The data on regression analysis (Table 3) resulted that In the D1 sown crop, the results revealed the weather parameters shown an influence of 44.7 per cent ($R^2 = 44.7$) on the *S. litura* population. At D2 75.1 per cent ($R^2 = 75.1$) influence of temperature maximum, and morning, evening relative humidity was observed. While there was a 58.7 per cent ($R^2 = 58.7\%$) influence of weather parameters on the incidence was noticed in D3 sown crop. Our results are more are less in line with the findings of Pazhanisamy et al. [13] who reported that there was an influence of 71.7 per cent ($R^2=71.7\%$) of weather parameters on incidence of *S. litura*.

3.2 Gram caterpillar *Helicoverpa armigera*

The first appearance of the *H. armigera* occurred during the 43rd SMW, with an average number of 0.2 larvae/plant, and continued until the 52nd SMW during D1 (Fig 2). Then it gradually increased to a maximum of 1.8 larvae/plant at the 47th SMW where the highest temperature was 29.1°C, the temperature minimum was

18.1°C, and the morning relative humidity was found to be 85.4% and in the evening was 84.4%, respectively. However, during the 52nd SMW the infestation was gradually reduced to 0.2 larvae/plant. In D2, the population of *H. armigera* was initially noticed at the 46th SMW with an average of 1.3 larvae/plant, Then it gradually increased and reached the peak on the 49th SMW with 2.9 larvae/plant where the temperature maximum of 29.7°C, temperature minimum of 18.2°C, and morning relative humidity of 87.6% and 79.9% in the evening. Furthermore, at the 3rd SMW, just 0.2 larvae/plants were detected, indicating a reduction. *H. armigera* was observed in the 50th SMW with a 1.1 larvae/plant and the infestation lasted until the 2nd SMW during D3. And then, it started to increase and reached a maximum during the 52nd SMW with (2.2 larvae/plant) where the temperature maximum was 28.2°C, the temperature minimum was 15°C, morning relative humidity 69% and evening 42.3% and then it started to decline at 2nd SMW with 0.2 larvae/plant (Table 4).

Table 4. Seasonal incidence of *H. armigera* in relation to weather parameters

| SMW | Temperature (oC) | | Relative Humidity (%) | | <i>H. armigera</i> larvae/plant | | |
|-----|------------------|------|-----------------------|--------|---------------------------------|-----|-----|
| | Tmax | Tmin | RH- I | RH- II | D1 | D2 | D3 |
| 42 | 32.0 | 21.0 | 91.1 | 90.0 | 00 | 00 | 00 |
| 43 | 31.6 | 19.2 | 87.9 | 60.4 | 0.2 | 00 | 00 |
| 44 | 31.2 | 20.2 | 85.1 | 48 | 0.5 | 00 | 00 |
| 45 | 29.8 | 19.6 | 87.6 | 69.7 | 0.2 | 00 | 00 |
| 46 | 29.1 | 19.1 | 84.1 | 69.4 | 1.1 | 1.3 | 00 |
| 47 | 29.1 | 18.1 | 85.4 | 84.4 | 1.8 | 1.7 | 00 |
| 48 | 28.9 | 18.7 | 92.3 | 92.7 | 1.0 | 2.0 | 00 |
| 49 | 29.7 | 18.2 | 87.6 | 79.9 | 0.7 | 2.9 | 00 |
| 50 | 29.7 | 17.3 | 82.4 | 65.3 | 0.4 | 1.5 | 1.1 |
| 51 | 29.3 | 17.3 | 80.7 | 62.0 | 0.4 | 1.0 | 1.4 |
| 52 | 28.2 | 15.0 | 69.0 | 42.3 | 0.2 | 1.0 | 2.2 |
| 1 | 29.1 | 15.1 | 66.1 | 30.9 | 00 | 0.7 | 1.0 |
| 2 | 29.4 | 15.5 | 75.9 | 52.7 | 00 | 0.4 | 0.2 |
| 3 | 29.5 | 16.8 | 76.4 | 70.9 | 00 | 0.2 | 00 |
| 4 | 29.2 | 16.3 | 77.3 | 55.3 | 00 | 00 | 00 |

SMW- Standard Meteorological Week, Tmax- Temperature maximum, Tmin- Temperature minimum, RH- I - Morning relative humidity morning, RH- II - Evening relative humidity

Table 5. Correlation coefficient of *H. armigera* in relation to the weather parameters

| Dates of sowing | Tmax(°C) | Tmin(°C) | RH- I % | RH- II % |
|-----------------|----------------------|----------------------|---------------------|---------------------|
| D1 | 0.256* | 0.097* | -0.186* | -0.501* |
| D2 | -0.464 ^{NS} | -0.090 ^{NS} | 0.179 ^{NS} | 0.389 ^{NS} |
| D3 | -0.452 ^{NS} | -0.596* | -0.626* | -0.528* |

*Significant at 5% level
 **Significant at 1% level
 NS- Non significant

Table 6. Multiple regression equation of *H. armigera* in relation to the weather parameters

| Different dates of sowings | Regression equation | R2 |
|----------------------------|---|------|
| D1 | Y=8.209+(0.4287) Tmax +(0.1608) Tmin +(0.0252) RH- I +(0.5225) RH- II | 42.6 |
| D2 | Y=11.527+(-0.4884) Tmax +(0.2342) Tmin + (0.0962) RH- I +(0.06673) RH- II | 68.0 |
| D3 | Y=7.6423+(-0.1452) Tmax +(0.2448) Tmin + (0.0232) RH- I + (- 0.9240) RH- II | 45.3 |

Y: incidence of *H. armigera*, Tmax: Temperature maximum, Tmin: Temperature minimum, RH- I : Morning relative humidity morning, RH- II: Evening relative humidity evening.

The results are in line with our current work Pawar et al. [17] who found that the *H. armigera* was more during the 44th and 45th SMW with 0.35 larvae/plant and it continued up to the 1st SMW with 0.15 larvae/plant. Sardar et al. [18] found the incidence of *H. armigera* was observed at 47th SMW with 0.14 larvae/plant on desi chickpea and 0.56 larvae/plant on Kabuli chickpea varieties. Chatar et al. [19] reported that the larval population of *H. armigera* reached a maximum during the 47th to 50th SMW on chickpea.

The correlation analysis in D1 revealed that the pest showed a positive significant correlation to Tmax (r= 0.256) and Tmin (r= 0.097) and a negative significant correlation towards RH- I (r= -0.186) and RH- II (r= -0.501). And in case of D2, the pest build-up was negatively non-significant towards the Tmax (r= -0.464) and Tmin (r= -0.090), and a positive non-significant correlation towards the relative RH- I (r= 0.179) and RH- II (r= 0.389). However, in D3, the pest showed a negative significant correlation with the RH- I (r= - 0.626) and RH- II (r=-0.528) and Tmax. And a negative, non-significant correlation with Tmax with r=-0.452 (Table 5). The present findings are in line with Gadhiya et al. [16] reported that the pest showed a positive response to mean temperature and significant negative correlation with relative humidity. Harish et al. [20,21] found that the incidence of *H. armigera* was significantly positively correlated to the temperature maximum.

The population of *H. armigera* in the D1 crop with 42.6 per cent (R² =42.6). In D2 sown crop, there was a positive association with all the weather parameters except the temperature maximum with the influence of 68 per cent (R²= 68%). While the D3 had a negative association to temperature maximum and evening relative humidity and a positive association to temperature minimum and morning relative humidity. There was an influence of 45.3 per cent (R² =45.3) with weather parameters (Table 6).

4. CONCLUSION

The study highlights the advantage of early sowing of the groundnut crop in the first week of October where the incidence of *S. litura* started in 42nd SMW and it lasted up to 1st week with a peak incidence of 11.2% when compared to other late sowings. On the other hand, the occurrence of *H. armigera* was first reported at the 42nd SMW and persisted until the 52nd SMW, with an average of 1.8/plant larvae. Thus, this aids in the scheduling of groundnut crop defoliator pest management techniques.

ACKNOWLEDGEMENTS

The authors are thankful to the Head of the Department. Department of Entomology, College of Agriculture, Rajendranagar, PJTSAU, Hyderabad and Associate Director of Research, RARS, Palem, PJTSAU for providing necessary facilities and support for conducting experiments.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Kandkoo SB, Khan HK, Gowda GB, Chakravarthy AK, Kumar CT. and Venkataravana P. The incidence and abundance of sucking insect pests on groundnut. *Current Biotica*. 2012;6(3): 342-348.
- Area, Production, Productivity of groundnut Available:www.agricoop.gov.in
- Singh TVK, Singh KM, Singh RN. Groundnut pest complex: III. Incidence of insect pests in relation to agro-climatic conditions as determined by graphical super imposition technique. *Indian Journal of Entomology*. 1990;52(4):686-692.
- Naresh T, Ramakrishna Rao A, Murali Krishna T, Devaki K, Khayum Ahammed S, Sumathi P. Seasonal incidence and effect of abiotic factors on population dynamics of thrips on groundnut (*Arachis hypogaea* L.) during rabi season. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(2):1600-1604.
- Amin PW. Insect Pests of groundnut in India and their management. *Plant protection in field crops: lead papers of the National Seminar on Plant Protection in Field crop*, 29-31 January 1986, CPPTI, Hyderabad; 1987.
- Devaki K, Prathima V, Murlikrishna T, Ramakrishna Rao A, Venkateshwarlu U. and Suma K. Impact of weather factors on the incidence of leaf miner, *Aproaerema modicella* Deventer on *kharif* groundnut in chitoor district of Andhra Pradesh. *Journal of Agrometeorology*. 2013;15(2): 212-216.
- Bajia R, Singh DS, Bairwa B, Padwal K. Major insect pests of groundnut (*Arachis hypogaea* L.), Nature of damage and their ecofriendly management; 2017.
- Naresh T, Ramakrishna Rao A, Murali Krishna T, Devaki K, Khayum Ahammed S, Sumathi P. Seasonal incidence of leaf miner *Aproaerema modicella* on groundnut (*Arachis hypogaea* L.) during *rabi* season. *Journal of Entomology and Zoology Studies*. 2017;5(6):92-96.
- Yadav MK. and Borad PK. Bio-efficacy of different insecticides against sucking pest of summer groundnut. *Indian Journal of Applied Entomology*. 2012;26(1):36-40.
- Ahir KC, Arti S, Rana BS and Dangi NL. Population dynamics of sucking pest in relation to weather parameters in groundnut (*Arachis hypogaea* L.). *Journal of Entomology and Zoological studies*. 2017;5(2):960-963.
- Arpit KM, Archana K, Awasthi AK, Chaure NK. and Shukla RK. Seasonal incidence of tobacco caterpillar (*Spodoptera litura* Fab.) on groundnut (*Arachis hypogaea* L.) and it's correlation with different abiotic factors. *Journal of Pharmacognosy and Phytochemistry*. 2021;10(2):1456-1459.
- Kumar V, Manglik VP, Battacharya AK. Estimation of population of some insect pests of soybean. *Journal of Insect Science*. 1998;11(1):14-18.
- Pazhanisamy M, Senthilkumar M, Sathyaseelan V. Seasonal incidence of leaf-eating caterpillar, *Spodoptera litura* (Fabricius) in groundnut ecosystem During *kharif* season. *Plant Archives*. 2019;19(2): 3351-3354.
- Monobrullah MD, Poonam B, Uma Shankar, Gupta RK, Srivastava K, Hafeez A. Trap catches and seasonal incidence of *Spodoptera litura* on cauliflower and tomato. *Annals of Plant Protection Sciences*. 2007;15(1):73-76.
- Nadaf AM, Kulkarni KA. Seasonal Incidence of the fruit borers, *Helicoverpa armigera* (Hubner) and *Spodoptera litura* Fabricius on chilli in Dharwad. *Karnataka Journal of Agricultural Sciences*. 2006; 19(3):549-552.
- Gadhiya HA, Borad PK, Bhut JB. Effectiveness of synthetic insecticides against *Helicoverpa armigera* (Hubner) and *Spodoptera litura* (Fabricius) infesting groundnut. *The Bioscan*. 2014;9(1):23-26.
- Pawar UA, Mahale AS, Ambhure KG. Seasonal incidence of gram pod borer, *Helicoverpa armigera* (Hub.) and natural enemies in Pigeonpea. *Green Farming*. 2015;6(1):140-143.
- Sardar SR, Bantewad SD, Jayewar NE. Seasonal incidence of *Helicoverpa armigera* influence by Desi and Kabuli Genotype of Chickpea. *International Journal of Current Microbiology and Applied Sciences*. 2018;6:536-541.
- Chatar VP, Raghvani KL, Joshi MD, Ghadge SM, Deshmukh SG, Dalave SK, Joshi MD. Conducted investigation on population dynamics of *H. armigera* on

- chickpea. *Journal of Plant Protection*. 2010;3(1):65-67.
20. Harish KN. Seasonal incidence of insects pests and their biocontrol agents on soybean. *Journal of Agriculture and Veterinary Science*. 2013;2:7-11.
21. Harish G, Nataraja MV, Jasrotia P, Holajjer P, Savaliya SD, Gejera M. 2015. Impact of weather on the occurrence pattern of insects pests on groundnut. *Legume Research*. 2015;38(4): 524-535.

© 2023 Priyanka et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/108770>