



# Influence of Various Organic Manures on the Growth and Yield of Radish at Different Growth Stages (*Raphanus sativus* L.)

M. E. Supriya <sup>a++\*</sup>, T. Bhagyalakshmi <sup>b#</sup>, S. B. Yogananda <sup>ct</sup>,  
G. S. Yogesh <sup>d‡</sup> and Chandrappa <sup>et</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, UAS GKVK, Bengaluru-560065, India.

<sup>b</sup> Department of Soil Science and Agricultural Chemistry, College of Agriculture, V. C. Farm, Mandya-571405, India.

<sup>c</sup> Department of Agronomy, College of Agriculture, V. C. Farm, Mandya-571405, India.

<sup>d</sup> Krishi Vigyan Kendra, Haradanahalli, Chamarajanagara-571127, India.

<sup>e</sup> Department of Horticulture, College of Agriculture, V. C. Farm, Mandya-571405, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i64656>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116715>

Original Research Article

Received: 06/03/2024

Accepted: 09/05/2024

Published: 14/05/2024

<sup>++</sup> Ph.D. Scholar;

<sup>#</sup> Assistant Professor;

<sup>†</sup> Professor and Head;

<sup>‡</sup> Senior Scientist and Head;

\*Corresponding author: E-mail: [supriyanv24021997@gmail.com](mailto:supriyanv24021997@gmail.com);

**Cite as:** Supriya, M. E., Bhagyalakshmi, T., Yogananda, S. B., Yogesh, G. S., & Chandrappa. (2024). Influence of Various Organic Manures on the Growth and Yield of Radish at Different Growth Stages (*Raphanus sativus* L.). *International Journal of Plant & Soil Science*, 36(6), 540–548. <https://doi.org/10.9734/ijpss/2024/v36i64656>

## ABSTRACT

Applying organic manures is crucial for optimal radish growth and yield due to their rich nutrient content, soil-enriching properties and ability to foster beneficial microbial activity, all of which contribute to healthier plants and improved harvests. A field experiment was conducted to evaluate the influence of different liquid organic manures on growth and yield of radish (*Raphanus sativus*) at organic farming block of Zonal Agricultural Research Station, V. C. Farm, Mandya during late Kharif 2020. The experiment was laid in Randomized Complete Block Design with nine treatments and three replications. The experiment consisted of nine treatments of three different levels of nitrogen was applied through FYM and different times of application of Panchagavya and Jeevamrutha. A significant variation was observed among the treatments. The 75% RDN through FYM + two time application of Panchagavya significantly increased the growth and yield attributes at 15, 30, 45 DAS and at harvest viz., plant height (8.39, 18.25, 28.16 and 33.06 cm), number of leaves per plant (6.22, 9.11, 16.04 and 18.43), leaf area index (0.08, 0.46, 0.70 and 1.01), root diameter (5.43cm at harvest), fresh shoot weight (8.53, 51.31, 115.29 and 132.43 g plant<sup>-1</sup>), fresh root weight (2.99, 61.19, 161.08 and 213.08 g plant<sup>-1</sup>), dry shoot weight (1.11, 10.76, 22.01 and 30.49 g plant<sup>-1</sup>), dry root weight (0.90, 13.13, 26.74 and 43.28 g plant<sup>-1</sup>), root yield (38.38 t ha<sup>-1</sup>) and shoot yield (30.26 t ha<sup>-1</sup>). In total, the results suggested that 75 per cent recommended dosage of nitrogen was applied through FYM with two times application of panchagavya is proved to be conducive for radish cultivation.

**Keywords:** Panchagavya; jeevamrutha; growth; yield; radish.

## 1. INTRODUCTION

“Radish (*Raphanus sativus* L.) is one of important root crop of family Brassicaceae. It is a popular root vegetable in both tropical and temperate regions of world. Radish is grown for its tender root which is consumed either cooked or raw. Leaves and pods of some cultivars can be boiled and eaten as a vegetable. Radish root develops from both the primary root and the hypocotyls” [1]. “It's a popular vegetable crop that's grown all across the globe. The major radish growing states in India are West Bengal, Haryana, Bihar, Assam, Punjab, Odisha, Chhattisgarh, Madhya Pradesh, Uttar Pradesh, Karnataka and Maharashtra. In India, it was grown in an area of 205 thousand hectares during 2019-20 with an annual production of 3170 thousand metric tons. Radish is a good source of vitamin- A and C and also rich source of minerals like calcium, potassium and phosphorus. Radish contains glucose as the major sugar and smaller quantities of fructose and sucrose. Pink-skinned radishes are generally richer in ascorbic acid than white-skinned ones” [2]. “It has *diuretic*, refreshing and cooling properties. It is also used for headache, neurological, sleeplessness and chronic diarrhea. The roots are also useful in urinary complaints, piles, liver and gall bladder complaints” [3]. “The leaves of radish are good source for extraction of protein on a commercial scale and radish seeds are potential source of

nondrying fatty oil suitable for soap making illuminating and edible purposes” [3].

“Growth, yield and quality of radish greatly depend on soil, climatic conditions and nutrient management. It is well documented fact that the increasing use of chemical fertilizers to increase vegetable production has been widely recognized but continuous and non-judicious use of inorganic inputs or commercial fertilizers have an adverse effect on soil health, environment, beneficial soil microorganisms, human being and leaving residual toxicity in the food products whereby reduces the quality of root and shoot” [4]. “In organic farming only organic manures or natural inputs like weeds, leaf litter and crop residues *etc.*, available on the farm are used. Thus, it reduces the cost of production against chemical inputs. Farm yard manure being a bulky organic material reduces the soil compaction and improves the aeration in addition to the supply of essential plant nutrients” [5]. “Apart from using conventional farm-based products there is an increasing demand for improvised materials like Panchagavya, Jeevamrutha and other liquid organic manures which mainly enrich the soil with indigenous microorganisms. These liquid formulations also contain macro nutrients, micro nutrients, many vitamins, essential amino acids, growth promoting factors like IAA, GA and thus helps in improving the growth, yield and quality of the crops” [6]. Keeping this in view the present experiment was conducted to study the influence

of various organic manures on the growth and yield of radish.

## 2. MATERIALS AND METHODS

A field experiment was conducted during late *Kharif* 2020 at organic farming block of Zonal Agricultural Research Station (ZARS) V. C. Farm, Mandya, University of Agricultural Sciences, Bangalore. It is located in the Southern Dry Zone of Karnataka (Agro Climatic Zone VI) and Region III at 11° 30' to 13° 05' North latitude and 76° 05' to 77° 45' East longitude with an altitude of 695 meters above mean sea level (MSL). The experiment consisted of nine treatments *viz.*, T<sub>1</sub> (100% RDN through FYM), T<sub>2</sub> (50% RDN through FYM + one time application of Jeevamrutha), T<sub>3</sub> (50% RDN through FYM + two time application of Jeevamrutha), T<sub>4</sub> (75% RDN through FYM + one time application of Jeevamrutha), T<sub>5</sub> (75% RDN through FYM + two time application of Jeevamrutha), T<sub>6</sub> (50% RDN through FYM + one time application of Panchagavya), T<sub>7</sub> (50% RDN through FYM + two time application of Panchagavya), T<sub>8</sub> (75% RDN through FYM + one time application of Panchagavya) and T<sub>9</sub> (75% RDN through FYM + two time application of Panchagavya). Panchagavya and Jeevamrutha were applied to soil one at the time of sowing and second application at 30 DAS at the rate of 500 L ha<sup>-1</sup>. The experiment was laid out in Randomized Block Design with three replications.

Radish variety *Arka Nishanth* developed at Indian Institute of Horticultural Research; Bangalore was used as a test crop. Prior to 15 days of sowing, well decomposed FYM was applied at the rate of 25 t ha<sup>-1</sup> to the respective experimental plots as per the treatments and it was incorporated with the top soil after application. Radish seeds were dibbled 2 cm down the ridges at a distance of 15 cm in the soil. Thinning was done at 15 days after sowing by rotation one seedling per hill. Seeds were sown in rows at 30 × 15 cm spacing. All the intercultural operations and plant protection measures recommended for the successful crop growth were followed and timely irrigation was given to maintain the proper moisture in the field for better growth and development of the plants.

Randomly five plants from each plot were selected to record the data on plant height (cm), number of leaves per plant, leaf length (cm), leaf area (cm<sup>2</sup>), leaf area index, , fresh root weight (g plant<sup>-1</sup>), fresh shoot weight (g plant<sup>-1</sup>), total fresh

weight (g plant<sup>-1</sup>), dry root weight (g plant<sup>-1</sup>), dry shoot weight (g plant<sup>-1</sup>), total dry weight (g plant<sup>-1</sup>), root length (cm), root diameter (cm), root yield (t ha<sup>-1</sup>), shoot yield (t ha<sup>-1</sup>) and total yield (t ha<sup>-1</sup>).

The obtained data were statistically analyzed as described by Gomez and Gomez [7]. The significance of variation among the treatments was observed by applying ANOVA. The level of significance used in "F" was P = 0.05. Critical difference (CD) values were calculated for the P = 0.05 whenever "F" test was found significant.

## 3. RESULTS AND DISCUSSION

### 3.1 Results

#### 3.1.1 Plant height

There was a significant difference in plant height due to application of organic manures. At 15, 30, 45 DAS and at harvest significantly highest plant height (8.39, 18.25, 28.16 and 33.06 cm respectively) was recorded in T<sub>9</sub> (75% RDN through FYM + two times application of Panchagavya) and was on par with the treatments that received 75 per cent RDN through FYM with one- or two-times application of Panchagavya and Jeevamrutha. The studied conducted by Umar et al. [3] and Gyewali et al. [8] also supported the influence of organic manures on plant height in radish.

#### 3.1.2 Number of leaves per plant

T<sub>9</sub> (75% RDN through FYM + two time application of Panchagavya) recorded highest number of leaves per plant (6.22, 9.11, 16.04 and 18.43) at 15, 30, 45 DAS and at harvest which was on par with T<sub>8</sub>, T<sub>5</sub> and T<sub>4</sub> which received 75 per cent RDN through FYM with one or two times application of Panchagavya and Jeevamrutha, respectively (Table 1). The findings of Mani and Anburani, (2010) as well as those of Umar et al. [3] and Gyewali et al. (2020) support the prior results.

#### 3.1.3 Leaf length, Leaf area and leaf area index

T<sub>9</sub> (75% RDN through FYM + two time application of Panchagavya) recorded highest leaf length (17.24, 26.96, 33.90 and 37.23 cm), leaf area (23.91, 136.77, 209.36 and 302.38 cm<sup>2</sup>) and leaf area index (0.08, 0.46, 0.70 and 1.01) over rest of the treatments (Table 2). These results closely correspond with the research conducted by Mani and Anburani, (2010) and Singh et al. [9].

### 3.1.4 Root length and root diameter

Significant difference with respect to root length and root diameter was observed at harvest (Fig 1). Highest root length of 27.91 cm was recorded in T<sub>9</sub> (75% RDN through FYM + two times application of Panchagavya) and was on par with T<sub>5</sub> (25.74 cm) but significant with rest of treatments. Maximum root diameter 5.43 cm was recorded in the treatment T<sub>9</sub> which received 75% RDN through FYM + two times application of Panchagavya and was on par with the treatments which received two times application of Panchagavya and Jeevamrutha *i.e.*, T<sub>5</sub> (5.10 cm), T<sub>7</sub> (4.95 cm) and T<sub>3</sub> (4.88 cm). The promotive effect of organic manures on root length and root diameter had also been observed by Uddain et al. (2010), Kumar et al. [10], Singh et al. [9] and Subedi et al. [11] in radish.

### 3.1.5 Fresh root weight, fresh shoot weight and total fresh weight

T<sub>9</sub> (75% RDN through FYM + two time application of Panchagavya) recorded highest fresh root weight (2.99, 61.19, 161.08 and 213.08 g plant<sup>-1</sup>), fresh shoot weight (8.53, 51.31, 115.29 and 132.43 g plant<sup>-1</sup>) and total fresh weight (11.50, 112.50, 276.37 and 345.51 g plant<sup>-1</sup>) and lowest was recorded in the treatment 50% RDN through FYM + one time application of Jeevamrutha (T<sub>2</sub>) (Table 3). These results were consistent with research by Kanaujia et al. [12], Singh et al. [9] and Gyewali et al. (2020) in radish.

### 3.1.6 Dry root weight, dry shoot weight and total dry weight

Highest dry root weight (0.90, 13.13, 26.74 and 43.28 g plant<sup>-1</sup>), dry shoot weight (1.11, 10.76, 22.01 and 30.49 g plant<sup>-1</sup>) and total dry weight (2.01, 23.89, 48.75 and 73.77 g plant<sup>-1</sup>) was recorded in T<sub>9</sub> (75% RDN through FYM + two time application of Panchagavya) and minimum values were observed with T<sub>2</sub> (50% RDN through FYM + one time application of Jeevamrutha) at all the stages of crop growth (Table 4). These findings corroborate with their results obtained by Bhaktavathsalam and Geetha [13], Sunandarani and Mallareddy [14], Kumar et al. [15], Kanaujia et al. (2010), Singh et al. [9] and Gyewali et al. [8].

### 3.1.7 Root, shoot and total yield of radish

The root, shoot and total yield of radish varied significantly with the application of Panchagavya

and Jeevamrutha at different stages (Table 5). Lowest root and shoot yield of 26.19 and 21.24 t ha<sup>-1</sup>, respectively in T<sub>2</sub> which was significantly increased to 38.38 and 30.26 t ha<sup>-1</sup>, respectively due to application 75% RDN through FYM + two times application of Panchagavya and it was on par with T<sub>5</sub> (36.08 and 27.35 t ha<sup>-1</sup>, respectively) which received (75% RDN through FYM + two times application of Jeevamrutha). The extent of root and shoot yield increase with application of 75% RDN through FYM +two times application of Panchagavya was 38.31 and 31.85 was higher than that recorded with the application of 100% RDN through FYM (T<sub>1</sub>). The positive response of radish to organic manures with respect to root and shoot yield was evidenced by Kanaujia et al. [12], Jadhav et al. [16], Khalid et al. [17], Ziaf et al. [18] and Singh et al. [9] in radish.

## 3.2 Discussion

Increase in growth parameters that could be obtained with application of Panchagavya and Jeevamrutha can be attributed to higher microbial load and growth hormones which might have enhanced the soil biomass thereby sustaining the availability and uptake of applied as well as native soil nutrients which ultimately resulted in better growth and yield (Boraiah et al., 2017). Panchagavya contains kinetin which has a role in enhancing chlorophyll content in plant leaves thus in turn, enhance photosynthetic activity, nitrogen metabolism in plants which ultimately improving the growth parameters [19]. Further it may be also due to solubilisation of nutrient in soil, absorption of nutrients and moisture due to soil application of Jeevamrutha [20]. Similar increase in the growth parameters upon application of organic manures was reported in radish by Uddain et al. [21], Zeid et al. [18], Satish, D, [22], Pathak et al. [23], Mani et al. [24], Subedi et al. (2018), Khatri et al. [25], Gyewali et al. (2020) and Basnet et al. [26].

Increase in root length and root diameter might be due to addition of organic manures helps soil micro-organisms to produce polysaccharides and thus lead to better soil structure which is useful for root growth and development [9]. Decrease in bulk density, better aggregation, increase in porosity and water holding capacity of the soil due to organic manures might have contributed in increasing the root length and root diameter of the plants. Similar results of root length and root diameter were also found in radish by Uddain et al. [21], Kumar et al. [10], Singh et al. [9], Subedi et al. [11], Mani et al. [24] and Gyewali et al. (2020).

Table 1. Effect of liquid organic manures on plant height and number of leaves at different growth stages of radish crop

Treatment	Plant height (cm)				Number of leaves per plant			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T <sub>1</sub>	7.28	15.55	20.09	23.95	4.91	7.82	9.88	12.78
T <sub>2</sub>	6.56	14.75	18.38	21.58	4.27	6.71	8.29	10.36
T <sub>3</sub>	6.66	14.81	23.23	27.68	4.29	6.83	12.88	15.03
T <sub>4</sub>	7.73	16.85	20.43	23.73	5.42	8.20	10.32	11.76
T <sub>5</sub>	7.84	16.91	25.85	30.47	5.51	8.34	14.81	17.27
T <sub>6</sub>	7.07	14.87	19.71	22.45	4.51	7.51	9.63	12.03
T <sub>7</sub>	7.11	14.99	24.69	28.40	4.64	7.66	13.63	15.80
T <sub>8</sub>	8.30	18.07	21.97	24.56	6.17	9.04	12.05	14.32
T <sub>9</sub>	8.39	18.25	28.16	33.06	6.22	9.11	16.04	18.43
<b>S.Em±</b>	<b>0.33</b>	<b>0.71</b>	<b>0.96</b>	<b>1.14</b>	<b>0.51</b>	<b>0.35</b>	<b>0.51</b>	<b>0.62</b>
<b>CD@ 5%</b>	<b>0.98</b>	<b>2.13</b>	<b>2.88</b>	<b>3.42</b>	<b>NS</b>	<b>1.05</b>	<b>1.53</b>	<b>1.85</b>

Table 2. Effect of liquid organic manures on leaf length, leaf area and leaf area index at different growth stages of radish crop

Treatment	Leaf length (cm)				Leaf area (cm <sup>2</sup> )				Leaf area index			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T <sub>1</sub>	13.82	22.02	26.42	30.58	20.65	119.96	145.31	231.40	0.07	0.40	0.48	0.77
T <sub>2</sub>	12.47	20.75	24.91	28.81	19.41	112.93	138.30	225.79	0.06	0.38	0.46	0.75
T <sub>3</sub>	12.56	20.83	26.60	32.06	19.47	113.66	152.50	251.73	0.06	0.38	0.51	0.84
T <sub>4</sub>	15.35	24.57	27.38	30.31	21.63	125.26	156.39	236.63	0.07	0.42	0.52	0.79
T <sub>5</sub>	15.39	24.65	30.03	35.33	21.96	126.62	190.88	290.02	0.07	0.42	0.64	0.97
T <sub>6</sub>	13.17	21.26	26.18	30.07	20.06	116.99	149.03	234.30	0.07	0.39	0.50	0.78
T <sub>7</sub>	13.26	21.33	29.76	32.71	20.32	117.75	166.27	261.91	0.07	0.39	0.55	0.87
T <sub>8</sub>	17.11	26.58	29.04	32.09	23.54	136.14	157.61	257.75	0.08	0.45	0.53	0.86
T <sub>9</sub>	17.24	26.96	33.90	37.23	23.91	136.77	209.36	302.38	0.08	0.46	0.70	1.01
<b>S.Em±</b>	<b>0.63</b>	<b>1.01</b>	<b>1.24</b>	<b>1.42</b>	<b>0.94</b>	<b>5.45</b>	<b>7.01</b>	<b>11.08</b>	<b>0.009</b>	<b>0.02</b>	<b>0.02</b>	<b>0.04</b>
<b>CD@ 5%</b>	<b>1.90</b>	<b>3.04</b>	<b>3.73</b>	<b>4.24</b>	<b>2.82</b>	<b>16.33</b>	<b>21.01</b>	<b>33.22</b>	<b>0.003</b>	<b>0.05</b>	<b>0.07</b>	<b>0.11</b>




Fig. 1. Root length and diameter of radish as influenced by application of liquid organic manures at harvest

Table 3. Effect of liquid organic manures on fresh root weight, fresh shoot weight and total fresh weight at different growth stages of radish crop

Treatment	Fresh root weight (g plant <sup>-1</sup> )				Fresh shoot weight (g plant <sup>-1</sup> )				Total fresh weight (root + shoot) (g plant <sup>-1</sup> )			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T <sub>1</sub>	2.79	52.14	123.03	158.69	6.89	44.31	85.91	102.73	9.68	96.45	208.94	261.42
T <sub>2</sub>	2.62	49.36	114.92	141.58	5.06	41.58	82.96	95.39	7.68	90.94	197.88	236.97
T <sub>3</sub>	2.74	49.53	138.55	169.88	5.11	41.87	95.61	112.57	7.85	91.40	234.16	282.45
T <sub>4</sub>	2.81	57.06	126.78	154.44	7.22	48.19	88.09	103.78	10.03	104.58	214.87	258.22
T <sub>5</sub>	2.88	57.98	150.67	189.33	7.47	49.66	105.52	126.60	10.35	106.64	256.19	315.93
T <sub>6</sub>	2.63	51.20	122.42	157.08	6.37	42.15	84.08	99.14	8.99	93.35	206.50	256.22
T <sub>7</sub>	2.69	51.87	140.53	181.48	6.53	42.74	99.02	116.62	9.22	94.61	239.55	298.10
T <sub>8</sub>	2.91	60.77	133.99	166.32	8.52	50.22	90.95	108.37	11.43	110.99	224.94	274.69
T <sub>9</sub>	2.99	61.19	161.08	213.08	8.53	51.31	115.29	132.43	11.50	112.50	276.37	345.51
<b>S.Em±</b>	<b>0.12</b>	<b>3.17</b>	<b>5.86</b>	<b>7.46</b>	<b>0.33</b>	<b>1.98</b>	<b>4.04</b>	<b>4.97</b>	<b>0.43</b>	<b>4.40</b>	<b>9.97</b>	<b>12.30</b>
<b>CD@ 5%</b>	<b>NS</b>	<b>NS</b>	<b>17.58</b>	<b>22.38</b>	<b>0.98</b>	<b>5.95</b>	<b>12.10</b>	<b>14.91</b>	<b>1.28</b>	<b>13.20</b>	<b>29.90</b>	<b>36.89</b>

**Table 4. Effect of liquid organic manures on dry root weight, dry shoot weight and total dry weight at different growth stages of radish crop**

Treatment	Dry root weight (g plant <sup>-1</sup> )				Dry shoot weight (g plant <sup>-1</sup> )				Total dry weight (root + shoot) (g plant <sup>-1</sup> )			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T <sub>1</sub>	0.75	10.48	19.45	34.67	0.79	8.57	15.78	22.81	1.54	19.05	35.23	57.48
T <sub>2</sub>	0.61	9.63	19.09	33.91	0.66	8.22	14.10	20.32	1.27	17.85	33.19	54.23
T <sub>3</sub>	0.64	9.88	22.17	36.39	0.69	8.33	18.39	24.67	1.33	18.21	40.56	61.06
T <sub>4</sub>	0.83	12.25	20.67	35.18	0.74	9.29	16.80	22.08	1.87	21.54	37.47	57.26
T <sub>5</sub>	0.88	12.86	24.22	41.77	1.05	9.32	20.29	28.49	1.93	22.18	44.51	70.26
T <sub>6</sub>	0.69	10.19	21.07	34.21	0.74	8.39	16.07	22.23	1.43	18.58	37.14	56.44
T <sub>7</sub>	0.73	10.84	23.70	37.47	0.77	8.46	19.65	26.20	1.50	19.30	43.35	63.67
T <sub>8</sub>	0.86	12.74	21.82	35.84	1.06	10.37	17.78	23.73	1.92	23.11	39.60	59.57
T <sub>9</sub>	0.90	13.13	26.74	43.28	1.11	10.76	22.01	30.49	2.01	23.89	48.75	73.77
<b>S.Em ±</b>	<b>0.08</b>	<b>1.04</b>	<b>0.96</b>	<b>1.62</b>	<b>0.24</b>	<b>0.40</b>	<b>0.77</b>	<b>1.07</b>	<b>0.07</b>	<b>0.89</b>	<b>1.73</b>	<b>2.69</b>
<b>CD @ 5%</b>	<b>NS</b>	<b>NS</b>	<b>2.88</b>	<b>4.86</b>	<b>0.71</b>	<b>1.19</b>	<b>2.32</b>	<b>3.22</b>	<b>0.21</b>	<b>2.66</b>	<b>5.19</b>	<b>8.07</b>

**Table 5. Effect of liquid organic manures on root, shoot and total yield of radish crop**

Treatment	Root yield (t ha <sup>-1</sup> )	Shoot yield (t ha <sup>-1</sup> )	Total yield (t ha <sup>-1</sup> )
T <sub>1</sub>	27.75	22.95	50.70
T <sub>2</sub>	26.19	21.24	47.43
T <sub>3</sub>	31.58	25.32	56.90
T <sub>4</sub>	28.71	22.88	51.59
T <sub>5</sub>	36.08	27.35	63.42
T <sub>6</sub>	27.01	22.11	49.12
T <sub>7</sub>	33.16	26.46	59.62
T <sub>8</sub>	29.66	23.73	53.39
T <sub>9</sub>	38.38	30.26	68.64
<b>S.Em±</b>	<b>1.34</b>	<b>1.08</b>	<b>2.42</b>
<b>CD @ 5%</b>	<b>4.02</b>	<b>3.25</b>	<b>7.27</b>

The maximum root and shoot weight could be attributed to the beneficial effect of organics in improving soil texture, aeration, reducing soil compaction and thus enhances the water and nutrient uptake by increasing the permeability of root cell membrane and stimulating the root growth or it may be attributed to solubilization of plant nutrients by addition of Panchagavya, Jeevamrutha and FYM leading to increase uptake of NPK. Fresh and dry weights of radish root and shoot results corroborate with their results obtained by Bhaktavathsalam and Geetha (2004), Sunandarani and Mallareddy [14], Kumar et al. [15], Kanaujia et al. (2010), Singh et al. [9] and Gyewali et al. (2020).

The observed variation in yield of radish with Panchagavya and Jeevamrutha might be attributed to the application of organic liquid manures enhances the availability of native nutrients to the crops and also improves the soil environment, which stimulated proliferous root system subsequent in better absorption of water and nutrients from the soil resulting in higher uptake and yield [14]. The positive response of radish to organic manures with respect to root and shoot yield was evidenced by Kanaujia et al.[12],Jadhav et al. [16], Khalid et al. [17], Ziaf et al. (2015), Singh et al. [9] and Gyewali et al. (2020) in radish [27-30].

#### 4. CONCLUSION

The organic manures had statistically significant variation ( $p < 0.05$ ) in all the growth and yield parameters. The performance of T<sub>9</sub> was found superior in all recorded growth and yield parameters followed by T<sub>5</sub>. Therefore, 75% RDN through FYM + two times application of Panchagavya and 75% RDN through FYM+ two times application of Jeevamrutha may be suggested to the radish growing farmers to get better results. This experiment should be conducted in a different agro - climatic zones for further validation.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Rawat S, Pant SC. Effect of organic manures, inorganic fertilizers and their combinations on yield of radish (*Raphanus*

2. Ghimire S, Adhikari B, Pandey S, Belbase, K, Lamichhane S, Pathak R. Effect of different organic manure on growth and yield of radish in Deukhuri, Dang, Nepal. Acta Sci. Agric. 2020;4:1-5.
3. Umar UM, Ibrahim I, Iro, Obidola SM., 2019, Growth and yield of radish (*Raphanus sativus* L.) as influenced by different levels of kalli organic fertilizer on the jos plateau. Asian J. Res. Crop Sci. 2019;4(4):1-8.
4. Mali DL, Singh V, Sarolia DK, Teli SK, Chittora A, Dhakar R. Effect of organic manures and bio-fertilizers on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese white. Int. J. Chem. Stud. 2020;6(2):1095-1098.
5. Chaudhary M, Narwal RP. Effect of long-term application of farmyard manure on soil micronutrient status, Arch. Agron. Soil Sci. 2005;51(3):351 – 359.
6. Amalraj Eld, Kumar GP, Ahmed MH. S. K, Abdul R, Kishore N. Microbiological analysis of panchagavya, vermicompost and FYM and their effect on plant growth promotion of pigeon pea (*Cajanus cajana* L.) in India. Org. Agric. 2013;3:23–29.
7. Gomez KA, Gomez AA. Statistical procedures for agricultural research. 2<sup>nd</sup> Ed. John Wiley Sons, New York; 1984.
8. Gyewali B, Maharjan B, Rana G, PANDEY, R, Pathak R, Poudel PR. Effect of different organic manures on growth, yield, and quality of radish (*Raphanus sativus*). Saarc J. Agric. 2020;18(2):101-114.
9. Singh KH, Naseeruddin, Rana DK. Effect of organic manures on growth, yield and quality of radish (*Raphanus sativus* L.) cv. Pusa Desi V. Hort. Flora Res. Spectrum. 2016;5(2):129-133.
10. Kumar S, Sutanu M, Kumar S, Singh HD. Efficacy of organic manures on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese White. Int. J. Plant Sci. 2014;9(1):57-60.
11. Subedi A, Srivastava MD, Sharma, Shah, SC. Effect of organic and inorganic nutrient sources on growth, yield and quality of radish (*Raphanus sativus* L.) varieties in Chitwan, Nepal. Saarc J. Agri. 2018;16 (1):61-69.
12. Kanaujia S, Singh SP, Singh VB, Singh A. K. INM for quality production of radish (*Raphanus sativus* L.) cv. Japanese White. Int. J. Recent Adv. Multidisciplinary Top. 2021;2 (2):13-15.



- (*Raphanus sativus* L.) in acid Alfisol. J. Soils Crops. 2010;20(1):1-9.
13. Bhaktavathsalam R, Geetha T. Macronutrient analysis of vermicompost and their effects on the growth of radish plant. Environ. Ecol. 2004;22(4):941-947.
  14. Sunandarani N, Mallareddy K., Effect of different organic manures and inorganic fertilizers on growth, yield and quality of carrot (*Daucus carota* L.). Karnataka J. Agric. Sci., 2007;20(3):686-688.
  15. Kumar M, Kumar S, Rattan P, Sharma J. P, Rai GK. Response of radish to the use of integrated nutrient management practices. Veg. Sci. 2009;36(3):406-408.
  16. Jadhav PB, Patel DJ, Kireeti A, Patil N. B., Dekhane SS, Harad NB., Effect of different levels of vermicompost on growth and yield of radish cv. Local. Int. J. Inf. Res. Rev. 2014;1(2):29-31.
  17. Khalid M, Yadav BK, Yadav MP., Studies on the effect of integrated nutrient management on growth and yield attributes of radish (*Raphanus sativus* L.). Ann. Hort. 2015;8(1):81-83.
  18. Zeid HA, Wafaa HM, Aboul Seoud ,Alhadad WAA., Effect of organic materials and inorganic fertilizers on the growth, mineral composition and soil fertility of radish plants (*Raphane's sativus*) grown in sandy soil. Middle East J. Agric. Res. 2015;4(01):77-87.
  19. Nileema SG, Sreenivasa MN., Influence of liquid organic manures on growth, nutrient content and yield of tomato (*Lycopersicon esculentum* Mill.) in the sterilized soil. Karnataka J. Agric. Sci. 2011;24(2):153-157.
  20. Yogananda SB, Devakumar N, Shruti MK. Ningaraju., Growth and yield of cowpea as influenced by different sources of organic manures, Nat. Symp. Org. Agric. Sust. Food Sec. Challenges Opp. Tamilnadu, India. 2015;113.
  21. Uddain J, Chowdhury S, Rahman MJ. Efficacy of different organic manures on growth and productivity of radish (*Raphanus sativus* L.). Int. J. Agric. Environ. Biotechnol. 2010;3(2):189-193.
  22. Satish D. Effect of different organic manure on growth and yield of Radish (*Raphanus sativus* L.). MSc. Thesis, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior R.A.K College of Agriculture Sehore 466001 (M.P.); 2016.
  23. Pathak M, Tripathy P, Dash SK, Sahu GS, Pattanayak SK., Effect of source of nutrient on growth, yield and quality of radish (*Raphanus sativus* L.) in radish - coriander cropping sequence. Pharma. Innov. J. 2017;6(12):496-499.
  24. Mani AP, Anburani A., Organic nutrient management technique for enhancing growth and physical parameters in radish (*Raphanus sativus* L.). J. Phytol. 2018;10: 40-42.
  25. Khatri KB, Ojha RB, Pande KR, Khanal B R. Effects of different sources of organic manures in growth and yield of radish (*Raphanus sativus* L.). Int. J. Appl. Sci. Biotechnol. 2019;7(1):39-42.
  26. Basnet B, Aryal A, Neupane A, Bishal KC, Rai NH, Adhikari S, Khanal P, Basnet M. Effect of integrated nutrient management on growth and yield of radish. J. Agric. Nat. Res. 2021;4(2):167-174.
  27. Boraiah B, Devakumar N, Shubha S, Palanna KB. Effect of panchagavya, jeevamrutha and cow urine on beneficial microorganisms and yield of capsicum (*Capsicum annuum* L. var. grossum). Int. J. Curr. Microbiol. Appl. Sci. 2017;6(9): 3226-3234.
  28. Hadley P, Swede, Turnip, Radish. In handbook of food science, food technology and nutrition (Macrae R, Robinson RK, MJ. Sadler eds.), Academic Press, London. 1993;4734.
  29. Subramani A, Anuburani A, Gayathiri M. Response of growth parameters of radish (*Raphanus sativus* L.) to various organic nutrients and biostimulants. Asian J. Hort. 2018;5(2):464-466.
  30. Ziaf K, Latif U, Amjad M, Shabir MZ. Combined use of microbial and synthetic amendments can improve radish (*Raphanus sativus*) yield. J. Environ. Agric. Sci. 2016;6:10-15.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/116715>