



Prediction of Lead Molecule Esculetin of *Tridax procumbens* against Zika Virus Envelope Protein

King Solomon Ebenezer ^{a*} and Rekha Manivanan ^a

^a Department of Drug Discovery and Microbiology, Bioscience Research Foundation, Chennai, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jpri/2024/v36i117619>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126154>

Original Research Article

Received: 03/09/2024

Accepted: 05/11/2024

Published: 12/11/2024

ABSTRACT

In the existing field of science, it is necessary for Siddha, Ayurveda therapists, to understand the interactions and to treat the target regions of the human system effectively. One of the most significant nightmares for scientists and medical professionals has been the Zika virus. Since the bite of the *Aedes* mosquito species has infected people worldwide with the Zika virus (Flavivirus). The human mechanism and infection control interactions of the Zika virus are still unclear. In addition, some herbal products may give a pathway to the treat the zika virus. Many herbs aim to regulate various diseases, but the herb *Tridax procumbens* has a key collection of pharmacological values for diseases. Traditional health practitioners have struggled to grasp the precise mechanism of linking existence and how to treat the disease, so the current Ayurinformatics approach and the evolving field are an idea of a modern medical science system. Here we examine the interaction of flavonoids procumbenetin and esculetin of *Tridax procumbens* with zika virus *in silico* docking and molecular simulation and ADME and toxicity profile for the drug candidate. Hereby we conclude

*Corresponding author: E-mail: solu.king@gmail.com;

that these two compounds have the property to suppress the ZIKA virus and are stable and active with ZIKA virus. Molecular docking of Esculetin and Procumbenetin with ZIKV 5JHM, they bind with each other on active site pockets ASP98, ARG99, GLY100, TRP101, GLY102, ASN103, GLY104, CYS105, GLY106, LEU107, PHE108, GLY109.

Keywords: *Tridax procumbens*; zika flavivirus; *In silico* docking; esculetin; ayurinformatics.

1. INTRODUCTION

Tridax procumbens a perennial, annual creeper herb that lives only for a short period and it was globally distributed. Stems are hairy and branched, ascending with the simple opposite petiole leaves. Flower heads were long stalks yellow rounded three toothed ray florets [1]. Several biological active compounds of herbs were isolated and involved in various pharmacological activities. Secondary metabolites have various medicinal activities like dysentery, wound healing, stomachache, malaria, blood pressure and some viruses [2] used in the Indian traditional medicine system [3]. Alkaloids, carotenoids, flavonoids, saponins, and tannins [4] from *Tridax procumbens* played a major role in drug discovery. The Zika virus's status both domestically and internationally has changed significantly in recent years. World health organization (WHO) identified that the Zika has mostly subsided after the severe outbreaks in 2015–2016, which affected more than 80 countries, while transmission is still occasionally active in some areas. There haven't been any reports of local mosquito-borne Zika transmission in the US since 2018, and the cases that are occurring now are either connected to travel or come from other sources, such as sexual transmission.

Nowadays, Zika viruses are a major threat of full disease in India, which persists in the human body for several days to a week, and once the individual was affected, they are protected from other infections such as neurologic complications [5]. So far, no vaccine or drugs to cure ZIKA virus-infected individuals. Zika virus is a mosquito borne flavivirus [6]. The flavivirus has several proteins responsible for the infection, but envelope glycoprotein is responsible for entering the virus for the infection and targets of neutralizing antibodies for other flaviviruses [7]. In the medical system field, medical informatics cannot be given to all emerging areas of Ayurveda because it varies from the modern system's concepts, though the aim was like that to give relief from pain or disease. So, the use of the recent emerging Ayurinformatics

approach will help to bridge a link between the ancient system and the modern system of medicine. The docking tool was used to predict the preferred orientation of one molecule with respect to another molecule when they bound each other for a stable complex with a minimum energy level [8]. Here, the aim of the present study was to investigate the interaction of *Tridax procumbens* flavonoids procumbenetin and esculetin with (ZIKV) Zika flavivirus viral envelope protein using *in silico* docking, molecular simulation and *in silico* toxicological profile prediction for the initial stage of drug development.

2. MATERIALS AND METHODS

Tridax procumbens flavonoids compounds as ligands selected for the study were esculetin and Procumbenetin. The esculetin ligand structure and physicochemical properties were retrieved from Pubchem database CID: 5281416 (Fig. 1). PubChem was a universal public database used to search a broad range of properties [9]. The Procumbenetin chemical structure was designed based on the chemical formula [10] using CHEMSKTECH software. The physical properties were identified, such as molecular formula, weight, composition, molar refractivity, molar volume, parachor, index refraction, surface tension, density, dielectric constant, polarity, and mass.

The 3-Dimensional structure of Zika virus (ZIKV) envelope glycoprotein was retrieved from the protein data bank (PDB) database [11]. The structure of Procumbenetin and Esculetin was represented in Fig. 1 and Fig. 2 respectively. The RCSB PDB operated by US data center for the global PDB archive enables the 3D structure with functional properties. The PDB ID used was 5JHM (Fig. 3).

Molecular docking studies were performed using commercial software suite SYBYL-X 1.3 software. This software expedites molecular modeling/computational chemistry and other discovery works from high throughput screening to late lead optimization. Here we used to predict

the small molecules such as drug candidate interaction and receptor binding. The ligands esculetin and procumbenetin were predicted for the binding orientation against the target Zika virus to estimate the affinity and activity of the molecule.

Discovery studio visualizer 3.1 software suite created by Accelrys to visualize, analyze, and share biological and chemical active substances. The software was used to view 3D graphic sequences, SMILES, Structure binding, editing and analysis functionally. This suite was very

useful for drug discovery. The suite generates receptor-ligand binding interaction plot and patterns between ligand and the protein. The Molecular dynamic and simulation annealing was performed to determine the ZIKV protein and esculetin complex's stability. The simulation was carried out for 150ns to identify the structural transition in envelope protein. The analysis was performed using g_rms, g_rmsf, and g_hbond GROMACS, root mean square fluctuation (RMSF), the root mean square deviation (RMSD), and the number of H-bonds between the ligand and the zika was obtained.




Fig. 1. Procumbenetin




Fig. 2. Esculetin




Fig. 3. 5JHM (ZIKV- Envelope Protein)

The *in silico* toxicological profiles were predicted using online software tools SwissADME and ProTox-II. The SwissADME was used to evaluate Physical chemical properties, Lipophilicity, water solubility, Pharmacokinetics, Drug likeness and medicinal chemistry [12]. ProTox-II for the prediction of toxicities like LD50, Toxicity Class, Organ toxicity, Carcinogenicity, Immunotoxicity, Mutagenicity, Cytotoxicity and Androgen receptor [13].

3. RESULTS AND DISCUSSION

Molecular docking of Esculetin and Procumbenetin showed interaction with ZIKV 5JHM envelope protein using SYBYL Software, which predicted the interactions between each other through some parameters. The total score was expressed as $-\log(K_d)$ and it includes the crash score between ligand atoms that are separated by rotatable bonds (Table 1 & 2).

The binding affinity was represented in which procumbenetin interacted with the residues like LYS-110, ASP-98 and ARG-99. LYS-110 was an essential compound involved in the formation of a double trimmer complex (Figs. 4a & b).

The observed bond length was 2.8, 2.9, 2.9 and 3.5. This shows the hydrogen bonding affinity with some close interaction. It was reported that LYS-110 is a crucial proliferating cell nuclear antigen (PCNA) target sequence for replication and repair foci. LYS-110 was located near the nuclear localization region. The activation of LYS-110 was known to cause mutations in the heterochemical region of PCNA. PCNA was involved in a wide spectrum of cellular functions, including DNA replication, repair, and epigenetic maintenance [14]. The docking result has shown the potential activity of procumbenetin in binding with LYS-110, predicting the inhibiting capacity of procumbenetin on LYS-110. ASP-98 structurally analogous to the proton-abstracting residues found in the zinc-containing enzymes, suggesting a related role for this residue. ASP-98 has been suggested to act as part of a proton shuttling pathway to this residue. ASP-98 is reported to alter nitrite binding. The activation of ASP-98 would result in the formation of mutation at the extracellular boundary of transmembrane helix 2 that reduces gonadotropin-releasing hormone activity. Nitrite was responsible for cell damage, increasing the methemoglobin level and leading to the development of cancer [15]. The docking result has shown the interaction of

procumbenetin with ASP-98. They must be further analyzed to find its potential or inhibiting activity so that the clear role of procumbenetin on ASP-98 can be explained. ARG-99 was a derivate of ARG that has a crucial role in the protein-protein interface. It can enhance hydration, maintain access to water, and contact polar head groups to stabilize the charged form. It has a unique capacity to retain its charge in environments expected to promote the neutral form. It was the right choice when a biochemical function requires the presence of charge in hydrophobic environments [16]. The interaction of procumbenetin with ARG-99 was observed in our study that could be further clarified to find its potential or inhibiting activity. The nitrating binding affinity of procumbenetin with LYS-110, ASP-98 and ARG-99 could be positive or negative, which must be further analyzed. Its interaction could help enhance the potential activity of the residues on interaction or the harmful activity of the residue. The pharmacological activities of procumbenetin include insecticidal, hypotensive, leishmanicidal, wound healing, hair growth-promoting, anti-inflammatory, immunomodulatory and hepatoprotective [17]. Procumbenetin was a very promising species that produced secondary metabolites reported to have a variety of medicinal uses. Its properties also include high blood pressure, anti-anemic, gastrointestinal, anti-diabetic respiratory infections, and anesthetic properties. This species has a long history of traditional use by different communities [18].

The predicted active sites are A/ASP98, A/ARG99, A/GLY100, A/TRP101, A/GLY102, A/ASN103, A/GLY104, A/CYS105, A/GLY106, A/LEU107, A/PHE108, A/GLY109. Our docking research article has discussed the importance of interacting residues with esculetin like GLY 109, ASP-98 and ARG-99. It has other potential active sites like ASP98, GLY100, TRP101, GLY102, ASN103, GLY104, CYS105, GLY106, LEU107 and PHE108. ASP98 was a protein trimmer present at 3 Å sphere [19]. GLY100 has the role in destabilizing the pyrophosphatases globular structure [20]. TRP101 plays a vital role in maintaining a defined phosphoserine aminotransferase microenvironment essentially required for optimal enzymatic activity [21]. A study has confirmed GLY 104 in the interaction, making it stable and emerging as hotspots in the antigen-EDE antibody association related to dengue fever [22]. The role of other residues is not yet clearly known A study on Zika virus

stated that the decoding LYS28, ASN17, PHE24, ARG213, LYS29, GLY81, SER150, SER215, SER56, GLY106, ARG84, TRP87, THR104, ASP146, GLY107, HIS110, ILE147 and GLY148 might be the potential target for zika virus drug [23].

Table 1. Properties of esculetin

Properties	Values	Properties
Molecular weight (g/mol)	178.143	Molecular weight (g/mol)
Hydrogen bond donor count	2	Hydrogen bond donor count
Hydrogen bond acceptor count	4	Hydrogen bond acceptor count
Complexity	248	Complexity
Topological polar surface area	66.8 A	Topological polar surface area

Table 2. Properties of envelope protein

Chain	Residue	Carbon	Nitrogen	Oxygen	Total	Water molecules
A	391	1867	521	575	2988	128
B	390	1860	519	571	2975	145




Fig. 4a. Docking interaction of procumbentin vs Zika




Fig. 4b. Binding pocket of procumbentin vs Zika

The Esculetin showed a total score of 2.8065 and a crash score of -0.7647 than procumbenetin of 1.7476 and -1.1283. This was due to the degree of penetration by the ligand into the protein. A crash score closes to 0 was favorable and a negative number indicates good penetration with the ligands Esculetin molecule. The D score of -42.8704 and -12.8443 chemscore of Esculetin denotes the charge and van der Waals good interactions between the protein and the ligand and hydrogen bonding, metal-ligand interaction, lipophilic contact, rotational entropy, along with an intercept than procumbenetin molecule. Esculetin had the interaction with the residues like GLY 109, ASP-98 and ARG-99 with the following bond length 3.5, 3.1, 3.2 and 3.0 (Fig. 5).

GLY-109 residue may forbid the formation of this bond due to steric hindrance [24]. It can also lead to its replacement by one of eight bulkier residues that could result in collagen disease termed osteogenesis imperfect [25]. Our docking result has shown the binding affinity of esculetin with GLY-109 that might have the potential in supporting the activity of GLY-109. Like procumbenetin, esculetin had also shown the interaction with ASP-98 and ARG-99. Its potential activity must be further analyzed. The bond length represents that the interaction was within the hydrogen bond length. The typical hydrogen bond length was between 2.8 to 3.5 Å, where above 3.0 was the critical configuration barrier. Lower Van der Waals energy denotes the impact of hydrogen bond activity of these compounds during protein or enzyme interaction. Hydrogen bond with the active site of the target macromolecule will result in a high Glide score. The formation of hydrogen bonds in docking results could predict its potential binding affinity that cannot be easily broken by external force [26]. Multiple molecular analysis has stated that the incidence of hydrogen bonds and ranges are key factors affecting the binding interactions of the ligand-receptor interaction [27]. In our docking result, the bond length varies from 3.0, 3.1, 3.2 and 3.5 (Fig. 5a, b & c). The simulated results show the bond length of 3.0, 3.0, 3.3 and 3.6. The bond length of 3.0 was difficult to compare to the bond length of 3.5. This shows the potential binding affinity. As discussed above, the bond could enhance the potential activity of the compound or could suppress the harmful effect of the compound that must be studied further. A recent study on esculetin has shown its anti-tumor efficacy on loading to Polylactide-co-glycolide nano-micelles and nano-encapsulation

against its cytotoxic activity on in vitro model [28]. Esculetin was reported to be an anti-inflammatory drug for treating bowel disease inflammation that was studied on the rat model through colon cytokine analysis [29]. It was also reported to have potential activity against anti-proliferative and cell death with other pharmacological properties like anti-edema, anti-inflammatory and anti-tumor [30].

Stable interaction was found in the ZIKV protein residues ASP 98 and ARG 99. Outcomes showed that esculetin has remained stable at 310 K with residues in ZIKV protein ASP 98 and ARG 99 for 150ns. No modifications are produced between the ZIKV protein and tannic acid in the bond distance and angle. Although the protein undergoes conformation change during annealing, it does not impact the interacted residues between the tannic acid-protein complexes. Also, the polar range did not go beyond 3.9 Å. Thus, the simulation analysis suggests that the structure was stable and not affected. The above Graph of RMSD and Radius of Gyration retained its values where RMSD did not cross 2Å from its initial configuration, and gyration was maintained at 20Å. So, the molecule was stable. Root mean square deviation (RMSD) of atoms ZIKA protein – Esculetin docked complexes at 310K (Fig. 6 a,b & c).

The rise in the binding energy represents the possibility of a new type of binding in the energetically favorable region. The shift in the value leads to a stronger binding of ligands. This binding flexibility was due to the high hydrophilic nature of the ligands and binding site [31]. The docking of a molecule in a protein-binding domain was a robust technique to elucidate the appropriate binding pose between a numbers of predicted compound poses. Molecular recognition of viral protease-ligand complex at the atomic level was also determined by analyzing the bound conformation of the ligands [32]. The compounds were well positioned at the active site, fully occupying the binding pocket. The docking study led to the prediction that the molecules interact with the zika hydrolase binding domine through the residues ASP 98 and ARG 99. Intermolecular H-binding between ligand and protein plays a crucial role in stabilizing protein-ligand compounds [33]. The stability of the hydrogen bond network formed at room temperature was calculated throughout the 150 ns simulation period.




Fig. 5a. Docking interaction of esculetin vs Zika (Annealing result)




Fig. 5b. Docking interaction of esculetin vs Zika (Simulation result)




Fig. 5c. Binding pocket of esculetin vs zika

ADME, which stands for Absorption, Distribution, Metabolism, and Excretion, refers to the pharmacokinetic processes that a drug or substance undergoes within the body. Understanding the ADME profile of a compound

was essential for assessing its potential therapeutic uses, as well as its toxicity. The physico-chemical properties indicated the molecular weight (524.43g/mol); number of heavy atoms (37); number of aromatic heavy

atoms (16); number of rotatable bonds (6); number of hydrogen bond acceptors (14); number of hydrogen donors (8); molar refractivity (123.14). Water solubility showed 3.28e-01mg/ml and soluble in nature. The pharmacokinetics exhibited the results on GI absorption was too low; no BBB permanent; with P-gp substrate and no CYP1A2, CYP2C19, CYP2C9, CYP2D6 and CYP3A4 inhibitors. The drug likeness bioavailability score showed 0.17.

The toxicity profile for the molecules revealed cytotoxicity and immunotoxicity. No hepatotoxicity. Carcinogenicity, mutagenicity, androgen receptor, estrogen receptor, peroxisome proliferator activated receptor gamma, heat shock factor response element, mitochondrial membrane potential. Further in-vivo studies fetch the exact scenario on the toxicity (Fig. 7).

Specific impact on esculletin's absorption was uncertain. However, with specific in-vitro data on

esculetin, was challenging to determine its distribution characteristics accurately. The specific isoforms involved and the resulting metabolites of esculletin remain largely unknown. The ADME properties and toxicity profile of Zika protein esculletin are well-established through computational approaches. Further research was needed to elucidate its ADME behavior and evaluate its toxicity, ensuring a comprehensive understanding of its safety and potential therapeutic applications. Previous literature revealed that some 17 biologically active quinazoline derivatives virtually screened with the Mpro, PLpro, RdRp, and Spike proteins of SARS-CoV-2. ADMET, toxicity, and drug-likeness properties of these compounds [34]. Prior to preclinical and clinical research, ayurinformatics research investigations can be crucial in the creation of novel Ayurvedic medicine moieties. since a well-established method for drug development is a demanding, protracted, and multidisciplinary endeavor [35].




Fig. 6a. Simulation pocket




Fig. 6b. Simulation with esculletin




Fig. 6c. Radius of Gyration and Retained value of RMSD all Ca atoms of the ZIKA protein and - ESCULETIN complex




Fig. 7. Radar chart for toxicity

4. CONCLUSION

In the current study, esculetin showed better scores on the interaction parameters compared to procumbenetin molecule. However, both Procumbenetin and Esculetin bind with zika virus envelope region. The ZIKA envelope docked active region sites with Esculetin were on the region of GLY 109, ASP 98 and ARG 99. Hereby we conclude that these two compounds have the property to suppress the ZIKA virus and are stable and active with ZIKA virus. Molecular docking of Esculetin and Procumbenetin with ZIKV 5JHM, they bind with each other on active site pockets ASP98, ARG99, GLY100, TRP101, GLY102, ASN103, GLY104, CYS105, GLY106, LEU107, PHE108, GLY109. The interaction results are neutral, which means not too good interaction nor not too bad interaction. The molecular dynamics and stimulation studies demonstrated that the complex of esculetin with zika at the residues ASP-98 and ARG 99 (esculetin) was stable in the time of 150 ns. The toxicity profile for the molecules revealed the adverse effects on cytotoxicity and immunotoxicity. No hepatotoxicity. Carcinogenicity, mutagenicity, androgen receptor, estrogen receptor. Further need an in-vitro and in-vivo model to know the exact mechanism.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

The authors are thankful to Bioscience Research Foundation, Chennai for the valuable support for completion of the research work.

COMPETING INTERESTS

Authors have declared that no competing

interests exist.

REFERENCES

1. Pittu V, KR, BB, Beevi S, Devi P, Avanapu SR. *In vitro* anti-cancer activity of aqueous and acetone extracts of *Tridax procumbens* leaf on PC 3 cell lines. Int. J. Pharm. Pharm. Sci. 2011;3:356–358.
2. Das S. Natural therapeutics for urinary tract infections—a review. Future J. Pharm. Sci. 2020;6: 64.
3. Ravishankar B, Shukla V, Indian Systems of Medicine: A Brief Profile. Afr. J. Tradit. Complement. Altern. Med. 2007;4:319–337.
4. Falade T, Ishola IO, Akinleye MO, Oladimeji-Salami JA, Adeyemi OO. Antinociceptive and anti-arthritic effects of aqueous whole plant extract of *Trianthema portulacastrum* in rodents: Possible mechanisms of action. J. Ethnopharmacol. 2019;238:111831.
5. Noorbakhsh F, Abdolmohammadi K, Fatahi Y, Dalili H, Rasoolinejad M, Rezaei F, Salehi-Vaziri M, Shafiei-Jandaghi NZ, Gooshki ES, Zaim M, Nicknam MH. Zika Virus Infection, Basic and Clinical Aspects: A Review Article. Iran. J. Public Health. 2019;48:20–31.
6. Ali A, Wahid B, Rafique S, Idrees M, Advances in research on Zika virus. Asian Pac. J. Trop. Med. 2017;10:321–331.
7. Plourde AR, Bloch EM. A Literature Review of Zika Virus. Emerg. Infect. Dis. 2016; 22:1185–1192.
8. Rath L. Ayurinformatics- The application of bioinformatics in the ayurvedic system of medicine, Proceedings - 9th International Conference on Information Technology, ICIT; 2006.
9. Available:www.ncbi.nlm.nih.gov/pubchem.
10. Mohammed Ali, Earla Ravinder, Ramidi Ramachandran., 2001.A new flavonoid from the aerial parts of *Tridax procumbens*. Fitoterapia. 2001;72,3,313-315
11. Available:www.rcsb.pdb
12. Antoine Daina, Olivier Michielin and Vincent Zoete. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. Sci. Rep. 2017;7:42717.
13. Banerjee P, Eckert OA, Schrey AK, Preissner R. ProTox-II: a webserver for the prediction of toxicity of chemicals. Nucleic Acids Res. 2018;2;46(W1):W257-W263.

14. Kim BJ, Lee H, Lys-110 is essential for targeting PCNA to replication and repair foci, and the K110A mutant activates apoptosis. *Biol. Cell.* 2008;100:675–686.
15. Flanagan C, Rodic V, Konvicka K, Yuen T, Chi L, Rivier J, Millar R, Weinstein H, Sealfon S, Multiple Interactions of the Asp 2.61(98) Side Chain of the Gonadotropin-Releasing Hormone Receptor Contribute Differentially to Ligand Interaction †. *Biochemistry.* 2000;39:8133–41.
16. Harms MJ, Schlessman JL, Sue GR, García-Moreno EB. Arginine residues at internal positions in a protein are always charged. *Proc. Natl. Acad. Sci. U. S. A.* 2011;108:18954–18959
17. Agrawal S, Mohale D, Talele G. Pharmacological activities of *Tridax procumbens* (Asteraceae). *Med. Plants - Int. J. Phytomedicines Relat. Ind.* 2010;2:73.
18. Beck S, Mathison H, Todorov T, Calderon-Juarez EA, Kopp, O., A Review of Medicinal Uses and Pharmacological Activities of *Tridax Procumbens* (L.). *J. Plant Stud.* 2018;7:19.
19. Sen K, Horrell S, Kekilli D, Yong CW, Keal TW, Atakisi H, Moreau DW, Thorne RE, Hough MA, Strange RW. Active-site protein dynamics and solvent accessibility in native *Achromobacter cycloclastes* copper nitrite reductase. *IUCr J.* 2017;4:495–505.
20. Moiseev VM, Rodina EV, Kurilova SA, Vorobyeva NN, Nazarova TI, Avaeva SM. Substitutions of glycine residues Gly100 and Gly147 in conservative loops decrease rates of conformational rearrangements of *Escherichia coli* inorganic pyrophosphatase. *Biochem. Biokhimiia* 2005;70: 858–866.
21. Mishra V, Kumar A, Ali V, Nozaki T, Zhang KYJ, Bhakuni V. Role of conserved active site tryptophan-101 in functional activity and stability of phosphoserine aminotransferase from an enteric human parasite. *Amino Acids* 2012;43:483–491.
22. Lapelosa M, Burrone O, Rocchia W. Specific Residue Interactions Regulate the Binding of Dengue Antigens to Broadly Neutralizing EDE Antibodies. *Chemistry Open.* 2018;7:604–610.
23. Chuang CH, Chiou S, Cheng TL, Wang YT. A molecular dynamics simulation study decodes the Zika virus NS5 methyltransferase bound to SAH and RNA analogue. *Sci. Rep.* 2018;8:6336.
24. Shinde RN, Karthikeyan S, Singh B. Molecular dynamics studies unravel role of conserved residues responsible for movement of ions into active site of DHBPS. *Sci. Rep.* 2017;7:40452.
25. Cheng H, Rashid S, Yu Z, Yoshizumi A, Hwang E, Brodsky B. Location of Glycine Mutations within a Bacterial Collagen Protein Affects Degree of Disruption of Triple-helix Folding and Conformation. *J. Biol. Chem.* 2011;286:2041–2046.
26. Manogar P, Vijayakumar S, Rajalakshmi S, Pugazhenti M, Praseetha PK, Jayanthi, S., In silico studies on CNR1 receptor and effective cyanobacterial drugs: Homology modelling, molecular docking and molecular dynamic simulations. *Gene Rep.* 2019; 17, 100505.
27. Pushyarag P Venugopal and Debashree Chakraborty. Molecular mechanism of inhibition of COVID-19 main protease by β -adrenoceptor agonists and adenosine deaminase inhibitors using in silico methods. *J Bio Struct Dyn.* 2020;1-16
28. Kacar AK, Bahadori F, Tekkeli SEK, Topcu G, Bolkent S. Investigation of cell death mechanism and activity of esculetin-loaded PLGA nanoparticles on insulinoma cells *In vitro*. *J. Pharm. Pharmacol.* 2020;72:592–606.
29. Witaicenis A, Luchini AC, Hiruma-Lima CA, Felisbino SL, Justulin LA, Garrido-Mesa N, Utrilla P, Gálvez J, Di Stasi LC. Mechanism and Effect of Esculetin in an Experimental Animal Model of Inflammatory Bowel Disease. *Eur. J. Inflamm.* 2013;11:433–446.
30. Jeon YJ, Jang JY, Shim JH, Myung PK, Chae JI. Esculetin, a Coumarin Derivative, Exhibits Anti-proliferative and Pro-apoptotic Activity in G361 Human Malignant Melanoma. *J. Cancer Prev.* 2015;20:106–112.
31. Ajmal Ali M. Molecular docking and molecular dynamics simulation of anticancer active ligand '3,5,7,3',5'-pentahydroxy-flavanonol-3-O- α -L-rhamnopyranoside' from *Bauhinia strychnifolia* Craib to the cyclin-dependent protein kinase. *J. King Saud Univ. - Sci.* 2020;32:891–895.
32. Choudhary MI, Shaikh M, tul-Wahab A, ur-Rahman A. *In silico* identification of potential inhibitors of key SARS-CoV-2 3CL hydrolase (Mpro) via molecular docking, MMGBSA predictive binding energy calculations, and molecular

- dynamics simulation. PLOS ONE. 2020; 15.
33. Priya R, Sumitha R, Doss CGP, Rajasekaran C, Babu S, Seenivasan R, Siva R. Molecular Docking and Molecular Dynamics to Identify a Novel Human Immunodeficiency Virus Inhibitor from Alkaloids of *Toddalia asiatica*. Pharmacogn. Mag. 2015;11:S414–S422.
34. Reza R, Dutta T, Ghosh NN. et al. Inhibitory efficiency of quinazoline derivatives against SARS-CoV-2: virtual screening and molecular dynamics study. Proc.Indian Natl. Sci. Acad; 2024.
35. Sahu R, Gupta PK, Mishra A, Kumar A. Ayurveda and in silico Approach: A Challenging Proficient Confluence for Better Development of Effective Traditional Medicine Spotlighting Network Pharmacology. Chin J Integr Med. 2023;29 (5):470-480.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/126154>