



Induction and Treatment of Polycystic Ovary Syndrome with Plant Extract in Rats: A Systematic Review

Wuraola Sarah Esan ^a, Christopher Oloruntoba Akintayo ^a,
Olurotimi Sanya ^a, Osiyemi A. Oluwaseun ^b,
Gbemisola Olubayo ^c and Ademola L. Adelekan ^{c*}

^a Department of Physiology, Afe Babalola University, Ado-Ekiti, Nigeria.

^b Biomedical Research Centre, Forestry Research Institute, Nigeria.

^c Department of Public Health and Biological Sciences, Blue Gate Research institute, Ibadan, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJMP/2023/v34i21124

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc., are available here: <https://www.sdiarticle5.com/review-history/96171>

Systematic Review Article

Received: 08/12/2022

Accepted: 15/02/2023

Published: 09/03/2023

ABSTRACT

Aims: To review the induction and treatment of polycystic ovarian syndrome (PCOS) with plant extract in rats in light of the rising incidence of the condition, its accompanying physical and mental issues, and the role that sex hormone alterations play in its development.

Methodology: An extensive literature search was carried out using relevant publications, which were published between 2012 and 2022 and indexed in Google Scholar, PubMed, Elsevier, Scientific Information Database, and Science Direct were investigated.

Results: This review shows that the majority of research examined the levels of sex hormones in the serum, the induction of PCOS, and the treatment of animal models. The review also found that

*Corresponding author: E-mail: ademola@bgri.org.ng, ademola.adelekan@gmail.com.;

43.75% of the studies induced PCOS with Letrozole, 31.25% with estradiol valerate, 6.25% with Testosterone Enanthate (TE), and 6.25% of the studies induced PCOS with Testosterone Propionate (TP).

Conclusion: This study showed that Letrozole, Estradiol valerate, Dhea, Testosterone Enanthate, and Testosterone Propionate effectively induce PCOS in rats. Also, plant extracts are effective in treating PCOS. More research into the mechanisms of action for plant extracts is required to complete our understanding of the reproductive endocrinological effects of plant extract treatment for PCOS.

Keywords: Endocrinological effects; plant extract treatment; polycystic ovarian syndrome.

1. INTRODUCTION

A very prevalent and diverse endocrine disease called polycystic ovarian syndrome (PCOS) causes roughly 75 percent of infertility cases [1,2]. "The Stein-Leventhal syndrome, first published in 1935, was the name given to the polycystic ovary condition at that time. Initially, the clinical triad of hirsutism, amenorrhea, and obesity, as well as pathognomonic ovarian abnormalities, were necessary for diagnosis" [3]. "In 1986, Brower used a single intramuscular (i.m.) injection of estradiol valerate (EV) in 8-week-old rats to induce PCOS in the rodents experimentally. The rats lost their ability to ovulate and manifested traits similar to those of PCOS in humans, such as big cystic follicles in the ovaries and altered luteinizing hormone concentrations" [4]. "Premature puberty, hirsutism, atypical menstrual cycles, early adulthood, middle age, and later life (diabetes mellitus and cardiovascular illnesses) are just a few examples of how PCOS symptoms can appear at any age" [5]. "Its primary characteristics are anovulation, hyperandrogenism, and polycystic ovarian morphology, and its main clinical symptoms are menstrual irregularities, infertility, hypertrichosis, and metabolic syndromes, which include obesity, dyslipidemia, and type 2 diabetes" [6]. "While the primary cause of this condition is still unknown, it has been hypothesized that genetics, insulin resistance, obesity, inflammation, and oxidative stress may contribute to its pathogenesis" [7,8]. "Acne, obesity, hair loss, hypertension, and irregular ovulation are symptoms of this condition in women" [9].

Due to underlying metabolic abnormalities, being overweight worsens all PCOS symptoms [10]. According to [11], these patients have changed endocrine hormone levels and are at risk for infertility and metabolic diseases, including type 2 diabetes. Treatment options for moderate syndromes (a situation in which the disease's symptoms are not severe) include oral

contraceptives or conventional herbal medicines [12]. Studies on theca cells, both *in vivo* and *in vitro*, indicated that ovarian theca cells in PCOS-afflicted women are significantly more active than those in healthy women in converting androgenic precursors into testosterone. As a result of theca cells producing androgen in response to luteinizing hormone (LH), individuals with PCOS have higher blood levels of androgens [13].

This disorder affects women's ability to conceive, mostly due to hormonal imbalances. High plasma androgen concentrations are common in patients. Patients with PCOS may experience increased androgen production due to metabolic abnormalities, including insulin resistance that causes high plasma androgen (HPA) axis activation [14,15]. Multiple cysts develop in PCOS patients' ovarian tissue due to hyperandrogenism and high levels of the luteinizing hormone [16,17]. "In contrast to follicular stimulating hormone (FSH), PCOS increased the frequency of gonadotropin-releasing hormone (GnRH) pulses" [18]. "Theca cells produce more androgens due to this rise in LH concentration, but a relative FSH shortage lowers granulosa cells' capacity to convert androgen into estrogen and hinders follicle development and ovulation" [19]. "PCOS has been treated with various therapeutic approaches, including lifestyle changes, surgery, and medications such as clomiphene citrate, metformin, letrozole, and tamoxifen" [20]. Our goal is to review the induction and treatment of PCOS with plant extract in rats in light of the rising incidence of the condition, its accompanying physical and mental issues, and the role that sex hormone alterations play in its development.

2. MATERIALS AND METHODS

2.1 Design

This systematic review was conducted and reported per the Preferred Reporting Items for

Systematic Reviews and Meta-Analyses (PRISMA).

2.2 Search Strategy

Different databases were searched to get the necessary materials needed for this review. "A scientific literature search was done using Elsevier, Google scholar, PubMed, and Springer link databases. Additionally, literature was systematically searched from research gate, Cochrane library, and the Directory of Open Access Journal (DOAL). For studies that may have been missed in the electronic search, cross reference was undertaken using reference lists of all identified articles [21]. The first search occurred between October 05 and 07, 2022, while the second occurred between October 15 and 20, 2022. "Detailed inclusion and exclusion criteria were developed cautiously to make sure that they match the review questions and involve sufficient details to help point out all relevant studies and exclude irrelevant ones" [21].

2.3 Inclusion Criteria

The inclusion criteria used for this review include qualitative and quantitative studies on PCOS in low and middle-income countries.

2.4 Exclusion Criteria

Articles were excluded if no data, editorials, or short commentaries were presented for the desired outcome. Other exclusion criteria were papers that were not peer-reviewed and those that the full text could not be assessed.

2.5 Data Extraction

Standardized forms developed by the authors were used for data extraction to minimize the risk of bias. One of the authors extracted data from the included studies, and two of the other authors checked these data. Discrepancies were resolved by referring to the original studies. Data were extracted on induction and treatment of PCOS plant extract in rats. One of the authors

independently considered the potential eligibility of each of the abstracts and titles from the retrieved citations and requested full-text versions of these potentially eligible studies. Two authors independently assessed the full text of the potentially eligible publications. Disagreements were resolved by consensus. The initial agreement was obtained on 80 items (95%), and discrepancies were discussed between authors until 100% agreement was obtained. The following information was extracted from the included studies: authors, title, study population, state, objective(s), study design, and findings.

Table 1. Search terms used in the literature search

AND	
PCOS*	Rat*
PCOS* Induction*	Plant extract * Rat*
PCOS* Treatment*	Plant extract* Rat*

2.6 Quality Assessment

"The articles used for the review were properly screened. They were first screened using the titles and their abstracts. The full papers were screened, and the similarities of the papers were determined. The selected papers were manually and automatically de-duplicated. The quality assessment of each study selected was based on the criteria used in previous studies [22-23].

2.7 Critical Appraisal

"The qualitative, quantitative, and mixed-method researches were evaluated using the Critical Appraisal Skills Programme [CASP] instrument [24] and Critical Appraisal Framework [25] criteria" [26].

2.8 List of Papers Reviewed

The search results are shown in Fig. 1, and a synopsis of the papers consulted (the PRISMA flow chat). Although the databases contained 57 research articles, only 16 met the inclusion criteria for this systematic review (Table 2).

Table 2. Literature review database

Author	Title	Study design	Country of study	Study duration	Objectives/Research question	Findings
[27]	<i>Crocus sativus</i> (saffron) petals extract and its active ingredient, anthocyanin, improves ovarian dysfunction, regulation of inflammatory genes, and antioxidant factors in testosterone-induced PCOS mice	Female NMRI mice (25–35 g, 6 weeks old) (n = 96) were divided into four groups (n = 12 control group, n = 12 testosterone enanthate (TE) group, n = 36 TE + SPE group, n = 36 TE + SPA group,	Iran	60 days	The effects of saffron petal extract (SPE) and saffron petal anthocyanins (SPA) on ovarian hormones, steroidogenic enzymes, ovarian dysfunction, regulation of anti-inflammatory genes, and antioxidant factors in female PCOS mice were studied	These results suggest that SPE and SPA ameliorate symptoms of PCOS by improving the dysregulation of ovarian steroids, steroidogenic, antioxidant enzymes, and inflammatory markers in PCOS mice.
[28]	Comparison of the effects of Ginger extract with clomiphene citrate on sex hormones in rats with the polycystic ovarian syndrome	63 adult female rats (170-200 g) were studied and divided randomly into 9 groups as control, sham, and 7 experimental groups (estradiol valerate used as PCOS-inducing agent, intramuscular) orally daily for 60 and 89 days.	Iran		To compare the effectiveness of Ginger with clomiphene on sexual hormones such as Luteinizing hormone (LH), Follicle stimulating hormone (FSH), estrogen, and progesterone to treat PCOS effectively with fewer side effects.	As the long-term administration of clomiphene citrate has some side effects, the use of Ginger as a herbal medicine without any side effects at high doses can be an effective and good alternative in improving PCOS
[29]	The ameliorative effects of marjoram in dehydroepiandrosterone-induced polycystic ovary syndrome in rats	75 postpubertal (42 days old) female Wistar rats were randomly assigned into five groups (control, dehydroepiandrosterone (DHEA) induced-PCOS model, marjoram-treated PCOS rats, metformin-treated PCOS rats and the combination of marjoram+metformin treated PCOS model)	Jordan		To investigate the ameliorative effects of marjoram extract on hormonal profiles, body and ovaries weight, insulin sensitivity, inflammation, and oxidative stress in a rat model of PCOS	The current study showed that marjoram significantly decreased ovaries' weight and estradiol levels (P-value < 0.05) compared to the DHEA group. Interestingly, marjoram improved insulin sensitivity as manifested by a significant increase in the adiponectin serum levels (P-value < 0.05).
[30]	Effect of <i>Aloe barbadensis</i> Mill. The formulation of Letrozole-induced polycystic ovarian syndrome rat model	The PCOS rat model was developed with 12-16 adult virgin Charles Foster female rats	India		To evaluate the efficacy of <i>Aloe barbadensis</i> gel formulation in a PCOS rat model.	<i>Aloe barbadensis</i> gel formulation exerts a protective effect against

Author	Title	Study design	Country of study	Study duration	Objectives/Research question	Findings
		weighing 200--225g.				the PCOS phenotype by restoring the ovarian steroid status and altering key steroidogenic activity. This can be attributed to phyto-components present in the extract.
[31]	Effect of <i>Citrullus colocynthis</i> hydro-alcoholic extract on hormonal and folliculogenesis process in estradiol valerate-induced PCOs rats model: An experimental study	40 female adult Wistar rats divided into five groups	Iran		To evaluate the effect of CCT hydro-alcoholic extract on hormonal and folliculogenesis process in estradiol valerate-induced PCOS rats' model.	Marked improvement in hormonal and histological symptoms of PCOS may be due to the CCT effect. Hence, CCT can potentially be considered an effective drug for treating PCOS.
[32]	Effect of <i>Vitex agnus-castus</i> ethanolic extract on hypothalamic KISS-1 gene expression in a rat model of polycystic ovary syndrome	Thirty-two female rats were distributed into: control, Vitagnus-treatment (365 mg/kg for 30 days), PCOS (Letrozole for 28 days) and PCOS animals treated with Vitagnus (30 days of Vitagnus after PCOS induction).	Iran	30 days	To investigate the Vitagnus effect on the expression of the kisspeptin gene in a rat model of PCOS	The results indicated that Vitagnus extract inhibited the downregulation of the KISS-1 gene in the hypothalamus of PCOS rats.
[33]	Effects of Chamomile extract on biochemical and clinical parameters in a rat model of polycystic ovary syndrome	Thirty virgin adults cycling Wistar rats, weighting 200 - 220 g, were divided into two groups	Iran		To evaluate the effects of Chamomile alcoholic extract on the biochemical and clinical parameters in a rat model of PCOS.	The histological and hormonal results showed that Chamomile could decrease the signs of PCOS in the ovarian tissue and help LH secretion in rats (p < 0.05)
[34]	Effects of <i>Fagonia indica</i> on letrozole-induced polycystic ovarian syndrome (PCOS) in young adult female rats	Twenty-five healthy female Wistar albino rats (150–200 g)	Pakistan		To investigate the effectiveness of ethanolic extract of <i>Fagonia indica</i> in letrozole-induced PCOS young adult female rats	This study validates the potential of <i>Fagonia indica</i> for the amelioration of metabolic and

Author	Title	Study design	Country of study	Study duration	Objectives/Research question	Findings
[35]	Hydroalcoholic extract of flax seed improves polycystic ovary syndrome in a rat model	Twenty-four rats were divided into four groups: negative control, positive control, PCOS, and treatment groups.	Iran		The effect of hydroalcoholic extract of flax seed was evaluated on ovarian hormones and histological changes of the uterus and ovary in a PCOS-induced rat model	hormonal disturbances that occur in PCOS. Hormonal profile and histomorphometric features of the ovary that were disturbed by PCOS induction were ameliorated by hydroalcoholic extract of flax seed
[36]	<i>Phyllanthus muellerianus</i> (Euphorbiaceae) restores ovarian functions in letrozole-induced polycystic ovarian syndrome in rats.	One hundred and eight adult female Wistar rats (180-200 g) were used.	Cameroon		To investigate the effects of <i>P. muellerianus</i> extracts on estrus cyclicity, lipid profile, oxidative stress-related markers, sex hormones, and ovarian architecture in Letrozole-Induced PCOS in rats.	This Plants could be useful in the management/treatment of reproductive and metabolic disorders related to PCOS.
[37]	Licorice ethanol extract improves symptoms of polycystic ovary syndrome in Letrozole-induced female rats.	Sprague Dawley rats were obtained from Daehan Biolink and divided into three groups of six rats each. For the PCOS rat model, a 90-day release pellet containing Letrozole (1.8 mg/pellet) was implanted subcutaneously for 4 weeks under anesthesia [Zoletile (30 mg/kg)–Rompun (5 mg/kg)–saline mixture (2:1:2)] in 6-week-old female rats.	Korea		To examine the effects of GRR (<i>Glycyrrhizae radix et rhizome</i>) extract on PCOS-like symptoms in female rats.	These results suggest that GRR extract inhibits the symptoms of PCOS by regulating imbalanced hormonal levels and irregular ovarian follicles
[38]	<i>Ocimum kilimandscharicum</i> L. restores ovarian functions in Letrozole - induced polycystic ovary syndrome (PCOS) in rats: comparison with metformin	Adult virgin female Wistar Albino rats (6 weeks old, weighing 150–200 g; total n = 54, n = 6)	Egypt		To investigate the therapeutic merits of <i>Ocimum kilimandscharicum</i> (Ok), in a letrozole PCOS rat model, and compare it to metformin	These results suggest that Ok extract and EA (ethyl acetate) fraction halt letrozole-induced reproductive dysfunctions and restore normal morphological and physiological functions in PCOS rats, even superior to metformin.
[39]	A novel potential reproductive effects of <i>Pterocarpus</i>	Cyclic, virgin female albino rats	India	15 days.	To investigate the potential	<i>Pterocarpus marsupium</i>

Author	Title	Study design	Country of study	Study duration	Objectives/Research question	Findings
	<i>marsupium</i> methanolic extract on testosterone propionate Induced Polycystic Ovary Syndrome in Female Albino Rat	(weighing 180-200g), All the experimental animals except the control group were injected with Testosterone Propionate (TP) intraperitoneally at a dose of 1mg/100gm b.w			reproductive effects of <i>Pterocarpus marsupium</i> methanolic extract on testosterone propionate-induced PCOS in female albino rats	showed potential reproductive effects on testosterone propionate-induced PCOS in female albino rats and could be used as an alternative therapy in the treatment of PCOS
[40]	The effect of hydroalcoholic extract of <i>Nigella Sativa</i> seed on the dehydroepiandrosterone-induced polycystic ovarian syndrome in rats: An experimental study	36 female Wistar rats (60 ± 10 g, aged 21 days)	Iran		To investigate the therapeutic effect of hydroalcoholic extract of <i>N. sativa</i> seed on dehydroepiandrosterone (DHEA)-induced PCOS rats.	The histopathological results, which are in accordance with biochemical findings, imply that <i>N. sativa</i> seed could be useful in the treatment of PCOS, the higher doses of the extract being more effective.
[41]	The effects of <i>Urtica dioica</i> extract on lipid profile, insulin resistance index and liver histology in polycystic ovary syndrome-induced Wistar rats	One hundred forty-four adult Wistar rats were divided into control, PCOS, and nettle-treated groups. The group PCOS was injected subcutaneously with 2 mg estradiol valerate after 60 days	Iran		To examine the <i>Urtica dioica</i> moderator effect on liver function in PCOS rats	The results show that <i>Urtica dioica</i> , by increasing insulin sensitivity and reducing hepatic necrosis, may reduce inflammation, improve metabolic symptoms in PCOS, and significantly protect the liver.
[42]	The aqueous leaf extract of <i>Myrianthus arboreous</i> P. Beauv. (Cecropiaceae) improved letrozole-induced polycystic ovarian syndrome-associated conditions and infertility in female Wistar rats	Sixty female Wistar rats were used to evaluate the effects of the aqueous extract of <i>M. arboreus</i> leaves on PCOS-associated symptoms and PCOS-related infertility.	Cameroon		To evaluate the effects of such an extract on an animal model of infertility caused by polycystic ovary syndrome (PCOS) to bring scientific evidence to the curative action of this plant against female infertility	the traditional use of <i>M. arboreus</i> against female infertility suggests that this plant could be a promising alternative treatment to improve symptoms associated with different PCOS phenotypes.

3. RESULTS

All publications reviewed were published between 2010 and 2022, with 21.42% published in 2017 (Fig.3) and 42.85% carried out in Iran (Fig. 2).

3.1 The Induction of Pcos with Plant Extract in Rats

Among the studies reviewed, it was found that 43.75% of the studies induced PCOS with Letrozole [30,34,36,37,38,42,32], 31.25% of the studies induced PCOS with estradiol valerate [28,31,33,35,41], 12.5% induced with Dhea [29,40], 6.25% induced with Testosterone Enanthate (TE) [27] and 6.25% of the studies

induced PCOS with Testosterone Propionate TP) [39].

3.2 The Treatment of PCOS with Plant Extract in Rats

A study [27] reveals that “in SPE (50, 200 and 600 mg/kg/BW) and SPA (20, 40 and 80 mg/kg/BW) treatment in TE-induced PCOS mice, the level(s) of LH, FSH, and steroid hormones, i.e., estrogen, progesterone, and testosterone, were estimated. The LH, estrogen, and testosterone concentration significantly increased ($P < 0.01$), and the FSH and progesterone concentration reduced ($P < 0.001$) following TE treatment as compared with the control”.




Fig. 1. PRISMA Flow chart of the study selected

According to [28], “the results of this study showed that LH serum level was significantly increased in the sham group (89 days) compared to control groups and sham group (60 days) ($p < 0.001$). These results showed that clomiphene and ginger extract could have positive and dose-dependent effects in PCOS treatment. In addition, experimental group 5 showed a higher increase in FSH serum level than experimental group 4. The results of serum concentration measurement of FSH showed that in experimental groups 1 and 2, FSH significantly reduced compared to control and sham groups ($p < 0.001$). In the experimental groups 3, 4, and 5, there was a significant decrease in serum estrogen and progesterone level, compared to experimental groups 1 and 2”.

Another study [29] found that “after 21 days of treatment, progesterone and estradiol levels were significantly higher in the DHEA group in comparison with the control group (P -value < 0.01). The serum progesterone level was significantly reduced in the metformin-treated group (group 4) compared to the DHEA group (P -value < 0.05). Furthermore, the levels of estradiol were significantly reduced following the administration of marjoram (group 3), metformin (group 4), or the combination treatment (group 5) in comparison to the DHEA group (P -value < 0.05). Conversely, the current study showed no significant changes in testosterone levels

detected among different study groups (P -value > 0.05)”.

In a study by [31], “it was revealed that there was a significant reduction in luteinizing hormone and testosterone in III-V groups compared to Sham group, whereas follicle-stimulating hormone in III-V groups was not significantly changed in comparison with Sham group.”

“It observed that vitagnus treatment in the PCOS group resulted in a rise in progesterone, estrogen, and FSH levels and a reduction in the levels of testosterone and LH” [32].

A study [33] found out that “the serum levels of estradiol and gonadotropins, LH and FSH, were significantly decreased in the Chamomile group relevant to the control group ($p < 0.05$)”.

In a study by [34], “it was seen that group II (PCOS) showed a significant increase ($p < 0.05$) in LH levels as compared to group I (normal control). LH levels decreased significantly in group III ($p < 0.05$) and group V ($p > 0.05$), while a decrease was observed in group IV when compared with group II. A significant ($p < 0.0001$) reduction was observed in FSH, estradiol, and progesterone concentrations in group II compared to group I. Estradiol levels were significantly improved ($p < 0.0001$) in all treated groups as compared with group II”.




Fig. 2. Study location




Fig. 3. Study Year

According to [35], “it was also reported that the level of estradiol in the PCOS and treatment groups increased compared to the positive and negative controls ($P < 0.05$). At the same time, there was no significant difference in estradiol levels between the PCOS and treatment groups. Progesterone concentration in the PCOS group decreased compared to the control and treatment groups ($P < 0.05$); however, the level of progesterone in the treatment group did not have a significant difference from the control groups ($P > 0.05$). In addition, the mean testosterone level in the PCOS group was higher than the control and treatment groups ($P < 0.05$). On the other hand, there was no significant difference in testosterone levels between the control and treatment groups. There was no significant difference between the groups regarding the level of DHEA ($P < 0.05$)”.

In a study [39], “it was observed that the hormonal parameters such as FSH, LH and estrogen levels in standard clomiphene citrate and *Pterocarpus marsupium* methanol extract treatment with low and high dose groups have exhibited a significant increase ($p < 0.01$) in their levels and a significant decrease ($p < 0.01$) in testosterone levels on Day 30”. “However, it was reported that the serum level of FSH was reduced only in the letrozole treatment group (PCOS), whereas significant recovery of FSH level was observed in the Letrozole and GRR co-treatment group (PCOS + GRR). Serum LH levels were not altered in any of the groups. Furthermore, the LH/FSH ratio (known biomarker

for PCOS) was elevated only in the letrozole treatment group (PCOS). In contrast, it was significantly reduced in the Letrozole and GR co-treatment group (PCOS + GRR)” [37].

“Furthermore, a study revealed deleterious effects of Letrozole on the hormonal profile, where significant elevation in testosterone and a significant reduction in estrogen and progesterone levels by 153%, 27%, and 9%, respectively, were reported, compared to control animals. Treatment of PCOS rats with metformin (Gp IIb), Ok extract (Gp IIc), and EA fraction (Gp II d) decreased testosterone levels by 40%, 42%, and 73%, significantly increased estrogen levels by 225%, 579%, and 94%, and also increased progesterone levels by 321%, 500%, and 813%, respectively, compared to PCOS untreated rats (Gp IIa)”. [38]. A study [36] reported that “*P.muellerianus* significantly decreased ($p < 0.001$) LH and testosterone (both extracts; 30, 60, and 120 mg/kg) levels, but increased ($p < 0.01$) estradiol (aqueous extract; 60 mg/kg) concentration”.

A study [40] also observed that “the treatment of PCO animals with metformin and doses of 50, 100, and 200 mg/kg of the *N. sativa* extract decreased LH levels ($p < 0.001$ and $p = 0.05$ respectively). The level of FSH decreased in PCOS rats, while FSH increased in other groups. The testosterone levels were elevated in PCOS rats ($p < 0.001$), and treatment with metformin and a dose of 200 mg/kg *N. sativa* reduced it. A 200 mg/kg dose of *N. sativa* extract increased

the progesterone level ($p = 0.03$). The present study's data showed that the estrogen level upregulated in the PCOS group while it decreased in all treatment groups”.

“However, it was reported that the aqueous extract of *M. arboreus* leaves induced a similar effect at doses of 20 mg/kg (37% induction; $p < 0.001$) and 110 mg/kg (95% induction; $p < 0.001$). The serum levels of LH increased by 310% ($p < 0.001$) in the LTZ group compared to the normal control group. The combined administration of clomiphene citrate and metformin decreased this parameter by 77% ($p < 0.001$) compared to the LTZ group. The aqueous extract of *M. arboreus* leaves induced a similar effect at tested doses. It was shown that estradiol serum levels of estradiol decreased by 84% ($p < 0.001$) in the LTZ group compared to the normal control group. The combined administration of clomiphene citrate and metformin increased this parameter by 764% ($p < 0.001$) compared to the LTZ group. The aqueous extract of *M. arboreus* leaves also induced a similar effect as it increased serum levels of estradiol by 74%, 219%, and 87% at doses of (20, 110, and 200) mg/kg, respectively” [42].

4. DISCUSSION

We carried out this review to introduce plants that have recently been studied for their effects on PCOS with animal models and to give evidence on the induction and treatment of PCOS. The majority of research examined the levels of sex hormones in the serum, the induction of PCOS, and the treatment of animal models.

Majority of the studies reviewed successfully administered Letrozole to induce PCOS in the rat model. It is a nonsteroidal and highly potent aromatase inhibitor [37]. Among the other studies, estradiol valerate was used to induce PCOS [35], as reviewed. “Following the induction of PCOS, estradiol and testosterone levels significantly increased, while the progesterone level significantly decreased in these groups; however, no significant change in levels of DHEA was observed. Other researchers also reported similar results following induction of PCOS by estradiol valerate” [43,44]. “In this present review, the level(s) of LH and FSH in the serum of TE-induced mice [27] was observed”. “Elevated level(s) of LH, estrogen, and testosterone significantly decreased following treatment with

SPE. In addition, a decrease in progesterone and FSH level(s) following treatment with TE was observed, and their reduction was restored by SPE treatment in TE-induced mice. According to research, premenstrual syndrome is caused by changes in estrogen and progesterone, such that the amount of progesterone compared to estrogen decreases. SPE causes a natural balance between estrogen and progesterone during menstruation. With its physiological-pharmacological effects, the saffron petal plant balances the decrease or increase of hormones in the body” [45].

“In the present review, previous studies reported the enhancement in serum level of LH and estrogen hormone and reduction in the level of FSH and progesterone. On the other hand, any agent that could reduce estrogen and LH and increase FSH and progesterone levels can be used to treat PCOS disorder. Previous studies stated that hyperandrogenism and increased serum levels of LH are very common in PCOS” [46]. “Some other reviews demonstrated that PCOS could increase secretion of testosterone and LH and reduce FSH hormones secretion” [47]. “One possible mechanism for an explanation of sex hormones changes in PCOS is relative lack of aromatase enzyme in the ovary which could increase androgen concentration” [48]. “Herbal medicine usage by women has increased over the past decade. Herbal remedies contain pharmacologically active constituents with physiological effects on female endocrinology. They are positively associated with reduced breast, bone, and cardiovascular disease incidence” [49,50].

“Previous studies demonstrated that inducing PCOS with letrozole led to higher amounts of testosterone, elevated ratio of LH/FSH, higher numbers of follicular cysts, the elevated thickness of the theca layer, and decreased thickness of granulosa layer” (Sun *et al.*, 2013). “All these key symptoms were observed in the current study, indicating successful induction of PCOS in our experimental animals. Increased levels of LH cause the increased production of testosterone, leading to the follicular arrest and increased AMH levels” [51]. “Polycystic ovarian syndrome is a complex endocrine disorder that is most frequently encountered gynecological endocrinopathy among reproductive-aged women” [52]. A variety of genetic and environmental variables causes the pathogenesis of this syndrome. An unhealthy

lifestyle and diet, as well as any infectious mediators, increase the likelihood of PCOS.

5. CONCLUSION

In conclusion, this study showed that Letrozole, Estradiol valerate, Dhea, Testosterone Enanthate, and Testosterone Propionate effectively induce PCOS in rats. Also, plant extracts are effective in treating PCOS.

More research into the mechanisms of action for plant extracts is required to complete our understanding of the reproductive endocrinological effects of plant extract treatment for PCOS. Also, a clinical trial is recommended to ascertain these plant extracts' efficacy in humans.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

The authors acknowledge the effort of the Blue Gate Research Institute staff to review this paper.

COMPETING INTERESTS

The authors have declared that no competing interests exist.

REFERENCES

1. Mohammadi S, Bardei LK, Hojati V, Ghorbani A, Nabiuni M. Anti-inflammatory effects of curcumin on insulin resistance index, levels of interleukin-6, C-reactive protein, and liver histology in polycystic ovary syndrome-induced rats. *Cell Journal (Yakhteh)*. 2017;19(3):425.
2. Bakhshalizadeh S, Amidi F, Shirazi R, Shabani Nashtaei M. Vitamin D3 regulates steroidogenesis in granulosa cells through AMP-activated protein kinase (AMPK) activation in a mouse model of polycystic ovary syndrome. *Cell biochemistry and function*. 2018;36(4):183-93.
3. Stein IF. Amenorrhea associated with bilateral polycystic ovaries. *Am J Obstet Gynecol*. 1935;29:181-91.
4. Brawer JR, Munoz M, Farookhi R. Development of the polycystic ovarian condition (PCO) in the estradiol valerate-treated rat. *Biology of reproduction*. 1986;35(3):647-55.
5. Norman R, Wu R, Stankiewicz M. Polycystic ovary syndrome; 2004.
6. Hu Q, Jin J, Zhou H, Yu D, Qian W, Zhong Y, Zhang J, Tang C, Liu P, Zhou Y, Wang X. Crocetin attenuates DHT-induced polycystic ovary syndrome in mice via revising kisspeptin neurons. *Biomedicine & Pharmacotherapy*. 2018;107:1363-9.
7. Goodarzi MO, Dumesic DA, Chazenbalk G, Azziz R. Polycystic ovary syndrome: Etiology, pathogenesis and diagnosis. *Nature reviews endocrinology*. 2011;7(4):219-31.
8. Murri M, Luque-Ramírez M, Insenser M, Ojeda-Ojeda M, Escobar-Morreale HF. Circulating markers of oxidative stress and polycystic ovary syndrome (PCOS): A systematic review and meta-analysis. *Human reproduction update*. 2013;19(3):268-88.
9. Hoeger KM, Legro RS, Welt CK, editors. A patient's guide: Polycystic ovary syndrome (PCOS). *The Journal of Clinical Endocrinology & Metabolism*. 2014;99(1):35A-6A.
10. Pages ES, Tarlatzis BC, Rebar RW. Consensus on women's health aspects of polycystic ovary syndrome (PCOS). *Hum Reprod*. 2012;27(1):14-24.
11. Lizneva D, Suturina L, Walker W, Brakta S, Gavrilova-Jordan L, Azziz R. Criteria, prevalence, and phenotypes of polycystic ovary syndrome. *Fertility and sterility*. 2016;106(1):6-15.
12. van Die MD, Burger HG, Teede HJ, Bone KM. Vitex agnus-castus extracts for female reproductive disorders: A systematic review of clinical trials. *Planta medica*. 2013;79(07):562-75.
13. Rosenfield RL, Ehrmann DA. The pathogenesis of polycystic ovary syndrome (PCOS): The hypothesis of PCOS as functional ovarian hyperandrogenism revisited. *Endocrine reviews*. 2016; 37(5):467-520.
14. Conway G, Dewailly D, Diamanti-Kandarakis E, Escobar-Morreale HF, Franks S, Gambineri A, Kelestimur F, Macut D, Micic D, Pasquali R, Pfeifer M. The polycystic ovary syndrome: A position statement from the European Society of

- endocrinology. *European Journal of Endocrinology*. 2014;171(4):P1-29.
15. Sirmans SM, Pate KA. Epidemiology, diagnosis, and management of polycystic ovary syndrome. *Clinical epidemiology*. 2014;6:1.
 16. Imani B, Eijkemans MJ, te Velde ER, Habbema JD, Fauser BC. A nomogram to predict the probability of live birth after clomiphene citrate induction of ovulation in normogonadotropic oligoamenorrhic infertility. *Fertility and sterility*. 2002;77(1):91-7.
 17. Tarlatzis BC, Fauser BC, Legro RS, Norman RJ, Hoeger K, Pasquali R, Franks S, Messinis IE, Casper RF, Homburg R, Lobo R. Consensus on infertility treatment related to polycystic ovary syndrome. *Human Reproduction*. 2008;23(3):462-77.
 18. Katsayal UA, Lamal RS. Preliminary phytochemical and antibacterial screening of the ethanolic stem bark extract of *Phyllanthus muellerianus*. *Nigerian Journal of Pharmaceutical Sciences*. 2009;8(2):121-5.
 19. Agyare C, Lechtenberg M, Deters A, Petereit F, Hensel A. Ellagitannins from *Phyllanthus muellerianus* (Kuntze) Exell.: Geraniin and furosin stimulate cellular activity, differentiation and collagen synthesis of human skin keratinocytes and dermal fibroblasts. *Phytomedicine*. 2011;18(7):617-24.
 20. Legro RS, Arslanian SA, Ehrmann DA, Hoeger KM, Murad MH, Pasquali R, Welt CK. Diagnosis and treatment of polycystic ovary syndrome: An endocrine Society clinical practice guideline. *The Journal of Clinical Endocrinology & Metabolism*. 2013;98(12):4565-92.
 21. Boland A, Cherry G, Dickson R, editors. *Doing a systematic review: A student's guide*; 2017.
 22. McCormick F, Cvetanovich GL, Kim JM, Harris JD, Gupta AK, Abrams GD, Romeo AA, Provencher MT. An assessment of the quality of rotator cuff randomized controlled trials: Utilizing the Jadad score and CONSORT criteria. *Journal of shoulder and elbow surgery*. 2013;22(9):1180-5.
 23. Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, Leeflang MM, Sterne JA, Bossuyt PM, QUADAS-2 Group*. QUADAS-2: A revised tool for the quality assessment of diagnostic accuracy studies. *Annals of internal medicine*. 2011;155(8):529-36.
 24. Singh J. Critical appraisal skills programme. *Journal of pharmacology and Pharmacotherapeutics*. 2013;4(1):76.
 25. Wright BE, Grant AM. Unanswered questions about public service motivation: Designing research to address key issues of emergence and effects. *Public administration review*. 2010;70(5):691-700.
 26. Cuschieri S. The STROBE guidelines. *Saudi journal of anaesthesia*. 2019;13(Suppl 1):S31.
 27. Moshfegh F, Balanejad SZ, Shahrokhbady K, Attaranzadeh A. *Crocus sativus* (saffron) petals extract and its active ingredient, anthocyanin improves ovarian dysfunction, regulation of inflammatory genes and antioxidant factors in testosterone-induced PCOS mice. *Journal of Ethnopharmacology*. 2022;282:114594.
 28. Atashpour S, Jahromi HK, Jahromi ZK, Maleknasab M. Comparison of the effects of Ginger extract with clomiphene citrate on sex hormones in rats with polycystic ovarian syndrome. *International Journal of Reproductive BioMedicine*. 2017;15(9):561.
 29. Rababa'h AM, Matani BR, Ababneh MA. The ameliorative effects of marjoram in dehydroepiandrosterone induced polycystic ovary syndrome in rats. *Life sciences*. 2020;261:118353.
 30. Maharjan R, Nagar PS, Nampoothiri L. Effect of Aloe barbadensis Mill. Formulation on Letrozole induced polycystic ovarian syndrome rat model. *Journal of Ayurveda and integrative medicine*. 2010;1(4):273.
 31. Barzegar MH, Khazali H, Kalantar SM, Khoradmehr A. Effect of Citrullus colocynthis hydro-alcoholic extract on hormonal and folliculogenesis process in estradiol valerate-induced PCOs rats model: An experimental study. *International Journal of Reproductive BioMedicine*. 2017;15(10):661.
 32. Feyzollahi Z, Kouchesfehiani HM, Jalali H, Eslimi-Esfahani D, Hosseini AS. Effect of Vitex agnus-castus ethanolic extract on hypothalamic KISS-1 gene expression in a rat model of polycystic ovary syndrome. *Avicenna Journal of Phytomedicine*. 2021;11(3):292.
 33. Zangeneh FZ, Minaee B, Amirzargar A, Ahangarpour A, Mousavizadeh K. Effects

- of chamomile extract on biochemical and clinical parameters in a rat model of polycystic ovary syndrome. *Journal of reproduction & infertility*. 2010;11(3):169.
34. Younas A, Hussain L, Shabbir A, Asif M, Hussain M, Manzoor F. Effects of *Fagonia indica* on letrozole-induced Polycystic Ovarian Syndrome (PCOS) in Young Adult Female Rats. *Evidence-Based Complementary and Alternative Medicine*. 2022;2022.
 35. Jelodar G, Masoomi S, Rahmanifar F. Hydroalcoholic extract of flaxseed improves polycystic ovary syndrome in a rat model. *Iranian journal of basic medical sciences*. 2018;21(6):645.
 36. Ndeingang EC, Defo Deeh PB, Watcho P, Kamanyi A. *Phyllanthus muellerianus* (Euphorbiaceae) restores ovarian functions in letrozole-induced polycystic ovarian syndrome in rats. *Evidence-Based Complementary and Alternative Medicine*. 2019;2019.
 37. Yang H, Kim HJ, Pyun BJ, Lee HW. Licorice ethanol extract improves symptoms of polycystic ovary syndrome in Letrozole-induced female rats. *Integrative medicine research*. 2018;7(3):264-70.
 38. Khaled N, El-Bahy AA, Radwan R, Handoussa H, AbdelMaksoud S. *Ocimum kilimandscharicum* L. restores ovarian functions in letrozole-induced Polycystic Ovary Syndrome (PCOS) in rats: Comparison with metformin. *Life sciences*. 2019;232:116640.
 39. L Hugar A, P Kanjekar A, L Londonkar R. A novel potential reproductive effects of *Pterocarpus marsupium* methanolic extract on testosterone propionate induced polycystic ovary syndrome in female albino rats. *Endocrine, Metabolic & Immune Disorders-Drug Targets (Formerly Current Drug Targets-Immune, Endocrine & Metabolic Disorders)*. 2017;17(4):317-23.
 40. Khani S, Abdollahi M, Khalaj A, Heidari H, Zohali S. The effect of hydroalcoholic extract of *Nigella Sativa* seed on dehydroepiandrosterone-induced polycystic ovarian syndrome in rats: An experimental study. *International Journal of Reproductive BioMedicine*. 2021;19(3):271.
 41. Zare S, Nabiuni M, Tayanloo A, Hoseini S, Karimzadeh-Bardei L. The effects of *Urtica dioica* extract on lipid profile, insulin resistance index and liver histology in polycystic ovary syndrome-induced Wistar rats. *Advanced Herbal Medicine*. 2015;1(2):23-33.
 42. Mvondo MA, Mzemdem Tsoplack FI, Awounfack CF, Njamen D. The leaf aqueous extract of *Myrianthus arboreus* P. Beauv.(Cecropiaceae) improved letrozole-induced polycystic ovarian syndrome associated conditions and infertility in female Wistar rats. *BMC Complementary Medicine and Therapies*. 2020;20(1):1-3.
 43. Linares R, Hernández D, Morán C, Chavira R, Cárdenas M, Domínguez R, Morales-Ledesma L. Unilateral or bilateral vagotomy induces ovulation in both ovaries of rats with polycystic ovarian syndrome. *Reproductive Biology and Endocrinology*. 2013;11(1):1-9.
 44. Azarnia M, Kamyab SZ, Mirabolghasemi SG, Saeidnia S. Effect of hydroalcoholic extract of *Melia azedarach* L. seeds on serum concentration of sex hormones in polycystic ovary syndrome induced in female wistar rats. *KAUMS Journal (FEYZ)*. 2015;19(2):111-7.
 45. Agha-Hosseini M, Kashani L, Aleyaseen A, Ghoreishi A, Rahmanpour HA, Zarrinara AR, Akhondzadeh S. *Crocus sativus* L.(saffron) in the treatment of premenstrual syndrome: A double-blind, randomised and placebo-controlled trial. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2008;115(4):515-9.
 46. Arif M, Thakur SC, Datta K. Implication of thymoquinone as a remedy for polycystic ovary in rat. *Pharmaceutical biology*. 2016;54(4):674-85.
 47. Karampoor P, Azarnia M, Mirabolghasemi G, Alizadeh F. The effect of hydroalcoholic extract of fennel (*Foeniculum vulgare*) seed on serum levels of sexual hormones in female wistar rats with polycystic ovarian syndrome (PCOS). *Journal of Arak University of Medical Sciences*. 2014;17(5):70-8.
 48. Requena A, Herrero J, Landeras J, Navarro E, Neyro JL, Salvador C, Tur R, Callejo J, Checa MA, Farré M, Espinós JJ. Use of Letrozole in assisted reproduction: A systematic review and meta-analysis. *Human reproduction update*. 2008;14(6):571-82.
 49. Smith CA, Bateson DJ, Weisberg E. A survey describing the use of complementary therapies and medicines by women attending a family planning clinic. *BMC Complementary and Alternative Medicine*. 2013;13(1):1-8.

50. Stankiewicz M, Smith C, Alvino H, Norman R. The use of complementary medicine and therapies by patients attending a reproductive medicine unit in South Australia: A prospective survey. *Australian and New Zealand Journal of Obstetrics and Gynaecology*. 2007;47(2):145-9.
51. Dinsdale NL, Crespi BJ. Endometriosis and polycystic ovary syndrome are diametric disorders. *Evolutionary applications*. 2021;14(7):1693-715.
52. Pourteymour Fard Tabrizi F, Hajizadeh-Sharafabad F, Vaezi M, Jafari-Vayghan H, Alizadeh M, Maleki V. Quercetin and polycystic ovary syndrome, current evidence and future directions: a systematic review. *Journal of Ovarian Research*. 2020;13(1):1-0.

© 2023 Esan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96171>