



Improving the Effectiveness of Therapeutic Measures for Generalized Periodontitis in Patients with Hypertension

**Alina Alexandrovna Serekova^a, Madina Georgievna Dzgoeva^{a*},
Zalina Georgievna Dzgoeva^a, Azamat Borisovich Kokoev^a,
Larisa Stanislavovna Kanukova^a and Olga Alexandrovna Farnieva^a**

^a North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia, Russia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i52B33623

Editor(s):

(1) Dr. S. Prabhu, Sri Venkateswara College of Engineering, India.

Reviewers:

(1) Golpar Radafshar, Guilan University of Medical Sciences, Iran.

(2) Luma de Oliveira Comini, Federal University of Viçosa, Brazil.

(3) Rajina Shakya, Nepal Health Research Council, Nepal.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:
<https://www.sdiarticle5.com/review-history/76378>

Original Research Article

Received 22 September 2021

Accepted 28 November 2021

Published 02 December 2021

ABSTRACT

The program of standard therapeutic measures in 45 patients with Generalized Chronic Periodontitis (GCP) and background hypertension, group I(a), included hirudotherapy, for comparison, the effectiveness of traditional treatment was evaluated in 40 patients, group I(b), who refused to use medical leeches. In the second comparison group (35 patients), patients with GCP, but normal blood pressure, also received only a set of traditional methods of treatment. The inclusion of hirudotherapy in the program of therapeutic measures in patients with GCP and hypertension makes it possible to optimize the onset of a persistent therapeutic effect, lengthen the period of remission of the pathological process in periodontal disease and significantly improves the quality of life of patients.

Keywords: Generalized chronic periodontitis; hypertension; hirudotherapy; medical leeches.

1. INTRODUCTION

At the present stage of cardiology development, a comprehensive approach to the treatment of hypertension is being implemented, which is based on drug therapy [1-3]. Despite the fact that the main successes in the treatment of hypertension are associated with pharmacotherapy, it is not always possible to achieve high treatment efficiency. In the last decade, non-drug treatment methods have been increasingly used for hypertension: magnetotherapy, physiotherapy, balneotherapy, etc [4-6]. An important component of the complex therapy of hypertension is an adequate correction of hemorheological and microcirculatory disorders characteristic of the disease in the tissues of the entire body [7]. Most researchers have established a more severe course of pathological processes in the tissues of the dental system with background vascular dystonia [8,9].

Unfortunately, the use of a standard arsenal of tools in modern clinical dentistry in the presence of background disorders of systemic hemodynamics cannot achieve complete rehabilitation of the pathological process in periodontitis [10-12]. The problems of effective and pathogenetically justified therapy of generalized chronic periodontitis (GCP) of hypertonia today remain not completely solved [13-16].

The emerging trend in clinical dentistry in recent years is the use of herbal preparations [17-20]. Clinicians over the past decade have also turned to the use of an undeservedly forgotten effective folk method -hirudotherapy [21,22]. All of the above served as the basis for the inclusion of hirudotherapy in the complex of standard methods of therapy for generalized chronic periodontitis .

The aim of the study was to evaluate the effectiveness of hirudotherapy in combination with standard methods of treatment of generalized chronic periodontitis in patients with hypertension.

2. MATERIALS AND METHODS

The study included middle-aged patients - 35-44 years old. Hirudotherapy was included in the comprehensive program of standard therapeutic measures in 45 patients with a chronic course of the inflammatory process in periodontal disease

and background hypertension (group I(a)). For comparison, the effectiveness of traditional treatment was evaluated in 40 patients with GCP on the background of hypertension (group I(b)) who refused to use medical leeches. In the second comparison group (35 patients), patients with CGP, but normal blood pressure, also received only a set of standard traditional methods of treatment.

Patients of both groups were prescribed comprehensive dental treatment, including traditional GCP therapy (removal of dental deposits, anti-inflammatory and antiseptic agents, closed curettage, splinting, etc.). Medical leeches grown in artificial conditions by the company "Girud I.N.", Balakovo, Saratov region, Russia, were used in the work. Leeches from a jar of water were deposited in a warm glass tube and its open end was applied to the inflamed vestibular surface of the alveolar processes of the jaws. The leeches sucked the blood for 30 to 35 minutes. After the procedure, the leech was revitalized - it was dipped in ethyl alcohol. After the procedure, the patient rinses the oral cavity with a solution of hydrogen peroxide (0.5%). It was recommended to abstain from eating for at least 3 hours. During the first application, one leech was used, for the second and subsequent procedures - two. The course of treatment was 10 sessions conducted in one day.

To assess the condition of periodontal tissues before and after treatment, the Schiller-Pisarev test was used, as well as indices: papillary-marginal-alveolar index (I.A. Barannikova, 1990), periodontal index (I.S. Ivanov, 1998), communal periodontal index [21-24].

All the information obtained during the study was statistically processed in MS Excel and Statistica programs.

In addition, the PMA index, the Russell periodontal index, etc. are used to assess the condition of patients. The same group includes hygienic indices (Fedorov-Volodkina, Green-Vermillion, Ramfjord, etc.). Irreversible indices characterize the severity of such symptoms of periodontal diseases as bone resorption of the alveolar process, gum atrophy. It is used to assess gum inflammation. The index is considered the most convenient both for epidemiological study and for individual studies in clinical practice. The assessment of the condition of the gums is carried out in the area of

all teeth. The Schiller-Pisarev test is a lifetime staining of gum glycogen, the amount of which increases with inflammation. With the help of this test, the presence and prevalence of inflammation are determined.

3. RESULTS AND DISCUSSION

The results of the studies showed that the inclusion of hirudo therapy in the complex of standard methods of treatment was more effective compared to therapy when medical leeches were not used.

The appointment of hirudotherapy in patients of group I(a) was accompanied in almost all by an improvement in the general condition the very next day after the first session of application of a medical leech. On the second and third days, the bleeding has stopped and hyperemia in the inflammatory process significantly decreased, unpleasant sensations, burning sensation and pain in the gum area disappeared, bad breath disappeared. On day 7, in 91.1% of patients, the Schiller-Pisarev test confirmed the absence of manifestations of the inflammatory process. The absence of separable from the dentoalveolar pockets and a clear tendency to their sclerosis were noted. The pathological mobility of the teeth was no longer determined.

In patients of group I(b), with GCP and background hypertension, who underwent a complex of traditional standard therapy, the results obtained differed significantly from patients with hirudotherapy and from the results of treatment of patients of group II. Positive dynamics in the clinic of GCP in patients of group I(b) was revealed only on the 11th-12th day of treatment, while against the background of improvement in general condition, cessation of pain and bleeding gums, burning and itching in 57.5% of patients, 52.5% retained separate areas of inflammation, confirmed by the Schiller-Pisarev test. 40.0% retained serous discharge from periodontal pockets. Pathological mobility of teeth was not noted [21].

In patients of the second group with normal blood pressure, whose treatment was also carried out according to the traditional standard program, without the use of hirudotherapy, positive dynamics in clinical manifestations of the disease in 62.8% of patients was noted after 9-10 days. At the same time, patients noted an improvement in well-being, cessation of itching, burning, soreness and bleeding in the gum area, almost

complete disappearance of hyperemia and swelling of the mucous membrane. However, some representatives of the group – 37.1% in the gingival margin retained areas of inflammation, decreased, but not completely disappeared serous discharge from periodontal pockets. The Schiller-Pisarev test also remained positive in 28.6% of group II patients.

Thus, the relief of inflammatory phenomena in periodontal disease in most patients of group I(a) with background GB with the inclusion of hirudotherapy in the complex of standard treatment methods occurred much earlier, as indicated by the indicators of papillary-marginal-alveolar index, periodontal index, communal periodontal index (Figure 1), significantly different from the initial values already on 6-7 days after the start of therapeutic measures with the use of medical leeches.

A similar positive dynamics of laboratory parameters (the level of phosphorus in the blood serum increased by 0.04 mmol/l, and in the elderly - by 0.06 mmol/l. In younger and older patients of the comparison group, the level of phosphorus in the blood serum increased by 0.05 mmol/l), indicating a decrease in inflammatory phenomena in periodontal disease in patients of group I(b) with hypertension, who did not undergo hirudotherapy, was recorded only after 11-12 days. However, the indicators of the activity of the pathological process (bleeding) significantly exceeded similar values of patients of both group I(a) with hirudotherapy and the second group (patients with GCP, but with normal blood pressure), which indicated a significant contribution of hemodynamic disorders in background hypertension to the nature of blood supply to periodontal tissues, manifested by deterioration of metabolic processes and a more torpid course of the disease to therapy [25,26].

Evaluation of the effectiveness of various treatment options for patients with GCP was carried out in the catamnesis, 6 and 12 months after the course of treatment. When examining patients of group I(a) with background hypertension treated with hirudotherapy, complete weakening the disease (patients were under observation for 14 days) after 6 months was maintained in 43 patients (95.5%), after 12 months - in 35 (77.7%). Of the 40 patients in the Ib group whose complex of therapeutic measures did not include hirudotherapy, remission during GCP persisted in 19 patients (47.5%) after 6

months, and in 16 patients (40.0%) after 12 months. In the second group - with GCP, but normal blood pressure, after 6 months, indicators of inflammatory process activity were absent in 22 patients (62.8%), after 12 months, 19 patients (54.2%) remained in clinical remission.

According to the data obtained, complete clinical remission persisted after 6 and 12 months in the vast majority of patients of group Ia with GB with the use of hirudotherapy in complex treatment, whereas in representatives of group I(b), without hirudotherapy, remission was observed in less than half of the patients.

Based on the studies carried out, remission was revealed in the vast majority of patients of group I(a) with hypertension when using hirudotherapy as part of a complex treatment. Measurements of the pocket probing depth (0.3 mm), the level of clinical attachment and reliable estimates of the bleeding index (with a decrease in the Hb level by ≥ 5 g / dL or Ht by $\geq 15\%$) were carried out.

In the studies, the authors also cite that hypertension and periodontitis are among the

most common infectious diseases worldwide. Scientists have studied and systematized the available data on the relationship between hypertension and periodontitis and the potential underlying mechanisms. We demonstrate that hypertension and periodontitis are concomitant diseases, and their relationship remains significant even when considering other factors. Age and periodontitis activity change the relationship, and key mechanisms include inflammation and oxidative stress, leading to deterioration of vascular function and structure. On the other hand, hemodynamic changes associated with hypertension can lead to remodeling of periodontal bones and vessels, which supports local inflammation. Only a few studies evaluate the effect of periodontitis treatment on blood pressure (BP). The results confirm that the assessment of the periodontal condition and specific treatment (if necessary) can improve blood pressure control. Individual and population cardiovascular effects of treatment of periodontal pathology should be evaluated during a comprehensive cardiovascular examination [27].




Fig. 1. Dynamics of PI indicators in patients with GCP in the course of therapy

Thus, the results of the work indicate the high efficiency of the use of hirudotherapy in the complex of traditional methods of rehabilitation of patients with generalized chronic periodontitis and background hypertension. The relief of the chronic inflammatory process in periodontal was characterized by a significant improvement and had a more persistent and prolonged character.

4. CONCLUSION

The inclusion of hirudotherapy in the program of therapeutic measures in patients with generalized chronic periodontitis and background hypertension made it possible to optimize the onset of a persistent therapeutic effect, lengthen the period of remission of the pathological process in periodontitis and significantly improve the quality of life of patients. These results allow us to recommend the widespread use of the method in clinical dentistry.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

ACKNOWLEDGEMENT

The work was carried out using the Center for Collective Use of the North Caucasus Federal University

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Remizova AA, Dzgoeva MG, Tingaeva YI, Hubulov SA, Gutnov VM, Bitarov PA, et al. Tissue Dental Status and Features of Periodontal Microcirculation in Patients with New COVID-19 Coronavirus Infection.. *Pharmacophore*. 2021;12(2): 6-13. Available: <https://doi.org/10.51847/5JlbnUb HkT>.
2. Gurevich S, Prins KW. The evolving role of interventional cardiology in the treatment of pulmonary hypertension. *Catheter CardiovascInterv*. 2021;97(4):E446-E453. DOI: 10.1002/ccd.29146.
3. Dzgoeva MG, Tibilova FL, Dzgoeva ZG, Khetagurov SK, Cerekova AA. Features of microcirculation in periodontal tissues in patients with arterial hypertension during orthopedic treatment using implants. *Modern orthopedic dentistry*. Voronezh. 2018:128-130. (In Russian)
4. Flack JM, Adekola B. Blood pressure and the new ACC/AHA hypertension guidelines. *TrendsCardiovascMed*. 2020 ;30(3):160-164. DOI: 10.1016/j.tcm.2019.05.003.
5. Dmitry Anatolyevich Domenyuk, Vladimir AlexandrovichZelensky, Igor Vladimirovich Rzhepakovsky, Oksana IvanovnaAnfinogenova, Application of Laboratory and X-Ray Genral Studies un Early Diagnostics of Metabolic Disturbances of Bone Tissue in Children with Autoimmune Diabetes Mellitus , *EntomolApplSci Lett*. 2018; 5(4):1-12
6. Blinov AV, Siddiqui SA, Nagdalian AA, Blinova AA, Gvozdenko AA, Raffa VV, et al. Investigation of the influence of Zinc-containing compounds on the components of the colloidal phase of milk. *Arab J Chem*. 2021;14(7):103229.
7. Rzhepakovskiy, Siddiqui SA, Avanesyan S, Benlidayi M, Dhingra K, Dolgalev A, Enukashvily N, Fritsch T, Heinz V, Kochergin S, Nagdalian A, Sizonenko M, Timchenko L, Vukovic M, Piskov S, Grimm WD. Anti- arthritic effect of chicken embryo tissue hydrolyzate against adjuvant arthritis in rats (X-ray microtomographic and histopathological analysis). *Food Science & Nutrition* 2021;00:1-22. Avilable:<https://doi.org/10.1002/fsn3.2529>
8. Remizova A A, Sakaeva Z U, Dzgoeva Z G, Rayushkin I I, Tingaeva Y I, Povetkin S

- N, et al. The Role Of Oral Hygiene In The Effectiveness Of Prosthetics On Dental Implants. *Ann. Dent. Spec.* 2021;9(1): 39-46.
Available: <https://doi.org/10.51847/HuTuWdD0mB>
9. Kaloeva ZD, Dzilikhova KM, Dzgoeva ZG, Cerekova AA, Dzgoeva MG, Ataeva MV, Tuaeava N.O. Features of hemodynamic disorders in adolescents with primary arterial hypotension. *G.N.Speransky Pediatrics.* 2017; 96(5):236-238 (In Russian).
 10. Orlandi M, Graziani F, D'Aiuto F. Periodontal therapy and cardiovascular risk. *Periodontol 2000.* 2020;83(1):107-124.
DOI: 10.1111/prd.12299.
 11. Avolio A, Kim MO, Adji A, Gangoda S, Avadhanam B, Tan I, Butlin M. Cerebral Haemodynamics: Effects of Systemic Arterial Pulsatile Function and Hypertension. *CurrHypertens Rep.* 2018 Mar 19;20(3):20.
DOI: 10.1007/s11906-018-0822-x.
 12. Valensise H, Farsetti D, Pisani I, Tiralongo GM, Lo Presti D, Gagliardi G, Vasapollo B, Novelli GP. Friendly help for clinical use of maternal hemodynamics. *J Matern Fetal Neonatal Med.* 2021;34(18):3075-3079.
DOI: 10.1080/14767058.2019.1678136.
 13. Galabueva AI, Biragova AK, Kotsoyeva GA, Borukayeva ZK, Yesiev RK, Dzgoeva ZG, et al. Optimization of Modern Methods of Treating Chronic Generalized Periodontitis of Mild Severity. *Pharmacophore.* 2020;11(1):47-51
 14. Yusupova MI, Mantikova KA, Kodzokova MA, Mishvelov AE, Paschenko AI, Ashurova ZAK, et al. Study Of The Possibilities Of Using Augmented Reality In Dentistry. *Ann. Dent. Spec.* 2021;9(2):17-21.
Available: <https://doi.org/10.51847/BG1ZAzqXRc>.
 15. Malezhik LP, Pinelislul, Malezhik MS. Some aspects of immune reaction in chronic generalized parodontal disease in elderly patients. *Stomatologiya (Moscow).* 2011;90(6):8-10 (In Russian).
 16. Seshima F, Nishina M, Namba T, Saito A. Periodontal Regenerative Therapy in Patient with Chronic Periodontitis and Type 2 Diabetes Mellitus: A Case Report. *BullTokyoDentColl.* 2016;57(2):97-104.
DOI: 10.2209/tdcpublication.2015-0041.
 17. Ali Redha, Ali & Siddiqui, Shahida & Ibrahim, Salam. Advanced extraction techniques for Berberis species phytochemicals: A review. *International Journal of FoodScience & Technology.* 2021:1-12, doi:10.1111/ijfs.15315.
 18. Siddiqui SA, Blinov AV, Serov AV, Gvozdenco AA, Kravtsov AA, Nagdalian AA, Raffa VV, Maglakelidze DG, Blinova AA, Kobina AV, et al. Effect of Selenium Nanoparticles on Germination of *HordéumVulgáre* Barley Seeds. *Coatings* 2021;11:862.
Available: <https://doi.org/10.3390/coatings11070862>
 19. Moghadam ET, Yazdanian M, Tahmasebi E, Tebyanian H, Ranjbar R, Yazdanian A, Seifalian A, Tafazoli A. Current herbal medicine as an alternative treatment in dentistry: In vitro, in vivo and clinical studies. *Eur J Pharmacol.* 2020;889: 173665.
DOI: 10.1016/j.ejphar.2020.173665.
 20. Leiva-Cala C, Lorenzo-Pouso AI, Centenera-Centenera B, López-Palafox J, Gándara-Vila P, García-García A, Pérez-Sayáns M. Clinical efficacy of an Aloe Vera gel versus a 0.12% chlorhexidine gel in preventing traumatic ulcers in patients with fixed orthodontic appliances: a double-blind randomized clinical trial. *Odontology.* 2020;108(3): 470-478.
DOI: 10.1007/s10266-019-00468-w.
 21. Jha K, Garg A, Narang R, Das S. Hirudotherapy in Medicine and Dentistry. *J ClinDiagn Res.* 2015;9(12):ZE05-7.
DOI: 10.7860/JCDR/2015/16670.6918.
 22. Gödekmerdan A, Arusan S, Bayar B, Sağlam N. Tibbisülüklervehirudoterapi (Medicinal leeches and hirudotherapy). *Türkiye Parazitol Derg.* 2011;35(4):234-9. Turkish.
DOI: 10.5152/tpd.2011.60.
 23. Guts luV. Rol' stepenike ratini zatsiislizistoï obolochki polostirta v otsenkere zull'tatovpro by Shillera—Pisareva (The role of the degree of oral mucosal keratinization in assessing the results of the Schiller-Pisarev test). *Stomatologiya (Moscow).* 1989;68(2):72-3. (In Russian).
 24. Nazaryan R, Kryvenko L, Zakut Y, Karnaukh O, Gargin V. Application of estimated oral health indices in adolescents with tobacco addiction. *PoIMerkurLekarski.* 2020;48(287): 327-330.
 25. Gillette WB. Periodontal index. *J Am Dent*

- Assoc. 1987;114(5):580-582.
DOI: 10.14219/jada.archive.1987.0140.
26. Benigeri M, Brodeur JM, Payette M, Charbonneau A, Ismaïl AI. Community periodontal index of treatment needs and prevalence of periodontal conditions. J ClinPeriodontol. 2000;27(5):308-12.
DOI:10.1034/j.1600-051x.2000.027005308.x.
27. Pogodina AV, Valyavskaya OV, Kolesnikova LR, Rychkova LV, Kolesnikov SI. Arterial hypertension and periodontitis: key aspects of comorbidity. Arterial'naya Gipertenziya ("Arterial Hypertension"). (In Russ.) 2019;25(6): 682-692.
Available: <https://doi.org/10.18705/1607-419X-2019-25-6-682-692>

© 2021 Serekova et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/76378>