



Effect of GA3 Plus Yogen Foliar Fertilization on Yield and Fruit Quality of Red Dragon Fruit at Lap Thanh District in Vietnam

Nguyen Minh Tuan^{1*}, Nguyen The Huan¹ and Nguyen Quoc Hung²

¹Department of Agronomy, Thai Nguyen University of Agriculture and Forestry, Quyet, Thang Commune, Thai Nguyen City, Vietnam.

²Fruit and Vegetable Research Institute, Ha Noi, Vietnam.

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2016/25129

Editor(s):

(1) Scd Victoria Anatolyivna Tsygankova, Department for Chemistry of Bioactive Nitrogen-Containing Heterocyclic Compounds, Institute of Bio-organic Chemistry and Petrochemistry, Plant Physiology, NAS, Ukraine.

Reviewers:

(1) Abdur Rab, KP Agricultural University, 25130, Peshawar, Pakistan.

(2) Isaac Kojo Arah, Ho Polytechnic, Ghana.

Complete Peer review History: <http://www.sciencedomain.org/review-history/15559>

Original Research Article

Received 19th February 2016

Accepted 28th March 2016

Published 29th July 2016

ABSTRACT

The present study aimed to evaluate the effect of GA₃ plus Yogen foliar fertilization application on fruit yield and quality of TL4 red dragon fruit from March to August 2015 at Lap Thach district, northern Vietnam. The experiment was conducted on complete block design with three replications. Fruit yield and different fruit quality parameters, as well as grade of fruit were determined. The results indicated that application of GA₃ 30 ppm plus Yogen 4 foliar fertilization significantly increased fruit set, fruit weight, fruit yield as well as fruit quality and reduced percentage grade S for red dragon fruit. Therefore, it could be concluded that application of GA₃ 30 ppm plus Yogen 4 foliar fertilization markedly fruit set, fruit quality for TL4 red dragon fruit.

Keywords: Red dragon fruit; GA₃; yogen foliar fertilization; fruit yield and quality.

*Corresponding author: E-mail: tuan_dhnl@yahoo.com;

1. INTRODUCTION

The red dragon fruit (*Hylocereus* spp.) is also known as pitaya or pitahaya (Latin America), strawberry pear, thang loy (Vietnamese), pitaya roja (Spanish), and la pitahaya rouge (French) and night blooming cereus (English) [1,2]. Dragon fruit is the fruit of several cactus species that have been classified as white (*Hylocereus undatus*), red (*H. polyrhizus*), and yellow (*Selenicereus megalanthus*) [3,4] and are native to Americas [5], with a wide distribution in the tropics and sub-tropics [6]. However, only the red and white flesh varieties are popular in VietNam. The dragon fruit is a large, oblong fruit with a red peel and large green scales. The flesh is sweet, delicate, red-purple, and contains numerous tiny black seeds. The fruit is nonclimacteric and has the best flavor when harvested at full red color [7]. Dragon fruit is rich in vitamins, minerals and fibers that helps the digestive process, prevent colon cancer and diabetes, neutralize toxic substances such as heavy metals, and helps to reduce cholesterol levels and high blood pressure [8]. Dragon fruit is enjoyed as a fresh fruit or juice, but also is valued as a natural food colorant [9]. Currently, dragon fruit is being grown commercially in Australia, Brazil, Colombia, Costa Rica, Egypt, Israel, Japan, Mauritius, Mexico, Nicaragua, Taiwan, the USA and Vietnam [6]. Exceptional tolerance to extreme drought, its micronutrient content and the vibrant color of the fruit itself has attracted consumer in the national and international markets. Its production could potentially create job and promote income for the nation that produces it [10,11,12].

Plant growth regulators (PGRs) have become an important component of agrotechnical procedures for most of the cultivated plants and especially for fruit plants [13]. According to reference [14] growth regulators may provide the means of bringing about required growth responses as their use can increase the yield of product. The use of plant growth regulators compounds (auxins, cytokinins and gibberellins) is becoming popular to ensure efficient production [15]. Moreover, many previous studies have shown that the application of PGRs can enhance the rapid changes in physiological and biochemical characters thus increasing crop productivity. GA3 (gibberellic acid) had been shown fruit development, reduce fruit drop, fruit cracks and improved fruit quality of wax apple under field conditions [16]. Reference [17]

indicated that among agricultural practices which may increase the fruit production and improve the quality of several fruit crops gibberellic acid. Gibberellic acid has been reported to influence vegetative growth, flowering, fruiting and various disorders in many fruit crops [18]. Sprays of GA3 have been widely adopted in commercial orchards because they have consistently been shown to increase fruit size and firmness of cherry [19]. Moreover, GA3 increased the yield of fruit in Balady mandarin [20], and increases soluble solids as well as fruit weight in sweet cherry [21]. Therefore, in order to increase the fruit production, plant growth regulators are one of the production tools that can enhance product quality and marketability.

Moreover, the plant nutrition is one of the most important factors responsible for the proper growth and development of the plants. The methods of nutrient application play an important role in supplying the nutrients to the plants because the efficacy of fertilizers applied soil low due to various losses and fixations. Foliar nutrition is designed to eliminate the above problems particularly with respect to macro nutrients. Nowadays application of N, P and K application in different ratios through foliar sprays is modern method of fertilization in vegetable crops due to the nature of heavy feeder of nutrients [22]. Foliar nutrition is more effective on young leaves, and shortage of macro and micro nutrients can be removed by this factor [23]. Rapid uptake of nutrients and no influence of pH and soil texture as well as providing cations such as Zn and Fe in the soil for plants that stabilize these elements and finally being cheaper than other methods are the advantages of foliar nutrition method [24]. Reference [25] studied the effects of folia application using nitrate and the results revealed that the yield of cucumber raised up to 14.5 tons per ha and there were an increase of NPK in plants applying foliar application.

Red dragon fruit cultivation has been recently introduced and fruit consumption has gained popularity in south VietNam. Although references are available in the literature and efforts have been made to improve fruit set as well as fruit quality by applying of chemical compounds but there is no available literature on the effect of plant growth regulator and foliar fertilizer on physiological and biochemical parameters of red dragon fruit in the North Vietnam. Therefore, this study was carried to investigate the effects of GA3 plus Yogen foliar application on growth,

yield and quality of red dragon fruit under field conditions.

2. MATERIALS AND METHODS

2.1 Plant Materials and Experiment Treatments

The experiment was conducted at red dragon fruit orchard in Lap Thach district, Vinh Phuc province, Northern VietNam from March to August 2015. The TL4 pitaya cultivar with the red peeled pulp was chosen for the experiment. The experiment was design in Randomized Complete Block Design (RCBD) with three replicated and three uniform trees were taken as an experiment unit. The experiment consisted of five treatments were applied for foliar application: Treatment 1 (GA₃30 ppm + Yogen1 NPK: 30:10:10); Treatment 2 (GA₃30 ppm + Yogen2 NPK: 10:30:10); Treatment 3 (GA₃30 ppm + Yogen3 NPK: 10:10:30); Treatment 4 (GA₃30 ppm + Yogen4 NPK compounds of Yogen1 + Yogen2 + Yogen 3); Treatment 5 (Control treatment (water spray). GA3 and Yogen were applied after fruit set stage on windless mornings with a truck-mounted monitorized sprayed until drip of and subsequently in 7 day intervals as 3 times.

2.2 Data Collection

The observations with regards to the growth, yield and quality was recorded from the randomly selected and tagged plants. The percentage of fruit setting was calculated using the following formula:

$$\text{Fruit set (\%)} = \frac{\text{Number of fruit}}{\text{Number of flowers}} \times 100.$$

At harvesting, final fruit length, fruit diameter, peel thickness was determined with the help of Vernier caliper. The fruit diameter measurement recorded was the average of two readings taken at two axes of the midsection of the fruit. Fruit length was measured from the part attached to the petiole to the base of the fruit. Peel thickness was determined at the equatorial point of fruit. Yield per treatment was recorded by weighing and counting the total number of fruits. Fruit fresh weight (FW) were measured using anelectronic balance (GF-6100; A&D Co. Ltd., Tokyo,Japan), and the edible portion of each fruit was calculated as:

$$\text{Edible percentage} = \left(\frac{\text{Pulp weight}}{\text{Fruit weight}} \right) \times 100.$$

TSS content was measured using a hand refractometer (model PAL-1, Atago, Tokyo, Japan). Fruit flesh was squeezed from a sample of the middle of freshly cut fruit and the result is expressed as °Brix. The fruits were peeled prior to determining the nutritional properties. Total soluble sugar was determined according to the phenol-sulphuric method by [26], vitamin C were done following standard procedures [27].

2.3 Statistical Analysis

The data obtained from the study were analyzed using SAS 9.1 statistical software for each cultivar separately. Differences between reatments were measured using Duncan's multiplerange test at a significance level of $P \leq 0.05$.

3. RESULTS AND DISCUSSION

3.1 Effect of GA3 and Yogen Foliar Fertilization on Fruit Set and Fruit Yield

The percentage fruit set was found to be statistically significant between the different ($P \leq 0.05$) treatments and control treatment, except GA₃30 ppm + Yogen1 application. From the results, it was observed that the highest (72.56%) fruit set was recorded in spraying with GA₃30 ppm + Yogen4, followed by GA₃30 ppm + Yogen3, GA₃30 ppm +Yogen2, GA₃30 ppm + Yogen1, with value of 65.91 %; 64.95%; 62.20% fruit set, respectively while, the lowest percentage fruit set (60.13%) was recorded in untreated control (Table 1). It seems be that external application of gibberellin induces fruit set in several species. In the case study, best result for increasing fruit set was achieved with spraying GA₃30 ppm + Yogen4 foliar fertilization combination. These results are in agreement with the findings of [28] who indicated that gibberellic acid is used widely in horticultural crops for improving fruit set. Moreover, enhance fruit set of dragon fruit in the study might be a reason of supplying more nutrients at the critical fruit set stage, which is in accordance with the finding reported by [29,30].

3.2 Fruit Yield

As shown in Table 1, there was significantly increased fruit yield in all treatment as compared to untreated control. In the case of this study, the highest fruit yield was found in GA₃30 ppm + Yogen4 treatment with 15.00 kg/tree, followed by GA₃30 ppm + Yogen3, GA₃30 ppm + Yogen1 application, whereas the control treatment showed the lowest value of 8,13 kg/tree. This implies that chemical compounds might be effective in improving fruit yield. In the current study, fruit yield was markedly enhanced by application GA₃30 ppm + Yogen4 in comparison with control. Our results were found to be in agreement with that of [31] who observed that, GA3 significantly influenced the fruit weight as well as yield in mandarin. Furthermore, the increase in the yield might be due to greater availability of nutrients, increased uptake of nutrients and water, resulting in more photosynthesis and enhanced food accumulation in edible part of the dragon fruits in the study (Table 1). The references [32,33] also reported similarly. However, application of GA30 ppm + Yogen2 showed slightly increased fruit set as compared to untreated control, with the significantly difference at ($p \leq 0.05$) (Table 1).

3.2.1 Effect of GA3 and Yogen foliar fertilization on fruit parameter

It has been documented in previous studies that growth regulators can have a significant effect on fruit size and fruit quality. In this study GA3 plus Yogen treatment was found to have a positive effect parameter of red dragon fruits as show bellow:

3.3 Fruit Weight

Data in Table 2 indicated that, there was significantly affected by application of GA3 + Yogen foliar fertilization on fruit weight among treatments as compared with control treatment, in term the control treatment had the lowest of fruit weight (406.3 g), whereas the highest fruit weight (534.6 g) were recorded with GA₃30 ppm + Yogen4 treatment, followed by GA₃30 ppm + Yogen3 application, GA₃30ppm + Yogen1 treatment in this study with value of 516.4 g and 488.2 g, respectively (Table 2). It seems that the role of GA3 was to multiply and to lengthen the meristem cells, which resulted in the increase of fruit weight. In addition, spraying with GA₃30 ppm + Yogen4 foliar fertilization combination resulted in significantly fruit weight enhancement as compared to untreated control (Table 2).

Table 1. Effect of GA3 and Yogen foliar fertilization on fruit set and yield of red dragon fruit

Treatment	Fruit set (%)	Yield (kg/tree)
GA ₃ 30 ppm + Yogen1 ^y	62.20c ^x	10.67c
GA ₃ 30 ppm + Yogen2	64.95b	9.30d
GA ₃ 30 ppm + Yogen3	65.91b	12.47b
GA ₃ 30 ppm + Yogen4	72.56a	15.00a
Control treatment (water spray)	60.13c	8.13e

^ymean foliar ratio with Yogen 1 NPK (30:10:10); Yogen 2 NPK (10:30:10); Yogen 3 NPK (10:10:30); Yogen 4 (Yogen 1 + Yogen 2 + Yogen 3 compounds)

^x mean in each column followed by the same letters are not significantly different at $P \leq 0.05$ according to Duncan's multiple range test

Table 2. Effect of GA3 and Yogen foliar fertilization on fruit parameter of red dragon fruit

Treatment	Fruit weight (g)	Fruit length (cm)	Fruit diameter (cm)	Peel thickness (cm)	Edible percentage (%)
GA ₃ 30 ppm + Yogen1 ^y	488.2ab ^x	11.37bc	8.61b	0.45a	82.52bc
GA ₃ 30 ppm + Yogen2	443.7bc	11.19bc	8.58b	0.46a	82.56bc
GA ₃ 30 ppm + Yogen3	516.4a	12.55b	8.73b	0.46a	83.61ab
GA ₃ 30 ppm + Yogen4	534.6a	14.22a	9.85a	0.45a	84.09a
Control treatment (water spray)	406.3c	10.45c	8.04b	0.43a	81.72c

^ymean foliar ratio with Yogen 1 NPK (30:10:10); Yogen 2 NPK (10:30:10); Yogen 3 NPK (10:10:30); Yogen 4 (Yogen 1 + Yogen 2 + Yogen 3 compounds)

^x mean in each column followed by the same letters are not significantly different at $P \leq 0.05$ according to Duncan's multiple range test

This is in agreement with [16] who found that spraying wax apple trees with GA3 significantly increased fruit weight. However, application of GA₃30 ppm + Yogen2 slightly increase fruit weight as compared untreated control although there was not significantly different at ($p \leq 0.05$) (Table 2). This results are in agreement with the report of [34], who indicated that the role of GA in improving fruit quantity namely, fruit weight and fruit size may be due to its role in increasing cell elongation.

3.4 Fruit Length and Fruit Diameter

The results summarized in Table 2 showed that, the application of different chemical compounds had significant effects on fruit length, in which the highest fruit length of 14.22 cm was recorded in GA₃30 ppm + Yogen4 treatment, followed by spraying GA₃30 ppm + Yogen3 with value of 12.55 cm. It consider that the increase in fruit size might be attributed to increase in cell division and cell elongation caused by auxins and GA3 which is in accordance with the finding reported by [35] who demonstrated that the increase in fruit size may be attributed to the increase in cell division and cell elongation caused by GA3. However, there was no effect of spraying with GA₃30 ppm + Yogen2, GA₃30 ppm + Yogen1 on fruit length than the untreated control (Table 2).

Furthermore, reference [36] indicated that the fruit size is one of the most important quality parameter in sweet cherry. For this reason, as the big fruits are much more fleshier, they are preferred more by the consumers. In the case of the study, there was significant differences fruit diameter between treatments (Table 2). In term, the treatment with GA₃30 ppm + Yogen4 application produced the highest fruit diameter (9.85 cm), whereas the lowest fruit diameter (8.04 cm) was recorded for untreated control (Table 2, Fig. 1). It seems that spray GA₃30 ppm + Yogen4 markedly increased fruit diameter

compared with the control. It consider that fruit size were significantly different due to the interaction effect of dragon fruit with GA3, indicating differential response in increasing size of red dragon fruit to GA3 application (Table 2). These results are in agreement with [37] who found that GA3 application increased cell division and elongation and had a positive effect on fruit size. However, application of GA₃30 ppm + Yogen3, GA₃30 ppm + Yogen2, GA₃30 ppm + Yogen1 slightly improve fruit diameter than the control treatment although there was not significantly different at ($p \leq 0.05$) (Table 2).

3.5 Peel Thickness of Fruit

The peel thickness for all treatment in this study is presented in Table 2. In term, spraying with GA₃30 ppm + Yogen3 gave the highest Peel thickness with values of 0,46 cm, while the control treatment produced the lowest Peel thickness (0,43 cm). The same was also observed concerning the peel thickness of remained treatment. In which, spraying with GA₃30 ppm + Yogen2, GA₃30 ppm + Yogen4, and GA₃30 ppm + Yogen1 had higher peel thickness (0,46 cm, 0,45 cm and 0,45 cm, respectively) in comparison with control, although the difference was not statistically significant ($p \leq 0.05$) (Table 2).

3.6 Edible Percentage of Fruit

The results summarize in Table 2 showed that there was greatly effect of chemical compounds application on percentage of edible among treatment. Spraying with GA₃30 ppm + Yogen4 had the maximum edible percentage (84.09%), followed by GA₃30 ppm + Yogen3 applications with value of 83.61% edible percentage. However, application of GA₃30 ppm + Yogen2 and GA₃30 ppm + Yogen1 showed slightly increased edible percentage as compared to untreated control, although there were not significantly different.




Fig. 1. Fruit size of red dragon TL4 cultivar (A: untreated control; B: GA₃30 ppm + Yogen4)

3.6.1 Effect of GA3 and Yogen foliar fertilization on fruit quality

TSS are considered an important quality parameter of any fruit. It has been reported that, chemical compounds can change the TSS content of fruits. As shown in Table 3, there was no significant difference TSS among treatment ($p \leq 0.05$), in terms of the GA₃30 ppm + Yogen4 application produced the highest TSS (14.19⁰Brix), followed by GA₃30 ppm + Yogen3 sprayed and the other treatments. In the same table data showed that application of chemical compounds also significantly increase vitamin C as compared to untreated control, which was recorded in GA₃30 ppm + Yogen4 application with value of 8.79 mg/100 g. It seem be that the increase in quality character might be due to the growth promoting substances which could have accelerated synthesis of carbohydrate, vitamins and other quality characters. These results are in conformity with those of [38,33]. However, spraying with GA₃30 ppm + Yogen3, GA₃30 ppm + Yogen2, GA₃30 ppm + Yogen1 showed slightly enhance vitamin C than the control treatment

with values of 7.95 mg/100 g, 7.93 mg/100 g, and 7.94 mg/100 g although there were not significantly different. It is well known now that plant growth regulators can play a role in increase in the sugar content in fruits. Data presented in Table 3 indicated that there were various responses among treatments with GA₃ + Yogen compounds at different concentration, although the difference was not statistically significant ($p \leq 0.05$).

Fruit are generally graded by size and color. The result of the study shows that there were significantly different graded of dragon fruit among treatment at different chemical compounds (Table 4). In term, the highest value of 5.95% and 52.38% was recorded with GA₃30 ppm + Yogen4 application compared to lowest value of 3.35% and 37.48% was found in untreated control in the case of grade XL and L, respectively. However, the lowest value of 8.10% was obtained in GA₃30 ppm + Yogen4, but highest value of (23.49%) was recorded in control treatment, which was achieved in the case of grade S (Table 4). It seem be that

Table 3. Effect of GA3 and Yogen foliar fertilization on fruit quality of red dragon fruit

Treatment	TSS ⁰ Brix	Vitamin C (mg/100 g)	Sugar content (%)
GA ₃ 30 ppm + Yogen1 ^y	13.26a ^x	7.94ab	10.88a
GA ₃ 30 ppm + Yogen2	13.60a	7.93ab	10.76a
GA ₃ 30 ppm + Yogen3	13.39a	7.95ab	10.89a
GA ₃ 30 ppm + Yogen4	14.19a	8.79a	11.08a
Control treatment (water spray)	13.01a	6.90b	10.91a

^y mean foliar ratio with Yogen 1 NPK (30:10:10); Yogen 2 NPK (10:30:10); Yogen 3 NPK (10:10:30); Yogen 4 (Yogen 1 + Yogen 2 + Yogen 3 compounds)

^x mean in each column followed by the same letters are not significantly different at $P \leq 0.05$ according to Duncan's multiple range test

Table 4. Effect of GA3 + Yogen on percentage of grading standard for dragon fruit

Treatment	XL ^z (%)	L (%)	M (%)	S (%)
GA ₃ 30 ppm + Yogen1 ^y	5.21ab ^z	48.40ab	33.6a	12.79c
GA ₃ 30 ppm + Yogen2	4.93b	48.33ab	32.40a	14.34c
GA ₃ 30 ppm + Yogen3	5.88a	44.41b	33.24a	16.47b
GA ₃ 30 ppm + Yogen4	5.95a	52.38a	33.57a	8.10d
Control treatment (water spray)	3.35c	37.48c	35.68a	23.49a

^y mean foliar ratio with Yogen 1 NPK (30:10:10); Yogen 2 NPK (10:30:10); Yogen 3 NPK (10:10:30); Yogen 4 (Yogen 1 + Yogen 2 + Yogen 3 compounds)

^x mean in each column followed by the same letters are not significantly different at $P \leq 0.05$ according to Duncan's multiple range test

^z grade 1 (XL – Extra Large fruit): above 700 g; grade 2 (L-large fruit): 500 to 700 g; grade 3 (M-regular fruit 330 to 500g); grade 4 (S-small fruit: 200 to 330 g) modify size grades suggested for Vietnam of [39]

gibberellic acid (GA3) plays a very important role in enlarging fruit size of red dragon fruit. This results are in agreement with [40] who indicated that strengthen the role of gibberellic acid (GA3) in the agricultural sector where previously it had been well known to be used to increase fruit set and fruit size of many plants and fruits: grapes (*Vitis vinifera*), lemon (*Citrus spp.*), banana (*Musa spp.*), currant (*Ribes aureum*), pineapple (*Ananas comosus*) and sweet cherry (*Prunus avium*). In the current study grade L gave the highest value of (52,38%) as compared to other grades (Table 4).

4. CONCLUSIONS

From the above mentioned results it can be concluded that the GA₃30 ppm + Yogen4 foliar fertilization compounds application greatly increased the percentage of fruit set. Moreover, application of GA₃30 ppm + Yogen4 foliar fertilization gave the best results in the physiological and biochemical parameters of red dragon fruit with improved fruit size, fruit weight, enhanced total soluble solids, total sugar, and vitamin C as well as increased fruit yield. Hence, application of GA₃30 ppm + Yogen 4 foliar fertilization could be a valuable tool in improving quality of red dragon fruit, based on both physical and biochemical quality characteristics.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Mizrahi Y, Nerd A, Nobel PS. Cacti as crops. Horticultural Reviews. 1997;18:291-319.
- Paull RE. Dragon fruit. In: Gross KC, Wang CY, Salveit M, (eds.). The commercial storage of fruits, vegetables and florist and nursery stocks. 2004. USDA, ARS Agricultural Handbook #66; 2008. Available<<http://usna.usda.gov/hb66/contents.html>>
- Hoa TT, Clark CJ, Waddell BC, Woolf AB. Postharvest quality of dragon fruit (*Hylocereus undatus*) following disinfecting hot air treatments. Postharvest Bio Tech. 2006;41:62–9.
- Nerd A, Sitrita Y, Kaushika RA, Mizrahi Y. High summer temperatures inhibit flowering in vine pitaya crops (*Hylocereus spp.*). Scientia Horticulturae. 2002;96:343–50.
- Barthlott W, Hunt DR. Cactaceae. In: Kubitzki K, editor. The families and the genera of vascular plants. Berlin (Germany): Springer. 1993;2:161–196.
- Merten S. A review of *Hylocereus* production in the United States. J. Prof. Assoc. Cactus Developmt. 2003;5:98–105.
- Nerd A, Gutman F, Mizrahi Y. Ripening and postharvest behaviour of fruits of two *Hylocereus* species (Cactaceae). Postharv. Biol. Technol. 1999;17:39-45.
- Zainoldin KH, Baba AS. The effect of *Hylocereus polyrhizus* and *Hylocereus undatus* on physicochemical, proteolysis, and antioxidant activity in yogurt. World Academy of Science, Engineering and Technol. 2009;60.
- Wybraniec S, Mizrahi Y. Fruit flesh betacyanin pigments in *Hylocereus cacti*. J. Agr. Food Chem. 2002;50:6086–6089.
- Cos P, Bruyne DT, Hermans N, Apers S, Berghe DV, Vlietinck AJ. Proanthocyanidins in health care: Current and new trends. Curr Med Chem. 2004;11: 1345–59.
- Stintzing FC, Schieber A, Carle R. Betacyanins in fruits from red-purple pitaya, *Hylocereus polyrhizus* (Weber) Britton & Rose. Food Chem. 2002;77:101–6.
- Yolanda Donají Ortiz-Hernández, José Alfredo Carrillo-Salazar. Pitahaya (*Hylocereus spp.*): A short review. Comunicata Scientiae. 2012;3(4):220-237.
- Monselise SP. The use of growth regulators in citriculture. A Review Sci. Hort. 1979;11:151-162.
- Morgan P. Synthetic growth regulators: Potential for development. Botanical Gazette. 1980;141:337–346.
- Guardiola JL, Garcia-Luis A. Increasing fruit size in citrus. Thinning and stimulation of fruit growth. Plant Growth Regul. 2000;31:121–132.
- Nguyen MT, Yen CR. Effect of Gibberellic Acid and 2,4-Dichlorophenoxyacetic Acid on fruit development and fruit quality of wax apple. World Academy of Science,

- Engineering and Technology. 2013;7:302-308.
17. Chao CT, Lovatt CJ. Foliar applied 3,5,6-trichloro-2-pyridoxylacetic acid (3,5,6-TPA) increases yield of commercially valuable large-size fruit of "Fina Sodea" clementine mandarin. *Acta Hort.* 2010;884:433-440.
 18. Paroussi G, Voyiatzis DG, Paroussi E, Drogour PD. Growth, flowering and yield responses to GA3 of strawberry grown under different environmental conditions. *Scientia Hort.* 2002;96:103–113.
 19. Clayton M, Biasi WV, Agar IT, Southwick SM, and Mitcham EJ. Sensory quality of 'Bing' sweet cherries following preharvest treatment with hydrogen cyanamide, calcium ammonium nitrate, or gibberellic acid. *HortScience.* 2006;41:745–748.
 20. El-Sese AMA. Effect of gibberellic acid 3 (GAs) on yield and fruit characteristics of Balady mandarin. *Assiut. J. Agri. Sci.* 2005;36:23-35.
 21. Basak A, Rozpara E, Grzyb Z. Use of bioregulators to reduce sweet cherry tree growth and to improve fruit quality. *Acta Hort.* 1998;468:719-723.
 22. Chaurasia SNS, Singh KP, Mathura R. Effect of foliar application of water soluble fertilizers on growth, yield, and quality of tomato (*Lycopersicon esculentum L.*). *Sri Lankan J. Agric. Sci.* 2005;42:66-70.
 23. Kashi AK. *Vegetables. Textbook of Horticulture College, Tehran University; 1994.*
 24. Lanauskas J, Kvikliene N. Effect of calcium foliar application on some fruit quality characteristics of Sinap Orlovskij apple. *Agronomy Research.* 2006;4(1):31-36.
 25. Karuppaiah P, Manivannank K, Sriramach, Rasekaran MV, Kuppusamy G. Responses of cucumber to foliar application of nutrients on lignite mine spoil. *Journal of the Indian Society of Soil Science.* 2001;49.
 26. Dubois MK, Gils JK, Hanniton PA, Robes, Smith F. Use of phenol reagent for the determination of total sugar. *Anl. Chem.* 1956;28:350.
 27. Sattar A. Effect of foliar application of urea as supplement on the fruit drop and quality of Kinnow Mandarin (*Citrus reticulata Blanco*). MSc. Thesis, Inst. of Hort. Sci., Univ. of Agri. Faisalabad, Pakistan; 1999.
 28. Taylor DR, Knight JN. Russetting and cracking of apple fruit and their control with plant growth regulators. *Acta. Hort.* 1986; 11:819-820.
 29. Naik LB, Prabhakar M, Tiwari RB. Influence of foliar sprays with water soluble fertilizers on yield and quality of Carrot (*Daucus carota L.*). *Proc, Int. Conf. Vegetables, Bangalore.* 2002;183.
 30. Vibhute CP. A process for manufacturing complex solid and liquid completely water soluble fertilizers. *Fert. News.* 1988;43(8): 63.
 31. Saraswathi T, Rangasamy P, Azhakiamaavalan RS. Effect of preharvest spray of growth regulators on fruit retention and quality of mandarins (*Citrus reticulata Balanco*). *South Ind. Hort.* 2003;51:110-112.
 32. Guvence I, Badem H. Effect of foliar application of different sources and levels of nitrogen on growth and yield of tomato (*Lycopersicon esculentum L.*). *Indian J. Agric. Sci.* 2000;72(2):104-105.
 33. Singh AB, Singh SS. Effect of various levels of nitrogen and spacing on growth, yield and quality of tomato. *Veg. Sci.* 1992;19(1):1-6.
 34. Eman AA, El-moneim MMMA, El- Migeed OA, Ismail MM. GA3 and zinc sprays for improving yield and fruit quality of Washington navel orange trees grown under sandy soil conditions. *Res. J. Agric. Biol. Sci.* 2007;3(5):498-503.
 35. Ranjan R, Purohit SS, Prasad V. *Plant hormones: Action and application.* Agrobios, India. 2003;183–189.
 36. Horvitz S, Godoy C, Lopez Camelo AF, Yommi A. Application of gibberellic acid to 'Sweetheart' sweet cherries: Effect on fruit quality at harvest and during cold storage. *Acta Horticult.* 2003;628:311-316.
 37. Usenik V, Kastelec D, Stampar F. Physicochemical changes of sweet cherry fruits related to application of gibberellic acid. *Food Chem.* 2005;90:663-671.
 38. Fageria MS, Arya PS, Jagmohan K, Singh AK. Effect of nitrogen levels on growth yield and quality of tomato. *Veg. Sci.* 1992;19(1):25-29.
 39. Le VT, Nguyen N, Nguyen DD, Dang KT, Nguyen TNC, Dang MVH, Chau NH, Trink

- NL. Quality assurance system for dragon fruit. ACIAR Proceedings. 2000;100:101-114.
40. Nor Shariah S, Razifah MR, Mamat AS, Adzemi MA. Application of gibberellic acid (Ga3) in stem cutting of dragon fruit (*Hylocereus Polyrhizus*): Effects on fruit quality and yield at harvest. Journal of Biology, Agriculture and Healthcare. 2014;4(21):51-55.

© 2016 Tuan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/15559>